

Composition and vulnerability of mixed bird flocks in tropical biodiversity hotspots

Everton Sousa Ferreira¹, Caio Graco Machado² and Alexandre Schiavetti^{1,3}

ABSTRACT

The extinction of ecological interactions represents the most dramatic scenario resulting from the massive loss of species. Essentially, mixed-species flocks constitute critical mutualistic relationships for the structure, cohesion, and diversity in bird communities, yet they are strongly threatened by the extensive destruction of primary forests. In tropical regions and their biodiversity hotspots, there appears to be no integrated assessment of mixed-species flocks in terms of composition, vulnerability, or habitat use. Thus, a systematic review was conducted based on 269 studies on mixed-species flocks, including 55 conducted in the tropics from 1963 to 2022. We highlight 367 bird species most frequently found in mixed-species flocks, among which 13 are threatened and 14 are near-threatened. According to global assessments, the Tropical Andes harbor the highest number of threatened species, followed by the Coastal Forest of East Africa, Western Ghats and Sri Lanka, and at a third level, the Himalayas. The Atlantic Forest did not present any threatened species, although it constitutes the focus of discussions concerning mixed-species flocks in tropical regions. Additionally, member species are more vulnerable compared to nuclear species, and overall, forests, savannas, and shrubland formations are the most critical habitats for the conservation of mixed-species flocks in these territories. Therefore, we emphasize that the birds evaluated herein should be at the forefront of global actions aimed at restoring ecosystems, preventing functional extinction, expanding the extent of protected areas, and mitigating the effects of climate change.

Keywords: Birds; Heterospecific Interactions; Mixed-species Flocks; Tropical Hotspots.

¹ Laboratório de Etnoconservação e Áreas Protegidas, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, km 16, 45662-900 Ilhéus, Bahia, Brasil.

² Laboratório de Ornitologia Sala 03, LABIO, Universidade Estadual de Feira de Santana, Avenida Transnordestina s/n, Novo Horizonte, 44036-900 Feira de Santana, Bahia, Brasil.

³ Investigador Asociado CESIMAR/CENPAT, Bv. Almirante Brown 2915, Puerto Madryn, Chubut, Argentina.

Corresponding author ✉. E-mail address: ESF (everton.ornito@gmail.com), CGM (gracom@uol.com.br).

SIGNIFICANCE STATEMENT

Here we present a global overview of the composition and vulnerability of mixed-species flocks of birds in tropical regions and their hotspots based on a systematic review. We demonstrate the focal points of discussion, identify key habitats for the conservation of mixed-species flocks as well as those harboring greater diversity of nuclear species and member species. We also emphasize the role of non-threatened species in the conservation of mixed-species flocks. Finally, we suggest that the species evaluated herein be at the forefront of global discussions focusing on the conservation of ecological interactions, prioritization of areas, and ecological restoration in the tropics and their hotspots.

INTRODUCTION

Mutualistic interactions are essential ecological processes in determining patterns in biological communities, particularly in terms of composition, foraging dynamics, fitness, and distribution (Chomicki *et al.*, 2020; Connor, 1995; Fowler *et al.*, 2023; Swynnerton, 1915). From this perspective, mutualistic dependence predicts that organisms living in isolation have lower adaptive success in the face of biotic and abiotic selective pressures. In this sense, species become, at different levels, dependent on these interactions to persist in ecosystems (Chomicki *et al.*, 2020). This theoretical framework is especially relevant when we seek to understand mutualistic interactions as vulnerable as mixed-species bird flocks, which, not coincidentally, have been studied for over a century (Moynihan, 1962; Powell, 1989, 1985; Swynnerton, 1915).

Mixed-species flocks consist of mutualistic associations in which two or more species forage and move synchronously along the same feeding route (Greenberg, 2000; Moynihan, 1962; Powell, 1989). Unlike fortuitous groupings that form around temporary food sources (e.g., fruit-bearing trees), mixed-species flocks are cohesive and mutually dependent interspecific associations (Greenberg, 2000; Machado, 1999, 1997; Moynihan, 1962; Sridhar *et al.*, 2009). In more detail, the formation, structure, and cohesion of mixed-species flocks in terms of composition depend on stimuli and behavioral responses from their members, especially nuclear species (Moynihan, 1963; Powell, 1989).

Essentially, nuclear species are gregarious, as they attract multiple birds to their vicinity (Moynihan, 1962; Powell, 1989). High movement frequency and vocalizations are also key characteristics of these species, which, in turn, exhibit their typical sentinel behavior in mixed-species flocks (Amaral and Ragusa-Netto, 2008; Bell, 1986; McClure, 1967; Moynihan, 1963; Powell, 1989). In particular, presence in mixed-species flocks results in higher chances of survival and reproduction for birds, as species optimize their foraging and reduce predation risk (Bohórquez, 2003; Greenberg, 2000; Jullien and Clobert, 2000; Machado, 1997; Moynihan, 1963; Sridhar *et al.*, 2009).

Mixed-species bird flocks have been the focus of

studies in ecology and conservation in the tropics, especially due to their vulnerability and dependence on primary forests (Bohórquez, 2003; Buskirk *et al.*, 1972b; Cordeiro *et al.*, 2015; Van Houtan *et al.*, 2006; Mokross *et al.*, 2013; Zou *et al.*, 2018). However, the synergistic impact of native vegetation suppression, expansion of pastures and monocultures, as well as natural and anthropogenic fires, has strongly threatened these regions (Giam, 2017; Hansen *et al.*, 2020; Van Houtan *et al.*, 2006; Symes *et al.*, 2018). This critical scenario justifies the fact that tropical forests are among the most threatened ecosystems on the planet (Symes *et al.*, 2018). Additionally, these impacts may have more devastating consequences in biodiversity hotspots, especially due to the high degree of vulnerability and irreparability of these territories (Brooks *et al.*, 2006; Fischer *et al.*, 2021; Laurance *et al.*, 2000; Myers *et al.*, 2000, 2004).

Specifically, advancing discussions with an emphasis on mixed-species bird flocks reinforces the need to prioritize ecological interactions and, consequently, multispecies in decision-making processes (Ceballos *et al.*, 2020; Cordeiro *et al.*, 2015; Muñoz and Jankowski, 2022; Sainz-Borgo *et al.*, 2018; Valiente-Banuet *et al.*, 2014; Zuluaga *et al.*, 2015). Currently, this scenario assumes an urgent character considering that birds have been assessed as the vertebrates most impacted by anthropogenic actions, constituting 65% of species on the brink of extinction (Ceballos *et al.*, 2020).

Birds are associated with multiple ecosystem services including pollination, seed and fruit dispersal, and pest control (Mariyappan *et al.*, 2023; Sekercioglu *et al.*, 2004; Whelan *et al.*, 2008, 2015). However, these and numerous ecological interactions dependent on multitaxa may be extinct due to the massive population decline of species (Valiente-Banuet *et al.*, 2014; Zhou *et al.*, 2019). From this perspective and the current scenario of mass extinction (Ceballos *et al.*, 2020), species would be living in such low densities as to become functionally extinct (Valiente-Banuet *et al.*, 2014).

In extensively fragmented landscapes, for example, it has been demonstrated that the functional extinction of birds directly impacts seed dispersal (Galett and Guevara, 2013), besides reducing the potential resilience of tropical forests

(Hatfield *et al.*, 2023). Specifically, mixed-species flocks are fundamental for maintaining birds in ecosystems, enabling greater taxon diversity and structuring multiple interaction networks (Goodale *et al.*, 2020; Powell, 1989). In this sense, the destruction of megadiverse ecosystems such as tropical forests represents the imminent decline of multispecies associated with mixed-species flocks and, consequently, the extinction of these interactions (Valiente-Banuet *et al.*, 2014). Additionally, this impact may be greater on birds that frequent mixed-species flocks during periods of food scarcity and, in turn, depend on this interaction to resist numerous selective pressures (da Luz *et al.*, 2022; Maldonado-Coelho and Marini, 2004, 2000, 2003; Mokross *et al.*, 2013; Rutt *et al.*, 2020).

Although representing less than 10% of the Earth's surface, tropical regions harbor two-thirds of all species on the planet, constituting one of the most biodiverse territories globally (DeFries *et al.*, 2007; Giam, 2017; Hansen *et al.*, 2020; Saatchi *et al.*, 2021). However, the historical scenario of primary forest suppression in the tropics has increasingly demanded urgency in decision-making processes (Edwards *et al.*, 2019). This includes careful land use planning (Hansen *et al.*, 2020), combined efforts in restoration (Brancalion *et al.*, 2019; Strassburg *et al.*, 2020), sustainable action direction, and expansion of protected area coverage (Cazalis *et al.*, 2020; Hansen *et al.*, 2020; Vidal *et al.*, 2016). This integrated effort has already proven effective in reducing deforestation in forest ecosystems as well as in the conservation of birds in at least eight of the planet's most threatened hotspots (Brooks, 2010; Cazalis *et al.*, 2020; Giam, 2017; Mittermeier *et al.*, 2011; Myers *et al.*, 2004). Mixed-species bird flocks have been the subject of numerous discussions in tropical regions and their hotspots (Bohórquez, 2003; Buskirk *et al.*, 1972a; Goodale and Kotagama, 2005; Machado, C.G. & Rodrigues, 2000; Machado, 2002; Martínez and Gomez, 2013; Richard, 1994; Swynnerton, 1915; Tien *et al.*, 2005; Zuluaga *et al.*, 2015), but apparently there has not yet been an integrated assessment in terms of composition, vulnerability, or habitat use of these birds.

Here, we analyze the composition and vulnerability of mixed-species flocks in the tropics and their biodiversity hotspots based on a systematic review. Thus, we present a list of the most frequent birds in mixed-species flocks in studies conducted in the tropics and indicate which hotspots are the focus of these discussions. Additionally, we emphasize the standout regions in terms of threatened birds in mixed-species flocks and, particularly, nuclear species. Finally, we demonstrate, from a general ecological perspective, the priority habitats in terms of restoration and, therefore,

conservation of mixed-species flocks.

MATERIAL AND METHODS

Literature research

The data were collected from technical publications, primarily focusing on articles, and also including monographs, master's theses, or doctoral dissertations on mixed-species bird flocks in the tropics. To avoid duplication of information, we assessed whether these recent studies had been published subsequently. In such cases, only the publications were considered. The studies were obtained from the following sources: Web of Science, Scopus, Academia.edu, ResearchGate, Google Scholar, and Mendeley. Generally, we searched for studies in English, French, Spanish, and Portuguese using the following keywords: mixed-species bird flocks, interspecific bird behavior, and heterospecific bird flocks.

Criteria for Inclusion or Exclusion of Studies

Only studies conducted in tropical regions were included. Specifically, the following countries and territories were included: Argentina, Bolivia, Brazil, Colombia, Colombia, Costa Rica, Cuba, Ecuador, French Guiana, Vietnam, India, Mexico, Panama, Republic of the Congo, Sri Lanka, Tanzania, Venezuela, India, and the island of Madagascar. Additionally, we did not set an initial period for the searches, and thus included both historical studies (from the second half of the 19th century) and those completed/published up to December 2022. Conversely, we excluded reviews and studies conducted in predominantly anthropized ecosystems, such as monocultures or pastures. The included studies included different types of documents: articles based on field studies, literature reviews, master's theses and doctoral dissertations, which in turn were our main source of data for the construction of statistical analyses and maps.

Representation of Studies and Key Species of Mixed-species Flocks in the Tropics

We analyzed the studies for two sets of information. First, in the "Materials and Methods" section, we consulted the locations and respective exact or approximate geographical coordinates provided by the authors. When not available, we obtained general coordinates from the described locations. Thus, we analyzed whether the studies were primarily

conducted in the tropics, and when applicable, in biodiversity hotspots. Next, we extracted from the “Results” and “Discussion” sections the most frequent birds in mixed-species flocks. This was possible because we noticed that authors generally use expressions such as “target species” and “species most frequently recorded” to emphasize patterns in flock composition. Additionally, according to authors’ classifications, we defined the birds as either member species (i.e., members of mixed-species flocks) or nuclear species.

Based on this information, we constructed maps demonstrating the distribution of historical studies and nuclear species in tropical biodiversity hotspots. We also assessed the representativeness of these regions focusing on nuclear species, particularly, as well as threatened mixed-species flock birds, in general. The maps were produced using the trial version of the ArcGIS software (Esri, 2023). All the functionalities available in the version used (e.g. definition of colour gradients and layer clipping) fully met the requirements for the construction, layout definition and finalisation of the maps.

Definition of Target Group

After assessing the threat category of species based on criteria from the International Union for Conservation of Nature (IUCN, 2024), we defined a target group for the analyses. In detail, the group consisted of species with two ecological profiles: 1) threatened, not at risk, or near threatened member species, and 2) nuclear species (under some degree of threat or not at risk). Predominantly, threatened species are targeted for monitoring, management, and conservation programs, being crucial in decision-making worldwide (Howard *et al.*, 2020; Strassburg *et al.*, 2020). Near threatened species, in turn, become similarly relevant due to the potential decline in their populations and reduction in their distribution in the face of extensive ecosystem destruction (Margules and Pressey, 2000; Zhou *et al.*, 2019).

Particularly, nuclear species directly influence the formation and cohesion of mixed-species flocks as well as the recruitment of individuals, even if member species are declining (Goodale and Beauchamp, 2010; Powell, 1989; Zou *et al.*, 2018). Species assessed as not at risk, on the other hand, are crucial for maintaining ecosystem services and resilience, especially as they are typically the most abundant (Baker *et al.*, 2019).

As a result of this assessment, the target group was composed of birds included in the following categories: Least Concern (LC), Near Threatened (NT), Vulnerable (VU), Endangered (EN), and Critically Endangered (CR). Based on these categories, we established a vulnerability ranking

ranging from one to five. Lower values represented a lower degree of vulnerability, while higher values represented a higher degree of vulnerability (IUCN, 2024).

Additionally, we sought information about the most common habitats for the presence of these species and constructed a general ecological profile for the target group (IUCN, 2023). As a result of this analysis, we evaluated the following habitats: Forests, Savannas, Shrublands, and Wetlands (IUCN, 2023). Lastly, we sought to identify the main impacts on threatened species in order to build a general vulnerability profile for these birds and their hotspots (IUCN, 2023). We used the IUCN’s taxonomic classification system as a criterion to compile the final list of birds, as well as to update nomenclature that has changed over the years.

Statistical Analyses

We conducted three statistical tests to assess the vulnerability and ecological preference of the target group. To analyze the level of vulnerability, we performed a non-parametric Mann-Whitney test (Nachar, 2008), comparing the values established in the ranking between nuclear species and member species ($p < 0.05$). Seeking to demonstrate a general pattern of habitat use based on a global assessment (IUCN, 2023), we performed two ordinations via NMDS - non-metric multidimensional scaling (Agarwal *et al.*, 2007), with a stress value defined between 0.1 and 0.3. In the first analysis, we considered only nuclear species, and in the second, all species in the target group were included. We used presence and absence matrices to assess these general ecological patterns in the ordinations. The matrices were constructed from categorical data, taking into account the habitat type mentioned on the IUCN platform. Thus, we assigned a value of ‘1’ to indicate the occurrence of species in phytophysognomic formations (habitats) such as forest formations, savannas, shrub formations or wetlands, and ‘0’ to indicate no correspondence to such habitats. From this point we constructed the ordinations and calculated the diversity indices.

Additionally, we calculated the Shannon-Wiener diversity index (H) for each habitat considering the usage profile of these formations by the target group. Then, we compared the diversity between habitats using a 95% confidence interval estimated via bootstrapping with 9999 randomizations (Johnson, 2001). The analyses and graphs were conducted using the software Past (v. 4.03) (Hammer *et al.*, 2001) and the trial version of Statistica (v.10) (Icn, 2011). Conventionally, some information contained in the figures and graphs is presented in English.

RESULTS

We identified 269 studies focusing on mixed-species flocks, of which 55 were conducted in tropical regions from 1963 to 2022 (Additional File 1), during which 367 species were the most frequently studied (Additional File 2). The evaluated studies consisted of 50 articles, two monographs, one master's thesis, and two doctoral dissertations. Specifically, the highest number of studies conducted in the Atlantic Forest makes this hotspot the center of historical discussions focusing on mixed-species flocks of birds in the tropics. Following this, the Tropical Andes, Mesoamerica, the Himalayas, the Southwest China Mountains, the Western Ghats, Sri Lanka, and lastly, the Eastern African Coastal Forest are highlighted (Figure 1). Nuclear species, on the other hand, exhibit a more pronounced concentration in the Tropical Andes, followed by the Atlantic Forest, Mesoamerica, the Himalayas, the Southwest China Mountains, the Western Ghats, Sri Lanka, and lastly, the Eastern African Coastal Forest (Figure 2).

According to recent literature records and global assessments (IUCN 2024), five tropical hotspots harbor threatened birds of mixed-species flocks. Particularly, the Tropical Andes harbor the highest number of threatened species, followed by the Eastern African Coastal Forest, the Western Ghats, Sri Lanka, and on a third level, the Himalayas (Table 1). In

relation to the total number of mixed-species flock species recorded in the studies, 92.6% are classified as Least Concern (LC), among which 15.2% are nuclear species. Additionally, we noted similar sets of impacts on threatened species and their respective hotspots. Logging, large-scale development, road construction, establishment of pastures, and wildfires constitute the main anthropogenic threats to these territories (IUCN, 2024).

Based on the target group (Table 2), we demonstrate that nuclear species and member species differ significantly in terms of vulnerability (via ranking). From this general perspective, we noticed that member species are more vulnerable (Figure 3) compared to nuclear species. Additionally, we observed that nuclear species are more associated with forests, savannas, and shrubland formations (Figure 4, a). We also noted that forests and shrublands represent the most critical habitats for the conservation of the target group overall (Figure 4, b).

The reflection of habitat association patterns was reinforced by diversity indices. Specifically, forests harbor the highest species diversity, followed by savannas and shrubland formations (Figure 5). Wetlands are at the lowest level, which, based on the literature reviewed here, featured a species classified as Near Threatened according to global criteria (Figure 5).




Figure 1. Number of studies on mixed flocks of birds in tropical hotspots. The light and dark shades representan ascending order from the lowest to the highest values.




Figure 2. Number of nuclear species of mixed flocks in tropical hotspots.

Table 1. Threatened mixed flock species, hotspots where they have been studied and global conservation overview.

Species	Hotspots	General scenario (IUCN, 2024).
<i>Basileuterus griseiceps</i> Sclater & Salvin, 1869 (EN)	Tropical Andes	No research or monitoring actions, but sites identified for conservation and occurrence in protected áreas (PAs).
<i>Diglossa venezuelensis</i> Chapman, 1925 (EN)		
<i>Premnoplex tatei</i> Chapman, 1925 (EN)		
<i>Bangsia melanochlamys</i> (Hellmayr, 1910), (VU)		
<i>Hypopyrrhus pyrohypogaster</i> (de Tarragon, 1847), (VU)	Western Ghates and Sri Lanka	(Idem).
<i>Argya cinereifrons</i> (Blyth, 1851), (VU)		
<i>Urocissa ornata</i> (Wagler, 1829), (VU)		(Idem).
<i>Sturnornis albofrontatus</i> (Layard, 1854), (VU)	Sri Lanka	
<i>Sitta formosa</i> Blyth, 1843 (VU)		(Idem).
<i>Phaenicophaeus pyrrhocephalus</i> (Pennant, 1769), (VU)		
<i>Anthreptes rubritorques</i> Reichenow, 1905 (VU)	Himalaya	(Idem).
<i>Artisornis moreaui</i> (W.L. Sclater, 1931), (CR)		Actions directed at systematic monitoring, sites identified for conservation and occurrence in PAs.
<i>Hedydipna pallidigaster</i> (Sclater & Moreau, 1935), (EN)		

Table 2. Target species group. *= nuclear species; a= Forest; b= Savannahs; c= Shrub formations; d= Wetlands

Species	English Name	IUCN	Habitat
<i>Artisornis moreaui</i> (W.L. Sclater, 1931)	Long-billed Forest-warbler	CR	a, c
<i>Hedydipna pallidigaster</i> (Sclater & Moreau, 1935)	Amani Sunbird	EN	a, b
<i>Basileuterus griseiceps</i> Sclater & Salvin, 1869	Grey-headed Warbler	EN	a
<i>Diglossa venezuelensis</i> Chapman, 1925	Venezuelan Flowerpiercer	EN	a, c
<i>Premnoplex tatei</i> Chapman, 1925	White-throated Barbtail	EN	a
<i>Argya cinereifrons</i> (Blyth, 1851)	Ashy-fronted Babbler	VU	a
<i>Anthreptes rubritorques</i> Reichenow, 1905	Banded Sunbird	VU	a, b
<i>Bangsia melanocephala</i> (Hellmayr, 1910)	Black-and-gold Tanager	VU	a
<i>Hypopyrrhus pyrohypogaster</i> (de Tarragon, 1847)	Red-bellied Grackle	VU	a
<i>Sitta formosa</i> Blyth, 1843	Beautiful Nuthatch	VU	a
<i>Sturnornis albofrontatus</i> (Layard, 1854)	White-faced Starling	VU	a
<i>Urocissa ornata</i> (Wagler, 1829)	Sri Lanka Blue Magpie	VU	a
<i>Xiphorhynchus pardalotus</i> (Vieillot, 1818)	Chestnut-rumped Woodcreeper	NT	a, b
<i>Vermivora chrysoptera</i> (Linnaeus, 1766)	Golden-winged Warbler	NT	a, c, d
<i>Amazona leucocephala</i> (Linnaeus, 1758)	Cuban Amazon	NT	a, b
<i>Arizelocichla milanjensis</i> (Shelley, 1894)	Stripe-cheeked Bulbul	NT	a, c
<i>Myiothlypis cinereicollis</i> Sclater, 1865	Grey-throated Warbler	NT	a
<i>Iridosornis porphyrocephalus</i> (Sclater, 1856)	Purplish-mantled Tanager	NT	a
<i>Leptasthenura setaria</i> (Temminck, 1824)	Araucaria Tit-spinetail	NT	a
<i>Melopyrrha nigra</i> (Linnaeus, 1758)	Cuban Bullfinch	NT	a, c
<i>Myrmotherula unicolor</i> (Ménétries, 1835)	Unicolored Antwren	LC	a, b
<i>Conirostrum binghami</i> (Chapman, 1919)	Giant Conebill	NT	a
<i>Patagioenas leucocephala</i> Linnaeus, 1758	White-crowned Pigeon	NT	a
<i>Setophaga cerulea</i> (Wilson, 1810)	Cerulean Warbler	NT	a
<i>Tauraco fischeri</i> (Reichenow, 1878)	Fischer's Turaco	NT	a, c
<i>Tangara cyanoptera</i> (Vieillot, 1817)	Azure-shouldered Tanager	NT	a
<i>Vireo atricapilla</i> Woodhouse, 1852	Black-capped Vireo	NT	a, c
<i>Phaenicophaeus pyrrhocephalus</i> * (Pennant, 1769)	Red-faced Malkoha	VU	a
<i>Epinecrophylla gutturalis</i> (Sclater & Salvin, 1881)	Brown-bellied Antwren	LC	a
<i>Thamnomanes ardesiacus</i> * (Sclater & Salvin, 1868)	Dusky-throated Antshrike	LC	a
<i>Basileuterus culicivorus</i> * (Deppe, 1830)	Stripe-crowned Warbler	LC	a
<i>Myiothlypis flaveola</i> * (Baird, 1865)	Flavescent Warbler	LC	a
<i>Basileuterus melanogenys</i> * Baird, 1865	Black-cheeked Warbler	LC	a
<i>Arremon torquatus</i> * (d'Orbigny & Lafresnaye, 1837)	White-browed Brush-finch	LC	a
<i>Chlorospingus canigularis</i> * (Lafresnaye, 1848)	Ashy-throated Bush-tanager	LC	a
<i>Myrmotherula menetriesii</i> * (d'Orbigny, 1837)	Grey Antwren	LC	a, b

to be continued...

Species	English Name	IUCN	Habitat
<i>Myrmotherula axillaris</i> * (Vieillot, 1817)	White-flanked Antwren	LC	a, d
<i>Thamnomanes caesi</i> us* (Temminck, 1820)	Cinereous Antshrike	LC	a
<i>Trichothraupis melanops</i> * (Vieillot, 1818)	Black-goggled Tanager	LC	a
<i>Myiothlypis bivittata</i> * (d'Orbigny & Lafresnaye, 1837)	Two-banded Warbler	LC	a
<i>Tangara inornata</i> * (Gould, 1855)	Plain-colored Tanager	LC	a
<i>Sittasomus griseicapillus</i> * (Vieillot, 1818)	Eastern Olivaceous Woodcreeper	LC	a, b, c
<i>Habia rubica</i> * (Vieillot, 1817)	Red-crowned Ant-tanager	LC	a
<i>Tangara labradorides</i> * (Boissonneau, 1840)	Metallic-green Tanager	LC	a
<i>Lepidocolaptes falcinellus</i> * (Cabanis & Heine, 1859)	Scalloped Woodcreeper	LC	a
<i>Myioborus melanocephalus</i> * (Tschudi, 1844)	Spectacled Whitestart	LC	a, c
<i>Setophaga pitiaiyumi</i> * (Vieillot, 1817)	Tropical Parula	LC	a, b
<i>Dendroma rufa</i> * (Vieillot, 1818)	Buff-fronted Foliage-gleaner	LC	a, d
<i>Syndactyla rufosuperciliata</i> * (Lafresnaye, 1832)	Buff-browed Foliage-gleaner	LC	a
<i>Pachysylvia semibrunnea</i> * (Lafresnaye, 1845)	Rufous-naped Greenlet	LC	a
<i>Hemithraupis ruficapilla</i> * (Vieillot, 1818)	Rufous-headed Tanager	LC	a
<i>Hylophilus amaurocephalus</i> * (Nordmann, 1835)	Grey-eyed Greenlet	LC	a, b, c
<i>Mecocerculus leucophrys</i> * (d'Orbigny & Lafresnaye, 1837)	White-throated Tyrannulet	LC	a, c
<i>Leiothlypis ruficapilla</i> * (Wilson, 1811)	Nashville Warbler	LC	a
<i>Myioborus brunniceps</i> * (d'Orbigny & Lafresnaye, 1837)	Brown-capped Whitestart	LC	a
<i>Myioborus miniatus</i> * (Swainson, 1827)	Slate-throated Whitestart	LC	a
<i>Phylloscartes oustaleti</i> * (Sclater, 1887)	Oustalet's Tyrannulet	LC	a
<i>Thamnomanes schistogynus</i> * Hellmayr, 1911	Bluish-slate Antshrike	LC	a
<i>Argya rufescens</i> * (Blyth, 1847)	Orange-billed Babbler	LC	a, c
<i>Anisognathus igniventri</i> * (d'Orbigny & Lafresnaye, 1837)	Fire-bellied Mountain-tanager	LC	a, c
<i>Anisognathus somptuosus</i> * (Lesson, 1831)	Blue-winged Mountain-tanager	LC	a
<i>Basileuterus hypoleucus</i> * Bonaparte, 1850	White-bellied Warbler	LC	a
<i>Buthraupis montana</i> * (d'Orbigny & Lafresnaye, 1837)	Hooded Mountain-tanager	LC	a
<i>Chlorornis riefferii</i> * (Boissonneau, 1840)	Grass-green Tanager	LC	a
<i>Chlorospingus flavigularis</i> * (Sclater, 1852)	Yellow-throated Bush-tanager	LC	a, c, d
<i>Chlorospingus flavopectus</i> * (Lafresnaye, 1840)	Common Bush-tanager	LC	a, c
<i>Diglossa cyanea</i> * (Lafresnaye, 1840)	Masked Flowerpiercer	LC	a, c
<i>Hemithraupis guira</i> * (Linnaeus, 1766)	Guira Tanager	LC	a, b
<i>Hylophilus poicilotis</i> * Temminck, 1822	Rufous-crowned Greenlet	LC	a
<i>Iridosornis jelskii</i> * (Cabanis, 1873)	Golden-collared Tanager	LC	a
<i>Lanio fulvus</i> * (Boddaert, 1783)	Fulvous Shrike-tanager	LC	a
<i>Lanio versicolor</i> * (d'Orbigny & Lafresnaye, 1837)	White-winged Shrike-tanager	LC	a
<i>Microrhophias quixensis</i> * (Cornalia, 1849)	Dot-winged Antwren	LC	a
<i>Orthogonys chloricterus</i> * (Vieillot, 1819)	Olive-green Tanager	LC	a

to be continued...

Species	English Name	IUCN	Habitat
<i>Phylloscartes ventralis</i> * (Temminck, 1824)	Mottle-cheeked Tyrannulet	LC	a
<i>Polioptila caerulea</i> * (Linnaeus, 1766)	Blue-grey Gnatcatcher	LC	a, b, c
<i>Polioptila plumbea</i> * (Gmelin, 1788)	Tropical Gnatcatcher	LC	a, b, c
<i>Islerothraupis rufiventer</i> * (Spix, 1825)	Yellow-crested Tanager	LC	a
<i>Tangara arthus</i> * Lesson, 1832	Chestnut-breasted Tanager	LC	a
<i>Tangara chilensis</i> * (Vigors, 1832)	Paradise Tanager	LC	a
<i>Tangara aurulenta</i> * Lafresnaye, 1843	Golden Tanager	LC	a
<i>Sporathraupis cyanocephala</i> * (d'Orbigny & Lafresnaye, 1837)	Blue-capped Tanager	LC	a, c
<i>Veniliornis spilogaster</i> * (Wagler, 1827)	White-spotted Woodpecker	LC	a, b
<i>Vireo olivaceus</i> * (Linnaeus, 1766)	Red-eyed Vireo	LC	a
<i>Xiphorhynchus pardalotus</i> * (Vieillot, 1818)	Chestnut-rumped Woodcreeper	LC	a,b

DISCUSSION

In this study, we provide a comprehensive overview of the composition and vulnerability of mixed-species flocks in tropical regions and their biodiversity hotspots (1), identify their general ecological preferences in terms of habitat use (2), and propose targets for restoration and conservation efforts (3). From this perspective, we proceed to illustrate the observed scenarios and emphasize the urgent need for national restoration policies, expansion of protected areas, and maintenance of connectivity in tropical regions and their hotspots. Specifically, forests, savannas, and shrubland habitats should be the primary targets for restoration and, consequently, conservation of mixed-species flocks. Therefore, these integrated actions assume urgency in light of the need to mitigate anthropogenic impacts and their effects on the tropics and on a global scale.

Vulnerability and Conservation Opportunities in Tropical Regions and their Hotspots

In a first scenario, we observed a low number of threatened or near-threatened nuclear species according to global criteria (IUCN, 2024). This outlook is particularly positive when considering the history of fragmentation and threat to primary forests in the Tropical Andes (Comer *et al.*, 2022) and the Atlantic Forest (Willrich *et al.*, 2019). This is because we demonstrated that these hotspots harbor the highest number of nuclear species recorded in the literature. Alternatively, this may reflect the persistence that birds exhibit in fragmented regions and their forest remnants (Pizo and Tonetti, 2020). In this sense, we assume that this may be the general scenario in both hotspots, as this persistence may be associated with the maintenance of original conditions of environmental heterogeneity in post-deforestation landscapes (Willrich *et al.*, 2019).

Additionally, we emphasize that national-level actions can have a broad scope of effectiveness in conserving mixed-species flocks, whose benefits may extend beyond political borders (Mason *et al.*, 2020; Wilson, 2021). In this regard, maintaining functional and structural connectivity between ecosystems (Beier and Noss, 1998; Belote and Wilson, 2020) seems to be the most promising approach in terms of conservation planning for the Tropical Andes, Atlantic Forest, and other tropical hotspots highlighted here (Wilson, 2021).

From this perspective, we highlight the establishment of minimum dynamic areas with a particular focus on adjacent ecosystems as a key action for the conservation of these regions (Belote

and Wilson, 2020; Blanco *et al.*, 2020; Riva and Fahrig, 2022). This scenario of opportunity and urgency also includes alternatives for the conservation of multiple habitat interfaces and their transitions (Belote and Wilson, 2020; Blanco *et al.*, 2020; Riva and Fahrig, 2022). Riparian corridors in fragmented landscapes, for example, have already proven to be effective for the movement of forest birds (Lees and Peres, 2007; Mendes, 2016; Sekercioglu, 2009), an ecological profile that predominates in species of mixed-species flocks in tropical regions (Bohórquez, 2003; Thiollay, 1999). Moreover, riparian forests integrate multiple mosaics, connecting habitats structurally and functionally, and are therefore essential for the conservation of mixed-species flocks (Lees and Peres, 2007; Mendes, 2016; Ribeiro and Walter, 2008; Sekercioglu, 2009).

We also demonstrate that mixed-species flocks recorded in literature harbor threatened birds in five tropical hotspots (IUCN, 2024). In this regard, the Tropical Andes constitute the region with the highest number of threatened species, for which there are no specific conservation or research actions (IUCN, 2024). Specifically, the impact of native vegetation loss has increasingly demanded urgency in expanding networks of protected areas in the Tropical Andes (Bax and Francesconi, 2019; Hrdina and Romportl, 2017). This is because in this hotspot, only 10% of all threatened species are effectively covered by protected areas, and moreover, 90% of endemics are unprotected and therefore imminently threatened (Bax and Francesconi, 2019; Comer *et al.*, 2022). From a broader perspective, assessments of the International Vegetation Classification (IVC) emphasize that the Tropical Andes have already lost between 50% to 70% of their forest cover, including up to 54% of shrub formations (Comer *et al.*, 2022). In this sense, we highlight that mixed-species flocks may be under imminent threat in the Tropical Andes, as forests and shrub formations are among the critical habitats for these interactions.

Surprisingly, despite extensive loss of primary formations and low coverage of protected areas, the Tropical Andes are among the global hotspots with the lowest restoration requirements (Brancalion *et al.*, 2019). On the other hand, this may represent a scenario of opportunities not only in the Tropical Andes but also in the other tropical hotspots evaluated here (Hrdina and Romportl, 2017; Ripple *et al.*, 2020; Tonetti *et al.*, 2022). As recently emphasized, natural ecosystem regeneration in tropical hotspots is crucial for preventing the extinction of numerous taxa and maintaining ecosystem services at different scales (Hrdina and Romportl, 2017; Ripple *et al.*, 2020; Tonetti *et al.*, 2022).

According to the studies evaluated here, none of the species in mixed-species flocks from the Atlantic




Figure 3. Mann-Whitney test comparing vulnerability levels between member species and nuclear species.




Figure 4. Ordinations evaluating the nuclear species (A) and the entire target group (B) showing their associations with the different habitats. a = Forest; b = Savannah; c = Shrub formations; d = Wetlands.

Forest are globally threatened (IUCN, 2024). However, we emphasize that from a local perspective, this scenario is critical in terms of habitat availability and connectivity between forest remnants. This is because the Atlantic Forest has already lost 70% of its

original coverage due to extensive fragmentation (Pizo and Tonetti, 2020; Rezende et al., 2018). Particularly, landscape homogenization (Jongman, 2002), reduction in home range size (Mortelliti and Lindenmayer, 2015), and increased isolation among fragments (Blanco et




Figure 5. Comparison of diversity between habitats based on the target group.

al., 2020; Turvey et al., 2015) are imminent threats to mixed-species flocks in this territory.

Even in the face of these multiple impacts, it has been reported that the persistence of birds in landscapes as drastically fragmented as the Atlantic Forest depends directly on landscape characteristics (Matos et al., 2018; Pizo and Tonetti, 2020; Tonetti et al., 2022). Among these, habitat heterogeneity, connectivity dynamics, edge extent, and potential for natural regeneration are critical for bird persistence in the Atlantic Forest and other similarly threatened tropical hotspots (Pizo and Tonetti, 2020; Riva and Fahrig, 2022; Tonetti et al., 2022). A highlight of this scenario is that mixed-species flocks are predominantly composed of insectivorous birds, which, however, represent the least abundant species in fragmented landscapes (Pizo and Tonetti, 2020). This reinforces the indispensable need for terrestrial ecosystem restoration (Abhilash, 2021; Ma et al., 2023) and the establishment of ecological corridors in tropical regions and their hotspots (Abhilash, 2021; Ma et al., 2023; Rutt et al., 2020).

This panorama justifies the fact that the Atlantic Forest has recently been identified as one of the global targets for restoration and, therefore, serves as a model for similarly threatened tropical hotspots due to fragmentation (Brancalion et al., 2019). In a broader context, the restoration and conservation of tropical forests are directly associated, for example, with mitigating the effects of climate change (Edwards et al., 2019; Franchito et al., 2012; Nelson and Chomitz, 2009; Sales et al., 2020). This is because the restoration of these ecosystems is one of the fastest and most promising ways to remove CO₂ from the

atmosphere, representing an essential solution against the accumulation of this greenhouse gas (Edwards et al., 2019; Franchito et al., 2012; Koch and Kaplan, 2022; Tonetti et al., 2022). In this sense, given the current perspective of climate emergency (Ripple et al., 2020), the Tropical Andes and the Atlantic Forest stand out as regions of global opportunities in forest restoration (Abhilash, 2021), conservation of mixed-species flocks, and carbon capture post-extensive restoration (Koch and Kaplan, 2022).

Despite these positive perspectives, we emphasize that the current scenario is concerning, especially because after extensive periods of deforestation, tropical forests become significant emitters of CO₂ into the atmosphere (Mills et al., 2023). In this sense, directing restoration and conservation programs in tropical regions and their hotspots is currently critical (Abhilash, 2021; Pizo and Tonetti, 2020). In general terms, this would represent the maintenance of viable minimum populations, conservation of ecological interactions and ecosystem services, as well as mitigation of climate change. This restoration perspective is especially relevant in the Tropical Andes, Western Ghats, Sri Lanka, Himalayas, and the Coastal Forest of Africa, which harbor threatened mixed-species flocks. Additionally, we highlight the Atlantic Forest as a hotspot with the highest number of nuclear species.

In tropical forests, approximately 130 million forest fragments were recorded by 2018 (Taubert et al., 2018), and recent projections demonstrate that the loss of forest cover in these systems can reduce extensive native forests to small fragments (Edwards et al., 2019; Ma et al., 2023). Included in this list of challenges

is the urgent task of maintaining heterogeneity in ecosystems, especially in landscapes increasingly homogeneous in terms of composition and community structure (Edwards *et al.*, 2019; Ma *et al.*, 2023). As we have demonstrated here, since mixed-species flocks are associated with and exhibit greater diversity in forests, savannas, and shrubland formations, we suggest that these habitats be prioritized for restoration (Abhilash, 2021; Brancalion *et al.*, 2019; IUCN, 2024; Leberger *et al.*, 2020; Marzluff and Ewing, 2001). This is essentially relevant as threatened mixed-species flocks in the Western Ghats, Sri Lanka, Coastal Forest of Africa, Tropical Andes, and the Himalayas are facing similar impacts (IUCN, 2023). In detail, logging, expansion of pastures, and monocultures threaten multiple habitats, their transitions (Kark, 2013), and consequently, mixed-species flocks in these territories.

The tolerance potential expressed by birds in the face of the expansion of rural and urban areas has been particularly evaluated in open tropical ecosystems, including savannas and shrubland formations (Mikula *et al.*, 2023). Essentially, the significant growth of the human population, from 6.4 to 7.7 billion inhabitants in just 17 years, has substantially contributed to the increased pressure on these ecosystems (Potapov *et al.*, 2022; Tollefson, 2019; United Nations, 2019). Specifically, shrubland formations have been intensely impacted by the expansion of monocultures, with estimated losses of nearly 50 thousand hectares in just five years, representing a direct threat to mixed-species flocks (Pool *et al.*, 2014). Even in the face of such impacts, savannas and shrubland formations constitute key habitats for biodiversity and species conservation and, therefore, deserve special attention in terms of restoration and spatial prioritization (Mikula *et al.*, 2023).

Imminent functional extinction and future perspectives for conservation

The functional extinction of ecological interactions such as mixed-species bird flocks is ultimately one of the most concerning consequences, particularly in biodiversity hotspots (Valiente-Banuet *et al.*, 2014). In general terms, the reduction in species population density can directly influence the functioning of ecological interactions (Valiente-Banuet *et al.*, 2014). In this sense, functional extinction can occur even before the disappearance of species (Valiente-Banuet *et al.*, 2014). This is because the massive decline in populations directly impacts ecosystems and their respective services at a faster rate than the actual extinction of species (Valiente-Banuet *et al.*, 2014). Thus, prioritizing the conservation of mixed-species flocks and their distribution areas in the tropics, particularly in hotspots, is an important

step in preventing functional extinction through the conservation of multispecies and their interaction networks (Root *et al.*, 2003; Valiente-Banuet *et al.*, 2014).

This context of ecological interaction extinction applies directly to bird communities since the decline in species results in the loss of functional diversity (Ali *et al.*, 2023). In the Atlantic Forest, for example, insectivorous birds are highly sensitive to the loss of functional diversity (Mariano-Neto and Santos, 2023). However, from an optimistic perspective, the high number of insectivorous species and their ecological redundancy result in better adaptive capacity in response to anthropogenic impacts (Luck *et al.*, 2013). Obviously, this scenario includes species whose populations are more abundant, generally crucial for the maintenance of ecological processes (Baker *et al.*, 2019; Julliard *et al.*, 2006).

In addition to the target group that was the subject of our evaluation, it is crucial to highlight the potential role of the other 282 species classified as of least concern (LC) in terms of mixed-species flock conservation in the tropics and their hotspots. These species can play a critical role in the resilience of mixed-species flocks in the face of habitat fragmentation and loss, mainly due to their greater abundance in ecosystems (Baker *et al.*, 2019). Furthermore, mixed-species flocks rely on populations in constant interaction, making least concern species crucial in terms of structure, species recruitment, and conservation, even in fragmented landscapes (Bates, 1863; Goodale and Beauchamp, 2010; Machado, 2002; Maldonado-Coelho and Marini, 2003; Powell, 1989).

From a vulnerability standpoint, even small declines in population size result in massive individual losses, yet these species may not be included in any IUCN threat category (Baker *et al.*, 2019). The ecological role of least concern species becomes even more relevant considering the vulnerability of threatened species in response to stochastic fluctuations in demographic or environmental factors, even in protected areas (Baker *et al.*, 2019; Evans *et al.*, 2022). Thus, we emphasize the essential role that least concern mixed-species flock birds can play in species recruitment and the maintenance of mixed-species flocks in vulnerable regions such as the tropics and their hotspots.

Looking ahead, we stress the need for a deeper understanding of the intrinsic ecological mechanisms of mixed-species flocks. This includes their stability, habitat selection, and dispersion across different feeding sites in fragmented landscapes (Batista *et al.*, 2013; Develey *et al.*, 2001; Martínez and Gomez, 2013). We also underscore the importance of analyzing functional connectivity between fragment networks through ecological corridors, using mixed-species

flocks as a model. Additionally, niche modeling focusing on nuclear species represents a key alternative in this process. The benefits of such theoretical predictions include filling gaps in the distribution and availability of habitat, as well as the vulnerability of mixed-species flocks to climate change (Beier and Noss, 1998; Borges *et al.*, 2019; Borges and Loyola, 2020; Correa Ayram *et al.*, 2016; Sekercioglu *et al.*, 2004).

Finally, we emphasize that mixed-species flock birds in tropical regions and their hotspots should be at the forefront of global efforts aimed at restoring ecosystems, preventing functional extinction, expanding protected area coverage, and mitigating the effects of climate change.

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

CONTRIBUTION STATEMENT

Conceived of the presented idea: ESF, AS, CGM.

Carried out the experiment: ESF, AS.

Carried out the data analysis: ESF, CGM, AS.

Wrote the first draft of the manuscript: ESF.

Review and final write of the manuscript: ESF, CGM, AS.

REFERENCES

Abhilash PC (2021) **Restoring the unrestored: Strategies for restoring global land during the un decade on ecosystem restoration (un-der).** *Land* 10:1-19.

Agarwal S, Lanckriet G, Wills J, Kriegman D, Cayton L, Belongie S (2007) **Generalized non-metric multidimensional scaling.** 2:1-8.

Ali JR, Blonder BW, Pigot AL, Tobias JA (2023) **Bird extinctions threaten to cause disproportionate reductions of functional diversity and uniqueness.** *Functional Ecology* 37: 162-175.

Amaral PP, Ragusa-Netto J (2008) **Bird mixed-flocks and nuclear species in a tecoma savanna in the Pantanal.** *Brazilian Journal of Biology* 68:511-518.

Baker DJ, Garnett ST, O'Connor J, Ehmke G, Clarke RH, Woinarski JCZ, McGeoch MA (2019) **Conserving the abundance of nonthreatened species.** *Conservation Biology* 33:319-328.

Bates HW (1863) **The Naturalist on the River Amazons.** *London: J. Murray.* 395.

Batista RO, Machado CG, Miguel R dos S (2013) **A composição de bandos mistos de aves em um fragmento de mata atlântica no litoral norte da Bahia.** *Bioscience Journal* 29:2001-2012.

Bax V, Francesconi W (2019) **Conservation gaps and priorities in the Tropical Andes biodiversity hotspot: Implications for the expansion of protected areas.** *Journal of Environmental Management* 232:387-396.

Beier P, Noss RF (1998) **Do Habitat Corridors Provide Connectivity?** *Conservation Biology* 12:1241-1252.

Bell HL (1986) **The social organization and foraging behaviour of three syntopic thornbills *Acanthiza* spp.** In: Keast JA, Recher HF, Ford HA, Saunders D (eds) *Birds of eucalypt forests and woodlands: ecology, conservation, management.* RAOU and Surrey-Beattie, Sydney, pp 151-163.

Belote RT, Wilson MB (2020) **Delineating greater ecosystems around protected areas to guide conservation.** *Conservation Science and Practice* 2:1-10.

Blanco J, Bellón B, Fabricius C, de O. Roque F, Pays O, Laurent F, Fritz H, Renaud PC (2020) **Interface processes between protected and unprotected areas: A global review and ways forward.** *Global Change Biology* 26:1138-1154.

Bohórquez C (2003) **Mixed-Species Bird Flocks in a Montane Cloud Forest of Colombia.** *Ornitologia Neotropical* 14:67-78.

Borges FJA, Loyola R (2020) **Climate and land-use change refugia for Brazilian Cerrado birds.** *Perspectives in Ecology and Conservation* 18:109-115.

Borges FJA, Ribeiro BR, Lopes LE, Loyola R (2019) **Bird vulnerability to climate and land use changes in the Brazilian Cerrado.** *Biological Conservation* 236:347-355.

Brancalion PHS, Niamir A, Broadbent E, Crouzeilles R, Barros FSM, Almeyda Zambrano AM, Baccini A, Aronson J, Goetz S, Leighton Reid J, Strassburg BBN, Wilson S, Chazdon RL (2019) **Global restoration opportunities in tropical rainforest landscapes.** *Science Advances* 5:1-11.

- Brooks T (2010) **Conservation planning and priorities.** *Conservation Biology for All*. 199-219.
- Brooks TM, Mittermeier RA, da Fonseca GAB, Gerlach J, Hoffmann M, Lamoreux JF, Mittermeier CG, Pilgrim JD, Rodrigues ASL (2006) **Global biodiversity conservation priorities.** *Science (New York, N.Y.)* 313:58–61.
- Buskirk W, Powell G V., Wittenberger J, Buskirk R, Powell T (1972a) **Interspecific Bird Flocks in Tropical Highland Panama.** *The Auk: Ornithological Advances* 89:612-624.
- Buskirk WH, Powell GV. N, Wittenberger JF, Buskirk RE, Powell TU. (1972b) **Interspecific bird flocks in tropical highland Panama.** *The Auk* 89:612–624.
- Cazalis V, Princé K, Mihoub JB, Kelly J, Butchart SHM, Rodrigues ASL (2020) **Effectiveness of protected areas in conserving tropical forest birds.** *Nature Communications*. 1-8.
- Ceballos G, Ehrlich PR, Raven PH (2020) **Vertebrates on the brink as indicators of biological annihilation and the sixth mass extinction.** *PNAS* 1–7.
- Chomicki G, Kiers ET, Renner SS (2020) **The Evolution of Mutualistic Dependence.** *Annual Review of Ecology, Evolution, and Systematics* 51:409-432.
- Comer PJ, Valdez J, Pereira HM, Acosta-Muñoz C, Campos F, García FJB, Claros X, Castro L, Dallmeier F, Rivadeneira EYD, Gill M, Josse C, Cartagena IL, Langstroth R, Larrea-Alcázar D, Masur A, Jaramillo GM, Navarro L, Novoa S, Prieto-Albuja F, Ortiz GR, Teran MF, Zambrana-Torrel C, Fernandez M (2022) **Conserving Ecosystem Diversity in the Tropical Andes.** *Remote Sensing* 14:1-17.
- Connor RC (1995) **The Benefits of Mutualism: A Conceptual Framework.** *Biological Reviews* 70:427-457.
- Cordeiro NJ, Borghesio L, Joho MP, Monoski TJ, Mkongewa VJ, Dampf CJ (2015) **Forest fragmentation in an African biodiversity hotspot impacts mixed-species bird flocks.** *Biological Conservation* 188:61-71.
- Correa Ayram CA, Mendoza ME, Etter A, Salicrup DRP (2016) **Habitat connectivity in biodiversity conservation: A review of recent studies and applications.** *Progress in Physical Geography* 40:7-37.
- DeFries R, Hansen A, Turner BL, Reid R, Liu J (2007) **Land use change around protected areas: Management to balance human needs and ecological function.** *Ecological Applications* 17:1031–1038.
- Develey PF, Von Matter S, Straube FC, Accordi IA, Piacentini V de Q, Cândido Jr JF (2001) **Ornitologia e Conservação: Ciência Aplicada, Técnicas de Pesquisa e Levantamento.** 516.
- Edwards DP, Socolar JB, Mills SC, Burivalova Z, Koh LP, Wilcove DS (2019) **Conservation of Tropical Forests in the Anthropocene.** *Current Biology* 29:1008-2020.
- Esri (2023) **ArcGis.** [<https://www.img.com.br/pt-br/arcgis/produtos/arcgis-pro/trial>] Accessed October 28, 2023.
- Evans MJ, Gordon IJ, Pierson JC, Neaves LE, Wilson BA, Brockett B, Ross CE, Smith KJ, Rapley S, Andrewartha TA, Humphries N, Manning AD (2022) **Reintroduction biology and the IUCN Red List: The dominance of species of Least Concern in the peer-reviewed literature.** *Global Ecology and Conservation* 38:1-13.
- Fischer R, Taubert F, Müller MS, Groeneveld J, Lehmann S, Wiegand T, Huth A (2021) **Accelerated forest fragmentation leads to critical increase in tropical forest edge area.** *Science Advances* 7:1-8.
- Fowler JC, Donald ML, Bronstein JL, Miller TEX (2023) **The geographic footprint of mutualism: How mutualists influence species' range limits.** *Ecological Monographs* 93:1-22.
- Franchito SH, Rao VB, Fernandez JPR (2012) **Tropical land savannization: Impact of global warming.** *Theoretical and Applied Climatology* 109:73-79.
- Galett M, Guevara R (2013) **Functional Extinction of Birds Drives Rapid Evolutionary Changes in Seed Size.** *Science* 340:1–32.
- Giam X (2017) **Global biodiversity loss from tropical deforestation.** *Proceedings of the National Academy of Sciences of the United States of America*. 114:5775-5777.
- Goodale E, Beauchamp G (2010) **The relationship between leadership and gregariousness in mixed-species bird flocks.** *Journal of Avian Biology* 41:99-103
- Goodale E, Kotagama SW (2005) **Testing the roles of species in mixed-species bird flocks of a Sri Lankan rain forest.** *Journal of Tropical Ecology* 21:669-676.
- Goodale E, Sridhar H, Sieving KE, Bangal P, Z GJC, Farine DR, Heymann EW, Jones HH, Krams I, Martínez AE, Montaña-Centellas F, Muñoz J, Srinivasan U, Theo A, Shanker K (2020) **Mixed**

- company: a framework for understanding the composition and organization of mixed-species animal groups. *Biological Reviews* 1–22.
- Greenberg R (2000) **Birds of Many Feathers: The Formation and Structure of Mixed- Species Flocks of Forest Birds.** pp. 521–559 .
- Hammer Ø, Harper DAT, Ryan PD (2001) **Past: Paleontological statistics software package for education and data analysis.** *Palaeontologia Electronica* 4..
- Hansen MC, Wang L, Song XP, Tyukavina A, Turubanova S, Potapov P V., Stehman S V. (2020) **The fate of tropical forest fragments.** *Science Advances* 6:1–9.
- Hatfield JH, Banks-Leite C, Barlow J, Lees AC, Tobias JA (2023) **Constraints on avian seed dispersal reduce potential for resilience in degraded tropical forests.** *Functional Ecology.* 00:1–12.
- Howard C, Flather CH, Stephens PA (2020) **A global assessment of the drivers of threatened terrestrial species richness.** *Nature Communications.* 11:1–9.
- Hrdina A, Romportl D (2017) **Evaluating Global Biodiversity Hotspots-Very Rich and even More Endangered.** *Journal of Landscape Ecology(Czech Republic)* 10:108–115.
- Icn S (2011) **STATISTICA.** .
- IUCN (2024) **The IUCN red list of threatened species.** *IUCN* .
- Johnson RW (2001) **An Introduction to the Bootstrap.** *Teaching Statistics* 23:.
- Jongman RHG (2002) **Homogenisation and fragmentation of the European landscape: Ecological consequences and solutions.** *Landscape and Urban Planning* 58:211–221.
- Julliard R, Clavel J, Devictor V, Jiguet F, Couvet D (2006) **Spatial segregation of specialists and generalists in bird communities.** *Ecology Letters.* 9:1237–1244.
- Jullien M, Clobert J (2000) **The survival value of flocking in neotropical birds: Reality or fiction?** *Ecology* 81:3416–3430.
- Kark S (2013) **Effects of Ecotones on Biodiversity.** *Encyclopedia of Biodiversity: Second Edition.* 142–148.
- Koch A, Kaplan JO (2022) **Tropical forest restoration under future climate change.** *Nature Climate Change* 12:279–283.
- Laurance WF, Vasconcelos HL, Lovejoy TE (2000) **Forest loss and fragmentation in the Amazon: Implications for wildlife conservation.** *ORYX* 34:39–45.
- Leberger R, Rosa IMD, Guerra CA, Wolf F, Pereira HM (2020) **Global patterns of forest loss across IUCN categories of protected areas.** *Biological Conservation* 241:1–11.
- Lees AC, Peres CA (2007) **Conservation value of remnant riparian forest corridors of varying quality for amazon birds and mammals.** *Conservation Biology* 22:439–449.
- Luck GW, Carter A, Smallbone L (2013) **Changes in Bird Functional Diversity across Multiple Land Uses: Interpretations of Functional Redundancy Depend on Functional Group Identity.** *PLoS ONE* 8:1–11.
- da Luz GS, Carvalho F, Zocche JJ (2022) **Composition and dynamics of mixed flocks of birds in a remnant of Submontane Atlantic Rain Forest in southern Brazil.** *Papeis Avulsos de Zoologia* 62:1–12.
- Ma J, Li J, Wu W, Liu J (2023) **Global forest fragmentation change from 2000 to 2020.** *Nature Communications* 14:1–10.
- Machado, C.G. and Rodrigues NMR (2000) **Alteração de altura de forrageamento de espécies de aves quando associadas a bandos mistos.** *Ornitologia Brasileira: perspectivas, conservação e pesquisa.* 231–239.
- Machado CG (1997) **Vireo olivaceus (Vireonidae): uma espécie migratória nos bandos mistos de aves na Mata Atlântica do sudeste brasileiro.** *Revista Brasileira de Ornitologia* 5:60–62.
- Machado CG (1999) **A composição dos bandos mistos de aves na Mata Atlântica da Serra de Paranapiacaba, no sudeste brasileiro.** *Revista Brasileira de Biologia* 59:75–85.
- Machado CG (2002) **As espécies-núcleo dos bandos mistos de aves da Mata Atlântica da Serra de Paranapiacaba, no Sudeste brasileiro.** *Sitientibus - Série Ciências Biológicas* 2:85–90.
- Maldonado-Coelho M, Marini MÂ. (2000) **Effects of forest fragment size and successional stage on mixed-species bird flocks in Southeastern Brazil.** *The Condor* 102:585–594.
- Maldonado-Coelho M, Marini MA (2003) **Composição de bandos mistos de aves em fragmentos de mata Atlântica no sudeste do Brasil.** *Papeis Avulsos de Zoologia* 43:31–54.
- Maldonado-Coelho M, Marini MÂ (2004) **Mixed-**

species bird flocks from Brazilian Atlantic forest: The effects of forest fragmentation and seasonality on their size, richness and stability. *Biological Conservation* 116:19-26.

Margules CR, Pressey RL (2000) **Systematic conservation planning.** *Nature*. 405: 243-253.

Mariano-Neto E, Santos RAS (2023) **Changes in the functional diversity of birds due to habitat loss in the Brazil Atlantic Forest.** *Frontiers in Forests and Global Change* 6:1-10.

Mariyappan M, Rajendran M, Velu S, Johnson AD, Dinesh GK, Solaimuthu K, Kaliyappan M, Sankar M (2023) **Ecological Role and Ecosystem Services of Birds: A Review.** *International Journal of Environment and Climate Change* 13:76-87.

Martínez AE, Gomez JP (2013) **Are mixed-species bird flocks stable through two decades?** *American Naturalist* 181:53-59.

Marzluff JM, Ewing K (2001) **Restoration of Fragmented Landscapes for the Conservation of Birds: A General Framework and Specific Recommendations for Urbanizing Landscapes.** *Restoration Ecology* 9:280–292.

Mason N, Ward M, Watson JEM, Venter O, Runting RK (2020) **Global opportunities and challenges for transboundary conservation.** *Nature Ecology and Evolution* 4:1-8.

Matos VPV de, Matos TPV de, Cetra M, Timo TP de C e, Valente RA (2018) **Forest fragmentation and impacts on the bird community.** *Revista Árvore* 42:1-13.

Mcclure HE (1967) **The composition of mixed species flocks in lowland and sub-montane forests of Malaya.** *The Wilson Bulletin* 79: 131-154.

Mendes AI da S (2016) **The use of riparian forests as ecological corridors by passerine birds in the south of portugal.** Universidade de Évora 190.

Mikula P, Tomášek O, Romportl D, Aikins TK, Avendaño JE, Braimoh-Azaki BDA, Chaskda A, Cresswell W, Cunningham SJ, Dale S, Favoretto GR, Floyd KS, Glover H, Grim T, Henry DAW, Holmern T, Hromada M, Iwajomo SB, Lilleyman A, Magige FJ, Martin RO, Marina MF, Nana ED, Ncube E, Ndaimani H, Nelson E, van Niekerk JH, Pienaar C, Piratelli AJ, Pistorius P, Radkovic A, Reynolds C, Røskaft E, Shanungu GK, Siqueira PR, Tarakini T, Tejeiro-Mahecha N, Thompson ML, Wamiti W, Wilson M, Tye DRC, Tye ND, Vehtari A, Tryjanowski P, Weston MA, Blumstein DT, Albrecht T (2023) **Bird tolerance to humans in open tropical**

ecosystems. *Nature Communications* 14:1-10.

Mills MB, Malhi Y, Ewers RM, Kho LK, Teh YA, Both S, Burslem DFRP, Majalap N, Nilus R, Huasco WH, Cruz R, Pillco MM, Turner EC, Reynolds G, Riutta T (2023) **Tropical forests post-logging are a persistent net carbon source to the atmosphere.** *Proceedings of the National Academy of Sciences of the United States of America* 120:1-7.

Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C (2011) **Global Biodiversity Conservation: The Critical Role of Hotspots.** *Biodiversity Hotspots*. 1-20.

Mokross K, Ryder TB, Côrtes MC, Wolfe JD, Stouffer PC (2013) **Decay of interspecific avian flock networks along a disturbance gradient in Amazonia.** *Proceedings of the Royal Society B: Biological Sciences* 281:1-10.

Mortelliti A, Lindenmayer DB (2015) **Effects of landscape transformation on bird colonization and extinction patterns in a large-scale, long-term natural experiment.** *Conservation Biology* 5: 1314-1326.

Moynihan M (1962) **The organization and probable evolution of some mixed species flocks of Neotropical birds** 143: 1-152.

Muñoz J, Jankowski JE (2022) **Neotropical mixed-species bird flocks in a community context.** *Philosophical Transactions of the Royal Society B2* 378:1–15.

Myers N, Mittermeier CG, Mittermeier RA (2004) **Hotspots: Earth's biologically richest and most endangered terrestrial ecoregions.** *Choice Reviews Online* 38:1-200.

Myers N, Mittermeier RA, Mittermeier CG, Fonseca GBA, Kent J (2000) **Biodiversity hotspots for conservation priorities.** *Nature* 403:853-858.

Nachar N (2008) **The Mann-Whitney U: A Test for Assessing Whether Two Independent Samples Come from the Same Distribution.** *Tutorials in Quantitative Methods for Psychology* 4:13-20.

Nelson A, Chomitz KM (2009) **Protected Area Effectiveness in Reducing Tropical Deforestation: A global analysis of the impact of Protections Status** 1-42.

Pizo MA, Tonetti VR (2020) **Living in a fragmented world: Birds in the Atlantic Forest.** *Condor* 122:1-14.

Pool DB, Panjabi AO, Macias-Duarte A, Solhjem DM (2014) **Rapid expansion of croplands in**

Chihuahua, Mexico threatens declining North American grassland bird species. *Biological Conservation* 170:274-281.

Potapov P, Turubanova S, Hansen MC, Tyukavina A, Zalles V, Khan A, Song XP, Pickens A, Shen Q, Cortez J (2022) **Global maps of cropland extent and change show accelerated cropland expansion in the twenty-first century.** *Nature Food* 3:19-28.

Powell (1989) **On the possible contribution of mixed species flocks to species richness in neotropical avifaunas.** *Behavioral Ecology and Sociobiology* 24:387-393.

Powell GVN (1985) **Sociobiology and Adaptive Significance of Interspecific Foraging Flocks in the Neotropics.** *Ornithological Monographs* 36:713-732.

Rezende CL, Scarano FR, Assad ED, Joly CA, Metzger JP, Strassburg BBN, Tabarelli M, Fonseca GA, Mittermeier RA (2018) **From hotspot to hopespot: An opportunity for the Brazilian Atlantic Forest.** *Perspectives in Ecology and Conservation* 16:208-214.

Ribeiro JF, Walter BMT (2008) **As principais fitofisionomias do bioma Cerrado.** *Cerrado: Ecologia e flora*. 1: 151-212.

Richard LH (1994) **The composition and social organization of mixed-species flocks in a tropical deciduous forest in Western Mexico.** *The Condor* 96:105-118.

Ripple WJ, Wolf C, Newsome TM, Barnard P, Moomaw WR (2020) **World Scientists' Warning of a Climate Emergency.** *BioScience* 70:1-5.

Riva F, Fahrig L (2022) **The disproportionately high value of small patches for biodiversity conservation.** *Conservation Letters* 15:1-7.

Root K V., Akçakaya HR, Ginzburg L (2003) **A multispecies approach to ecological valuation and conservation.** *Conservation Biology* 17:196-206.

Rutt CL, Mokross K, Kaller MD, Stouffer PC (2020) **Experimental forest fragmentation alters Amazonian mixed-species flocks.** *Biological Conservation* 242:1-9.

Saatchi S, Longo M, Xu L, Yang Y, Abe H, André M, Aukema JE, Carvalhais N, Cadillo-Quiroz H, Cerbu GA, Chernela JM, Covey K, Sánchez-Clavijo LM, Cubillos I V., Davies SJ, De Sy V, De Vleeschouwer F, Duque A, Sybille Durieux AM, De Avila Fernandes K, Fernandez LE, Gammino V, Garrity DP, Gibbs DA, Gibbon L, Gowae GY, Hansen M, Lee Harris N, Healey SP, Hilton RG, Johnson CM, Kankeu RS,

Laporte-Goetz NT, Lee H, Lovejoy T, Lowman M, Lumbuenamo R, Malhi Y, Albert Martinez JMM, Nobre C, Pellegrini A, Radachowsky J, Román F, Russell D, Sheil D, Smith TB, Spencer RGM, Stolle F, Tata HL, Torres D del C, Tshimanga RM, Vargas R, Venter M, West J, Widayati A, Wilson SN, Brumby S, Elmore AC (2021) **Detecting vulnerability of humid tropical forests to multiple stressors.** *One Earth* 4:988-1003.

Sainz-Borgo C, Koffler S, Jaffé K (2018) **On the adaptive characteristics of bird flocks: Small birds form mixed flocks.** *Ornitologia Neotropical* 29:289-296.

Sales LP, Galetti M, Pires MM (2020) **Climate and land-use change will lead to a faunal "savannization" on tropical rainforests.** *Global Change Biology* 26:7036-7044.

Sekercioglu CH (2009) **Tropical Ecology: Riparian Corridors Connect Fragmented Forest Bird Populations.** *Current Biology* 19:210-213.

Sekercioglu CH, Daily CG, Ehrlich, R P (2004) **Ecosystem consequences of bird declines.** *PNAS* 101:18042-18047.

Sridhar H, Beauchamp G, Shanker K (2009) **Why do birds participate in mixed-species foraging flocks? A large-scale synthesis.** *Animal Behaviour* 78:337-347.

Strassburg BBN, Iribarrem A, Beyer HL, Cordeiro CL, Crouzeilles R, Jakovac CC, Braga Junqueira A, Lacerda E, Latawiec AE, Balmford A, Brooks TM, Butchart SHM, Chazdon RL, Erb KH, Brancalion P, Buchanan G, Cooper D, Díaz S, Donald PF, Kapos V, Leclère D, Miles L, Obersteiner M, Plutzer C, Carlos CA, Scarano FR, Visconti P (2020) **Global priority areas for ecosystem restoration.** *Nature* 586:724-729.

Swynnerton CF. (1915) **The naturalist on the amazons.** 334-354 Symes WS, Edwards DP, Miettinen J, Rheindt FE, Carrasco LR (2018) **Combined impacts of deforestation and wildlife trade on tropical biodiversity are severely underestimated.** *Nature Communications* 9:1-9.

Taubert F, Fischer R, Groeneveld J, Lehmann S, Müller MS, Rödiger E, Wiegand T, Huth A (2018) **Global patterns of tropical forest fragmentation.** *Nature* 554:519-522.

Thiollay J-M (1999) **Frequency of Mixed Species Flocking in Tropical Forest Birds and Correlates of Predation Risk: An Intertropical Comparison.** *Journal of Avian Biology* 30:282-294.

Tien ML, Soh MCK, Sodhi N, Lian PK, Lim SLH

- (2005) **Effects of habitat disturbance on mixed species bird flocks in a tropical sub-montane rainforest.** *Biological Conservation* 122:193-204.
- Tollefson J (2019) **Humans are driving one million species to extinction.** *Nature* 569:171-171.
- Tonetti V, Niebuhr BB, Ribeiro M, Pizo MA (2022) **Forest regeneration may reduce the negative impacts of climate change on the biodiversity of a tropical hotspot.** *Diversity and Distributions* 28: 2956-2970.
- Turvey ST, Crees JJ, Di Fonzo MMI (2015) **Historical data as a baseline for conservation: Reconstructing long-term faunal extinction dynamics in Late Imperial-modern China.** *Proceedings of the Royal Society B: Biological Sciences* 282:1-9.
- United Nations (2019) **Population Division World Population Prospects 2019.** *World Population Prospects - 2019 Revision* .
- Valiente-Banuet A, Aizen MA, Alcantara JM, Juan Arroyo A, Cocucci MG, García MB, García D, Gómez JM, Jordano P, Medel R, Navarro L, Obeso JR, Oviedo R, Ramírez N, Rey PJ, Traveset A, Verdú M, Zamora R (2014) **Beyond species loss: the extinction of ecological interactions in a changing world.** *Functional Ecology* 29: 299-307.
- Van Houtan KS, Pimm SL, Bierregaard RO, Lovejoy TE, Stouffer PC (2006) **Local extinctions in flocking birds in Amazonian forest fragments.** *Evolutionary Ecology Research* 8:129-148.
- Vidal CY, Manguera JR, Farah FT, Rother DC, Rodrigues RR (2016) **Biodiversity Conservation of Forests and their Ecological Restoration in Highly-modified Landscapes.** *Biodiversity in Agricultural Landscapes of Southeastern Brazil* 342: 136-150.
- Whelan CJ, Şekercioğlu ÇH, Wenny DG (2015) **Why birds matter: from economic ornithology to ecosystem services.** *Journal of Ornithology*. 156:227-238.
- Whelan CJ, Wenny DG, Marquis RJ (2008) **Ecosystem Services Provided by Birds.** *Annals of the New York Academy of Sciences* 1134:25–60.
- Willrich G, Lima MR, dos Anjos L (2019) **The role of environmental heterogeneity for the maintenance of distinct bird communities in fragmented forests.** *Emu* 119: 1-9.
- Wilson O (2021) **Putting nature centre stage? The challenges of ‘mainstreaming’ biodiversity in the planning process.** *Journal of Environmental Planning and Management* 66:549-571.
- Zhou L, Peabotuwage I, Gu H, Jiang D, Hu G, Jiang A, Mammides C, Zhang M, Quan RC, Goodale E (2019) **The response of mixed-species bird flocks to anthropogenic disturbance and elevational variation in southwest China.** *Gerontologist* 10:1-13.
- Zou F, Jones H, Colorado Z. GJ, Jiang D, Lee TM, Martínez A, Sieving K, Zhang M, Zhang Q, Goodale E (2018) **The conservation implications of mixed-species flocking in terrestrial birds, a globally-distributed species interaction network.** *Biological Conservation* 224:267–276.
- Zuluaga C, Rodewald GJ, D. A (2015) **Response of mixed-species flocks to habitat alteration and deforestation in the Andes.** *Biological Conservation* 188:72–81.

Received: 01 January 2025

Accepted: 01 March 2025

Published: 18 June 2025

Editor: Rômulo Alves



Additional Files

Add File 1. General list of studies

- dos Anjos L, Collins CD, Holt RD, Volpato GH, Lopes E V., Bochio GM (2015) **Can habitat specialization patterns of Neotropical birds highlight vulnerable areas for conservation in the Atlantic rainforest, southern Brazil?** *Biological Conservation* 188:
- Arbeláez-Cortés E, Rodríguez-Correa HA, Restrepo-Chica M (2011) **Mixed bird flocks: Patterns of activity and species composition in a region of the Central Andes of Colombia.** *Revista Mexicana de Biodiversidad* 82:.
- Ávila BEV (2019) **Variación en bandanas mixtas de aves en un paisaje altoandino del sur del Ecuador.** Universidad Del Azuay,.
- Batista RO, Machado CG, Miguel R dos S (2013) **A composição de bandos mistos de aves em um fragmento de mata atlântica no litoral norte da Bahia.** *Bioscience Journal* 29:2001–2012.
- Brandt CS, Hasenack H, Laps RR, Hartz SM (2009) **Composition of mixed-species bird flocks in forest fragments of southern Brazil.** *Zoologia* 26:488–498.
- Buitrón-Jurado G, Tobar M (2007) **Posible asociación de la ardilla enana *Microsciurus flaviventer* (rodentia: sciuridae) y bandadas mixtas de aves en la Amazonia Ecuatoriana.** *Mastozoología Neotropical* 14:235–240.
- Buskirk W, Powell G V., Wittenberger J, Buskirk R, Powell T (1972) **Interspecific Bird Flocks in Tropical Highland Panama.** *The Auk: Ornithological Advances* 89:.
- Cestari C (2007) **A atração de aves em resposta ao playback de *Habia rubica*: implicações complementares sobre o papel da espécie para coesão de bandos mistos na Estação Ecológica Juréia-Itatins – SP.** *Atualidades Ornitológicas* 136:.
- Chaparro UCC (2012) **Composición de bandadas mixtas durante la estación seca en el bosque nublado montano tropical de la Estación Biológica Wayqecha, Cusco.** Universidad Nacional Agraria La Molina, .
- Colorado GJ, Rodewald AD (2015) **Assembly patterns of mixed-species avian flocks in the Andes.** *Journal of Animal Ecology* 84:.
- Cordeiro NJ, Borghesio L, Joho MP, Monoski TJ, Mkongewa VJ, Dampf CJ (2015) **Forest fragmentation in an African biodiversity hotspot impacts mixed-species bird flocks.** *Biological Conservation* 188:.
- Develey PF, Peres CA (2000) **Resource seasonality and the structure of mixed species bird flocks in a coastal Atlantic forest of southeastern Brazil.** *Journal of Tropical Ecology* 16:16–53.
- Fanjul ME, Echevarria A, Martínez MV (2021) **Estructura y composición de las bandadas mixtas de aves invernales a lo largo del gradiente latitudinal en las selvas montanas de las Yungas, Argentina.** *Acta Zoologica Lilloana* 65:268–286.
- Gannon GR (1934) **Associations of Small Insectivorous Birds.** *Emu - Austral Ornithology* 34:122–129.
- Ghizoni-Jr. IR (2009) **Composição de bandos mistos de aves no Parque Estadual das Araucárias, oeste de Santa Catarina, Brasil.** *Biotemas* 22:143–148.
- Ghizoni-Jr IR, Azevedo MAG (2006) **Composição de bandos mistos de aves florestais de sub-bosque em áreas de encosta e planície da Floresta Atlântica de Santa Catarina, sul do Brasil.** *Biotemas* 19:.
- Godoy FI (2011) **Composição e estrutura de bandos mistos de aves em uma área de plantio de eucalipto (*Eucalyptus* sp.).** *Atualidades Ornitológicas* 193:.
- Goodale E, Kotagama SW (2005) **Alarm calling in Sri Lankan mixed-species bird flocks.** *Auk* 122:.

- Goodale E, Kotagama SW (2006) **Vocal mimicry by a passerine bird attracts other species involved in mixed-species flocks.** *Animal Behaviour* 72:471–477.
- Goodale E, Kotagama SW, Raman TRS, Sidhu S, Goodale U, Parker S, Chen J (2014) **The response of birds and mixed-species bird flocks to human-modified landscapes in Sri Lanka and southern India.** *Forest Ecology and Management* 329:384–392.
- Graves GR, Gotelli NJ (1993) **Assembly of avian mixed-species flocks in Amazonia.** *Proceedings of the National Academy of Sciences of the United States of America* 90:.
- Hamel PB, Kirkconnell A (2005) **Composition of mixed-species flocks of migrant and resident birds in Cuba.** *Cotinga* 24:28–34.
- Van Houtan KS, Pimm SL, Bierregaard RO, Lovejoy TE, Stouffer PC (2006) **Local extinctions in flocking birds in Amazonian forest fragments.** *Evolutionary Ecology Research* 8:.
- Jones HH, Robinson SK (2020) **Vegetation structure drives mixed-species flock interaction strength and nuclear species roles.** *Behavioral Ecology* 31:1–13.
- Jullien M, Clobert J (2000) **The survival value of flocking in neotropical birds: Reality or fiction?** *Ecology* 81:.
- Jullien M, Thiollay JM (1998) **Multi-species territoriality and dynamic of neotropical forest understorey bird flocks.** *Journal of Animal Ecology* 67:.
- Kajiki LN, Montaña-Centellas F, Mangini G, Colorado Z. GJ, Fanjul ME (2018) **Ecology of mixed-species flocks of birds across gradients in the neotropics.** *Revista Brasileira de Ornitologia* .
- da Luz GS, Carvalho F, Zocche JJ (2022) **Composition and dynamics of mixed flocks of birds in a remnant of Submontane Atlantic Rain Forest in southern Brazil.** *Papeis Avulsos de Zoologia* 62:.
- Machado, C.G. Rodrigues NMR (2000) **Alteração de altura de forrageamento de espécies de aves quando associadas a bandos mistos.** *Ornitologia Brasileira: perspectivas, conservação e pesquisa.* p. .
- Machado CG (1999) **A composição dos bandos mistos de aves na Mata Atlântica da Serra de Paranapiacaba, no sudeste brasileiro.** *Revista Brasileira de Biologia* .
- Machado CG (2002) **As espécies-núcleo dos bandos mistos de aves da Mata Atlântica da Serra de Paranapiacaba, no Sudeste brasileiro.** *Sitientibus - Série Ciências Biológicas* 2:.
- Maldonado-Coelho M, Marini MÂ. (2000) **Effects of forest fragment size and successional stage on mixed-species bird flocks in Southeastern Brazil.** *The Condor* 102:585–594.
- Maldonado-Coelho M, Marini MA (2003) **Composição de bandos mistos de aves em fragmentos de mata Atlântica no sudeste do Brasil.** *Papeis Avulsos de Zoologia* .
- Maldonado-Coelho M, Marini MÂ (2004) **Mixed-species bird flocks from Brazilian Atlantic forest: The effects of forest fragmentation and seasonality on their size, richness and stability.** *Biological Conservation* 116:.
- Mammides C, Chen J, Goodale UM, Kotagama SW, Sidhu S, Goodale E (2015) **Does mixed-species flocking influence how birds respond to a gradient of land-use intensity?** *Proceedings of the Royal Society B: Biological Sciences* 282:.
- Mangini GG (2017) **Las bandadas mixtas de aves como estrategia: comportamiento y estacionalidad en la Selva Pedemontana de las Yungas Australes.** Universidad Nacional de Tucumán, .
- Mangini GG, Fanjul ME (2013) **Conociendo las bandadas mixtas de aves y los efectos de la fragmentación en bosques y selvas de la provincia de Salta.** *Temas BGN* 3:68–76.
- Martínez AE, Parra E, Muellerklein O, Vredenburg VT (2018) **Fear-based niche shifts in neotropical birds.** *Ecology*.
- Matthysen E, Cahill JRA (2008) **Mixed flock composition and foraging behavior of insectivorous birds in undisturbed and disturbed fragments of high-Andean Polylepis woodland.** *Ornitologia Neotropical* 19:403–416.

- McDermott ME, Rodewald AD, Matthews SN (2014) **Managing tropical agroforestry for conservation of flocking migratory birds.** *Agroforest Syst* 1–14.
- Mokross K, Potts JR, Rutt CL, Stouffer PC (2018) **What can mixed-species flock movement tell us about the value of Amazonian secondary forests? Insights from spatial behavior.** *Biotropica*.
- Mokross K, Ryder TB, Côrtes MC, Wolfe JD, Stouffer PC (2013) **Decay of interspecific avian flock networks along a disturbance gradient in Amazonia.** *Proceedings of the Royal Society B: Biological Sciences* 281:.
- Moynihan M (1963) **The organization and probable evolution of some mixed species flocks of Neotropical birds.** *Auk* 80:.
- Munoz J (2011) **The role of facilitation in the structure of tropical bird communities: a case study of mixed-species flocks.** The University of British Columbia. The University of British Columbia, .
- Perón G, Crochet P-A (2009) **Edge effect and structure of mixed-species bird flocks in an Afrotropical lowland forest.** *J Ornithol* 150:585–599.
- Powell (1979) **Structure and Dynamics of Interspecific Flocks in a Neotropical Mid-Elevation Forest.** *The Auk* .
- Powell LL, Cordeiro NJ, Stratford JA (2015) **Ecology and conservation of avian insectivores of the rainforest understory: A pantropical perspective.** *Biological Conservation* 1–10.
- Richard LH (1994) **The composition and social organization of mixed-species flocks in a tropical deciduous forest in Western Mexico.** *The Condor* 96:105–118.
- Rodrigues M, Machado CG, Alvares SMR, Galetti M (1994) **Association of the Black-Goggled Tanager (*Trichothraupis melanops*) with Flushers.** *Biotropica* 26:.
- Rodríguez EVB (2019) **Comportamiento agonístico de *Thammomanes ardesiacus* asociado a la teoría de defensa económica en bandadas mixtas (Madre de dios, Perú)**. Universidad Nacional Agraria La Molina, .
- Sridhar H, Sankar K (2008) **Effects of habitat degradation on mixed-species bird flocks in Indian rain forests.** *Journal of Tropical Ecology* 24:135–147.
- Srinivasan U, Raza RH, Quader S (2012) **Patterns of species participation across multiple mixed-species flock types in a tropical forest in northeastern India.** *Journal of Natural History* 46:.
- Stouffer PC, Bierregaard RO (1995) **Use of Amazonian Forest Fragments by Understory Insectivorous Birds Published.** *America* 76:.
- Stratford JA, Stouffer PC (2015) **Forest fragmentation alters microhabitat availability for Neotropical terrestrial insectivorous birds.** *Biological Conservation* 188:.
- Thiollay J-M (1999) **Frequency of Mixed Species Flocking in Tropical Forest Birds and Correlates of Predation Risk: An Intertropical Comparison.** *Journal of Avian Biology* 30:.
- Thiollay JM, Jullien M (1998) **Flocking behaviour of foraging birds in a neotropical rain forest and the antipredator defence hypothesis.** *Ibis* .
- Williams SM, Lindell CA (2018) **Nuclear species in Peruvian Amazonian mixed-species flocks are differentially attractive to transient species and to each other.** *Wilson Journal of Ornithology* 130:.
- Zou F, Jones H, Colorado Z. GJ, Jiang D, Lee TM, Martínez A, Sieving K, Zhang M, Zhang Q, Goodale E (2018) **The conservation implications of mixed-species flocking in terrestrial birds, a globally-distributed species interaction network.** *Biological Conservation* 224:267–276.
- Zuluaga C, Rodewald GJ, D. A (2015) **Response of mixed-species flocks to habitat alteration and deforestation in the Andes.** *Biological Conservation* 188:72–81.

Add File 2. List of the most frequently recorded species in studies of mixed flocks in tropical regions. * =nuclear species; a= Forest; b= Savanna; c= Shrubland; d= Wetlands; LC= Least Concern; NT= Near Threatened; VU= Vulnerable; EN= Ameaçada; CR= Critically Endangered; (Complementary study list) - 1 = Jullien and Thiollay (1998); Jullien (2000); Houtan et al (2006); 2 = Jullien and Thiollay (1998); Jullien (2000); Houtan et al (2006); Markoss et al (2018); 3 = Maldonado-Coelho and Marini (2004); Ghizoni-Jr and Azevedo (2006); Cestari (2007); 4 = Ghizoni-Jr and Azevedo (2006, 2009); 5 = Jullien and Thiollay (1998); Jullien (2000); Houtan et al (2006); Makross et al (2013); Markoss et al (2018); 6 = Jullien and Thiollay (1998); Jullien (2000); Houtan et al (2006); Buitrón-Jurado (2007); Munoz (2011); Martinez et al (2018); Rodriguez (2019); 7 = Anjos et al (2015); Mangini (2017); Fanjul et al (2021); Jullien and Thiollay (1998); Houtan et al (2006); Munoz (2011); Martinez et al (2018); Markoss et al (2018); 8 = Jullien (2000); Houtan et al (2006); Munoz (2011); Makross et al (2013).

Species	English Name	IUCN	Habitat	Works
<i>Artisornis moreaui</i>	Long-billed Forest-warbler	CR	a, c	Cordeiro et al (2015).
<i>Hedydipna pallidigaster</i>	Amani Sunbird	EN	a, b	Cordeiro et al (2015).
<i>Basileuterus griseiceps</i>	Grey-headed Warbler	EN	a	Zuluaga e Rodeward (2015).
<i>Diglossa venezuelensis</i>	Venezuelan Flowerpiercer	EN	a, c	Zuluaga e Rodeward (2015).
<i>Premnoplex tatei</i>	White-throated Barbtail	EN	a	Zuluaga e Rodeward (2015).
<i>Argya cinereifrons</i>	Ashy-fronted Babbler	VU	a	Goodale e Kotagama (2005); Mammides et al (2015).
<i>Anthreptes rubritorques</i>	Banded Sunbird	VU	a, b	Cordeiro et al (2015).
<i>Bangsia melanochlamys</i>	Black-and-gold Tanager	VU	a	Zuluaga e Rodeward (2015).
<i>Hypopyrrhus pyrohypogaster</i>	Red-bellied Grackle	VU	a	Zuluaga e Rodeward (2015).
<i>Phaenicophaeus pyrrhocephalus</i> *	Red-faced Malkoha	VU	a	Goodale e Kotagama (2005).
<i>Sitta formosa</i>	Beautiful Nuthatch	VU	a	Srinivasan et al (2012).
<i>Sturnus albobfrontatus</i>	White-faced Starling	VU	a	Mammides et al (2015).
<i>Urocissa ornata</i>	Sri Lanka Blue Magpie	VU	a	Goodale e Kotagama (2006).
<i>Epinecrophylla gutturalis</i>	Brown-bellied Antwren	NT	a	Powell (1979); Stouffer e Bierregaard (1995).1
<i>Xiphorhynchus pardalotus</i>	Chestnut-rumped Woodcreeper	NT	a, b	Powell (1979); Stouffer e Bierregaard (1995).2
<i>Vermivora chrysoptera</i>	Golden-winged Warbler	NT	a, c, d, f	Buskirk et al (1972); Zuluaga e Rodeward (2015).
<i>Amazona leucocephala</i>	Cuban Amazon	NT	a, b	Hamel e Kinkkconnell (2005).
<i>Arizelocichla milanensis</i>	Stripe-cheeked Bulbul	NT	a, c	Cordeiro et al (2015).
<i>Myiothlypis cinereicollis</i>	Grey-throated Warbler	NT	a	Zuluaga e Rodeward (2015).
<i>Iridosornis porphyrocephalus</i>	Purplish-mantled Tanager	NT	a	Zuluaga e Rodeward (2015).
<i>Leptasthenura setaria</i>	Araucaria Tit-spinetail	NT	a	Anjos et al (2015).
<i>Melopyrrha nigra</i>	Cuban Bullfinch	NT	a, c	Hamel e Kinkkconnell (2005).
<i>Myrmotherula unicolor</i>	Unicolored Antwren	NT	a, b	Ghizoni-Jr e Azevedo (2006); Cestari (2007).
<i>Conirostrum binghami</i>	Giant Conebill	NT	a	Mathysen (2008).
<i>Patagioenas leucocephala</i>	White-crowned Pigeon	NT	a	Hamel e Kinkkconnell (2005).

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Setophaga cerulea</i>	Cerulean Warbler	NT	a	Zuluaga e Rodeward (2015).
<i>Tauraco fischeri</i>	Fischer's Turaco	NT	a, c	Cordeiro et al (2014).
<i>Tangara cyanoptera</i>	Azure-shouldered Tanager	NT	a	Machado e Rodrigues (2000).
<i>Vireo atricapillus</i> *	Black-capped Vireo	NT	a, c	Hutto (1994)
<i>Anisognathus igniventris</i> *	Fire-bellied Mountain-tanager	LC	a, c	Chaparro (2012)
<i>Anisognathus somptuosus</i> *	Blue-winged Mountain-tanager	LC	a	Jones e Robinson (2020)
<i>Basileuterus culicivorus</i> *	Stripe-crowned Warbler	LC	a	Godoy (2011); Da Luz et al., (2022)
<i>Myiothlypis flaveola</i> *	Flavescent Warbler	LC	a	Batista et al (2013)
<i>Basileuterus hypoleucus</i> *	White-bellied Warbler	LC	a	Maldonado-Coelho e Marini (2000)
<i>Basileuterus melanogenys</i> *	Black-cheeked Warbler	LC	a	Moynihan (1963)
<i>Arremon torquatus</i> *	White-browed Brush-finch	LC	a	Mangini e Fanjul (2013)
<i>Buthraupis montana</i> *	Hooded Mountain-tanager	LC	a	Chaparro (2012)
<i>Chlorornis riefferii</i> *	Grass-green Tanager	LC	a	Chaparro (2012)
<i>Chlorospingus canigularis</i> *	Ashy-throated Bush-tanager	LC	a	Jones e Robinson (2020)
<i>Chlorospingus flavigularis</i> *	Yellow-throated Bush-tanager	LC	a, c, d	Munoz (2011)
<i>Chlorospingus flavopectus</i> *	Common Bush-tanager	LC	a, c	Moynihan (1963); Buskirk et al (1972).
<i>Diglossa cyanea</i> *	Masked Flowerpiercer	LC	a, c	Chaparro (2012).
<i>Habia rubica</i> *	Red-crowned Ant-tanager	LC	a	Develey e Peres (2000); Maldonado-Coelho e Marini (2003).3
<i>Hemithraupis guira</i> *	Guira Tanager	LC	a, b	Ghizoni-Jr (2009).
<i>Hemithraupis ruficapilla</i> *	Rufous-headed Tanager	LC	a	Maldonado-Coelho e Marini (2000); Maldonado-Coelho e Marini (2003).
<i>Hylophilus amaurocephalus</i> *	Grey-eyed Greenlet	LC	a, b, c	Maldonado-Coelho e Marini (2003); Batista et al (2013).
<i>Hylophilus poicilotis</i> *	Rufous-crowned Greenlet	LC	a	Machado (1999); Machado (2002).
<i>Iridosornis jelskii</i> *	Golden-collared Tanager	LC	a	Chaparro (2012).
<i>Lanio fulvus</i> *	Fulvous Shrike-tanager	LC	a	Jullien e Thiollay (1998).
<i>Lanio versicolor</i> *	White-winged Shrike-tanager	LC	a	Munoz (2011).
<i>Lepidocolaptes falcinellus</i> *	Scalloped Woodcreeper	LC	a	Ghizoni-Jr e Azevedo (2006); Ghizoni-Jr (2009); Anjos et al (2015).
<i>Mecocerculus leucophrys</i> *	White-throated Tyrannulet	LC	a, c	Mathysen (2008); Mangini e Fanjul (2013); Fanjul et al (2021).
<i>Microrhophias quixensis</i> *	Dot-winged Antwren	LC	a	Munoz (2011).

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Vireo olivaceus</i> *	Red-eyed Vireo	LC	a	Machado (1999); Mangini (2017).
<i>Leiothlypis ruficapilla</i> *	Nashville Warbler	LC	a	Hutto (1994).
<i>Veniliornis spilogaster</i> *	White-spotted Woodpecker	LC	a, b	Ghizoni-Jr (2009).
<i>Argya rufescens</i> *	Orange-billed Babbler	LC	a, c	Goodale e Kotagama (2006); Mammides et al (2015).
<i>Trichothraupis melanops</i> *	Black-goggled Tanager	LC	a	Rodrigues et al (1994); Maldonado-Coelho e Marini (2000; 2003).4
<i>Sporothraupis cyanocephala</i> *	Blue-capped Tanager	LC	a, c	Chaparro (2012).
<i>Thamnomanes schistogynus</i> *	Bluish-slate Antshrike	LC	a	Munoz (2011).
<i>Thamnomanes caesius</i> *	Cinereous Antshrike	LC	a	Powell (1979); Stouffer e Bierregaard (1995). 5
<i>Thamnomanes ardesiacus</i> *	Dusky-throated Antshrike	LC	a	Powell (1979); Stouffer e Bierregaard (1995). 6
<i>Tangara labradorides</i> *	Metallic-green Tanager	LC	a	Jones e Robinson (2020).
<i>Tangara inornata</i> *	Plain-colored Tanager	LC	a	Moynihan (1963).
<i>Tangara arthus</i> *	Chestnut-breasted Tanager	LC	a	Munoz (2011).
<i>Tangara chilensis</i> *	Paradise Tanager	LC	a	Munoz (2011).
<i>Tangara aurulenta</i> *	Golden Tanager	LC	a	Jones e Robinson (2020).
<i>Islerothraupis rufiventer</i> *	Yellow-crested Tanager	LC	a	Munoz (2011)
<i>Syndactyla rufosuperciliata</i> *	Buff-browed Foliage-gleaner	LC	a	Anjos et al (2015); Mangini e Fanjul (2013); Fanjul et al (2021).
<i>Sittasomus griseicapillus</i> *	Eastern Olivaceous Woodcreeper	LC	a, b, c	Machado (1999); Ghizoni-Jr (2009).7
<i>Polioptila plumbea</i> *	Tropical Gnatcatcher	LC	a, b, c	Batista et al (2013).
<i>Polioptila caerulea</i> *	Blue-grey Gnatcatcher	LC	a, b, c	Hutto (1994); Hamel e Kinkkconnell (2005).
<i>Phylloscartes ventralis</i> *	Mottle-cheeked Tyrannulet	LC	a	Mangini e Fanjul (2013); Fanjul et al (2021).
<i>Phylloscartes oustaleti</i> *	Oustalet's Tyrannulet	NT	a	Machado (1999); Machado e Rodrigues (2000); Machado (2002).
<i>Philydor rufum</i> *	Buff-fronted Foliage-gleaner	LC	a, d	Machado (1999); Machado (2002); Maldonado-Coelho e Marini (2000).
<i>Setophaga pitaiayumi</i> *	Tropical Parula	LC	a, b	Ghizoni-Jr e Azevedo (2006); Brandt et al (2009); Ghizoni-Jr (2009).
<i>Pachysylvia semibrunnea</i> *	Rufous-naped Greenlet	LC	a	Jones e Robinson (2020).

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Orthogonys chloricterus</i> *	Olive-green Tanager	LC	a	Machado (1999); Machado (2002).
<i>Myrmotherula menetriesii</i> *	Grey Antwren	LC	a, b	Powell (1979); Stouffer e Bierregaard (1995).8
<i>Myrmotherula axillaris</i> *	White-flanked Antwren	LC	a, d	Stouffer e Bierregaard (1995); Jullien e Thiollay (1998).9
<i>Myiothlypis bivittata</i> *	Two-banded Warbler	LC	a	Mangini e Fanjul (2013). Buskirk et al (1972);
<i>Myioborus miniatus</i> *	Slate-throated Whitestart	LC	a	Munoz (2011); Jones e Robinson (2020).
<i>Myioborus melanocephalus</i> *	Spectacled Whitestart	LC	a, c	Munoz (2011); Chaparro (2012); Ávila (2019).
<i>Myioborus brunnicaps</i> *	Brown-capped Whitestart	LC	a	Mathysen (2008); Mangini e Fanjul (2013); Fanjul et al (2021).
<i>Alcippe poioicephala</i> *	Brown-cheeked Fulvetta	LC	a,c	Sridhar and Sankar (2008).
<i>Deleornis axillaris</i> *	Grey-headed Sunbird	LC	a	Perón and Crochet (2009).
<i>Dicrurus ludwigii</i> *	Square-tailed Drongo	LC	a,b	Cordeiro et al (2015).
<i>Phyllastrephus icterinus</i> *	Icterine Greenbul	LC	a,b	Perón and Crochet (2009).
<i>Teretistris fernandinae</i> *	Yellow-headed Warbler	LC	a,c	Hamel and Kirkconnell (2005).
<i>Teretistris fornsi</i> *	Oriente Warbler	LC	a,c	Hamel and Kirkconnell (2005).
<i>Myrmotherula longipennis</i>	Long-winged Antwren	LC	-	-
<i>Philydor erythrocercum</i>	Rufous-rumped Foliage-gleaner	LC	-	-
<i>Automolus infuscatus</i>	Olive-backed Foliage-gleaner	LC	-	-
<i>Conirostrum speciosum</i>	Chestnut-vented Conebill	LC	-	-
<i>Tunchiornis ochraceiceps</i>	Tawny-crowned Greenlet	LC	-	-
<i>Hypothymis azurea</i>	Black-naped Monarch	LC	-	-
<i>Acritillas indica</i>	Yellow-browed Bulbul	LC	-	-
<i>Leptopogon amaurocephalus</i>	Sepia-capped Flycatcher	LC	-	-
<i>Mniotilta varia</i>	Black-and-white Warbler	LC	-	-
<i>Xenops minutus</i>	White-throated Xenops	LC	-	-
<i>Xiphorhynchus fuscus</i>	Lesser Woodcreeper	LC	-	-
<i>Arremon dorbignii</i>	Stripe-crowned Sparrow	LC	-	-
<i>Automolus leucophthalmus</i>	White-eyed Foliage-gleaner	LC	-	-
<i>Ceuthmochares aereus</i>	Chattering Yellowbill	LC	-	-
<i>Chiroxiphia caudata</i>	Blue Manakin	LC	-	-
<i>Culicicapa ceylonensis</i>	Grey-headed Canary-flycatcher	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Certhiasomus stictolaemus</i>	Spot-throated Woodcreeper	LC	-	-
<i>Setophaga fusca</i>	Blackburnian Warbler	LC	-	-
<i>Setophaga virens</i>	Black-throated Green Warbler	LC	-	-
<i>Dicrurus remifer</i>	Lesser Racquet-tailed Drongo	LC	-	-
<i>Dysithamnus mentalis</i>	Plain Antvireo	LC	-	-
<i>Glyphorhynchus spirurus</i>	Wedge-billed Woodcreeper	LC	-	-
<i>Harpactes fasciatus</i>	Malabar Trogon	LC	-	-
<i>Myiarchus swainsoni</i>	Swainson's Flycatcher	LC	-	-
<i>Myiobius barbatus</i>	Whiskered Flycatcher	LC	-	-
<i>Pachyramphus polychopterus</i>	White-winged Becard	LC	-	-
<i>Pycnonotus barbatus</i>	Common Bulbul	LC	-	-
<i>Tangara cyanocephala</i>	Red-necked Tanager	LC	-	-
<i>Terpsiphone paradisi</i>	Indian Paradise-flycatcher	LC	-	-
<i>Tangara sayaca</i>	Sayaca Tanager	LC	-	-
<i>Tolmomyias sulphureus</i>	Yellow-olive Flatbill	LC	-	-
<i>Turdus rufiventris</i>	Rufous-bellied Thrush	LC	-	-
<i>Cardellina pusilla</i>	Wilson's Warbler	LC	-	-
<i>Xiphocolaptes albicollis</i>	White-throated Woodcreeper	LC	-	-
<i>Abroscopus albogularis</i>	Rufous-faced Warbler	LC	-	-
<i>Agelaius humeralis</i>	Tawny-shouldered Blackbird	LC	-	-
<i>Schoeniparus castaneiceps</i>	Rufous-winged Fulvetta	LC	-	-
<i>Alcippe nipalensis</i>	Nepal Fulvetta	LC	-	-
<i>Chamaetylas fuelleborni</i>	White-chested Alethe	LC	-	-
<i>Alophoixus flaveolus</i>	White-throated Bulbul	LC	-	-
<i>Anabazenops dorsalis</i>	Dusky-cheeked Foliage-gleaner	LC	-	-
<i>Anabazenops fuscus</i>	White-collared Foliage-gleaner	LC	-	-
<i>Anairetes flavirostris</i>	Yellow-billed Tit-tyrant	LC	-	-
<i>Anairetes parulus</i>	Tufted Tit-tyrant	LC	-	-
<i>Stelgidillas gracilirostris</i>	Slender-billed Greenbul	LC	-	-
<i>Eurillas gracilis</i>	Grey Greenbul	LC	-	-
<i>Andropadus masukuensis</i>	Shelley's Greenbul	LC	-	-
<i>Eurillas virens</i>	Little Greenbul	LC	-	-
<i>Hedydipna collaris</i>	Collared Sunbird	LC	-	-
<i>Anthreptes neglectus</i>	Uluguru Violet-backed Sunbird	LC	-	-
<i>Apalis melanocephala</i>	Black-headed Apalis	LC	-	-
<i>Apaloderma vittatum</i>	Bar-tailed Trogon	LC	-	-
<i>Asthenes dorbignyi</i>	Creamy-breasted Canastero	LC	-	-
<i>Asthenes modesta</i>	Cordilleran Canastero	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Atlapetes citrinellus</i>	Yellow-striped Brush-finch	LC	-	-
<i>Atlapetes melanolaemus</i>	Black-faced Brush-finch	LC	-	-
<i>Attila spadiceus</i>	Bright-rumped Attila	LC	-	-
<i>Automolus ochrolaemus</i>	Buff-throated Foliage-gleaner	LC	-	-
<i>Batara cinerea</i>	Giant Antshrike	LC	-	-
<i>Batis mixta</i>	Forest Batis	LC	-	-
<i>Bleda syndactylus</i>	Red-tailed Bristlebill	LC	-	-
<i>Bradypterus lopezi</i>	Evergreen-forest Warbler	LC	-	-
<i>Cryptolybia olivacea</i>	Green Barbet	LC	-	-
<i>Camaroptera brachyura</i>	Bleating Camaroptera	LC	-	-
<i>Campephilus guatemalensi</i>	Pale-billed Woodpecker	LC	-	-
<i>Campethera mombassica</i>	Mombasa Woodpecker	LC	-	-
<i>Pardipicus nivosus</i>	Buff-spotted Woodpecker	LC	-	-
<i>Campylorhamphus falcularius</i>	Black-billed Scythebill	LC	-	-
<i>Chamaeza campanisona</i>	Short-tailed Antthrush	LC	-	-
<i>Chamaeza ruficauda</i>	Rufous-tailed Antthrush	LC	-	-
<i>Chlorochrysa calliparaea</i>	Orange-eared Tanager	LC	-	-
<i>Riccordia ricordii</i>	Cuban Emerald	LC	-	-
<i>Habia carmioli</i>	Carmioli's Tanager	LC	-	-
<i>Cissa chinensis</i>	Common Green Magpie	LC	-	-
<i>Cnemoscopus rubrirostris</i>	Grey-hooded Tanager	LC	-	-
<i>Colaptes auratus</i>	Yellow-shafted Flicker	LC	-	-
<i>Colaptes melanochloros</i>	Green-barred Woodpecker	LC	-	-
<i>Columbina passerina</i>	Common Ground-dove	LC	-	-
<i>Conopophaga lineata</i>	Rufous Gnateater	LC	-	-
<i>Contopus caribaeus</i>	Cuban Pewee	LC	-	-
<i>Coracina azurea</i>	Blue Cuckooshrike	LC	-	-
<i>Ceblepyris caesi</i>	Grey Cuckooshrike	LC	-	-
<i>Coracina melaschistos</i>	Black-winged Cuckooshrike	LC	-	-
<i>Craniolaema erythroptera</i>	Red-faced Spinetail	LC	-	-
<i>Criniger calurus</i>	Red-tailed Greenbul	LC	-	-
<i>Cryptospiza reichenovii</i>	Red-faced Crimsonwing	LC	-	-
<i>Cyanerpes cyaneus</i>	Red-legged Honeycreeper	LC	-	-
<i>Cyanocorax chrysops</i>	Plush-crested Jay	LC	-	-
<i>Dacnis cayana</i>	Blue Dacnis	LC	-	-
<i>Deconychura longicauda</i>	Northern Long-tailed Woodcreeper	LC	-	-
<i>Dendrocincla fuliginosa</i>	Plain-brown Woodcreeper	LC	-	-
<i>Dendrocincla merula</i>	White-chinned Woodcreeper	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Dendrocincla turdina</i>	Plain-winged Woodcreeper	LC	-	-
<i>Dendrocolaptes certhia</i>	Amazonian Barred Woodcreeper	LC	-	-
<i>Dendrocolaptes picumnus</i>	Black-banded Woodcreeper	LC	-	-
<i>Dendrocolaptes platyrostris</i>	Planalto Woodcreeper	LC	-	-
<i>Setophaga caerulescens</i>	Black-throated Blue Warbler	LC	-	-
<i>Setophaga discolor</i>	Prairie Warbler	LC	-	-
<i>Setophaga dominica</i>	Yellow-throated Warbler	LC	-	-
<i>Setophaga magnolia</i>	Magnolia Warbler	LC	-	-
<i>Setophaga palmarum</i>	Palm Warbler	LC	-	-
<i>Setophaga pityophila</i>	Olive-capped Warbler	LC	-	-
<i>Setophaga tigrina</i>	Cape May Warbler	LC	-	-
<i>Dendropicos fuscescens</i>	Cardinal Woodpecker	LC	-	-
<i>Dendroplex picus</i>	Straight-billed Woodcreeper	LC	-	-
<i>Dicrurus aeneus</i>	Bronzed Drongo	LC	-	-
<i>Dicrurus atripennis</i>	Shining Drongo	LC	-	-
<i>Dicrurus paradiseus</i>	Greater Racquet-tailed Drongo	LC	-	-
<i>Drymophila malura</i>	Dusky-tailed Antbird	LC	-	-
<i>Dryoscopus cubla</i>	Black-backed Puffback	LC	-	-
<i>Dumetella carolinensis</i>	Grey Catbird	LC	-	-
<i>Elaenia pallatangae</i>	Sierran Elaenia	LC	-	-
<i>Eleoscytalopus indigoticus</i>	White-breasted Tapaculo	LC	-	-
<i>Elminia albonotata</i>	White-tailed Crested-flycatcher	LC	-	-
<i>Empidonax minimus</i>	Least Flycatcher	LC	-	-
<i>Epinecrophylla erythrura</i>	Rufous-tailed Antwren	LC	-	-
<i>Epinecrophylla leucophthalma</i>	White-eyed Antwren	LC	-	-
<i>Erpornis zantholeuca</i>	White-bellied Erpornis	LC	-	-
<i>Erythrocercus mccallii</i>	Chestnut-capped Flycatcher	LC	-	-
<i>Eubucco bourcierii</i>	Red-headed Barbet	LC	-	-
<i>Euphonia pectoralis</i>	Chestnut-bellied Euphonia	LC	-	-
<i>Euphonia violacea</i>	Violaceous Euphonia	LC	-	-
<i>Falco sparverius</i>	American Kestrel	LC	-	-
<i>Formicivora grisea</i>	Southern White-fringed Antwren	LC	-	-
<i>Geothlypis trichas</i>	Common Yellowthroat	LC	-	-
<i>Glaucidium siju</i>	Cuban Pygmy-owl	LC	-	-
<i>Glyphorhynchus spirurus</i>	Wedge-billed Woodcreeper	LC	-	-
<i>Grallaria varia</i>	Variegated Antpitta	LC	-	-
<i>Gymnopithys rufigula</i>	Rufous-throated Antbird	LC	-	-
<i>Harpactes erythrocephalus</i>	Red-headed Trogon	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Heliobletus contaminatus</i>	Sharp-billed Treehunter	LC	-	-
<i>Helmitheros vermivorum</i>	Worm-eating Warbler	LC	-	-
<i>Hemipus picatus</i>	Bar-winged Flycatcher-shrike	LC	-	-
<i>Kleinothraupis atropileus</i>	Black-capped Hemispingus	LC	-	-
<i>Thlypopsis superciliaris</i>	Eyebrowed Hemispingus	LC	-	-
<i>Hemitriccus margaritaceiventer</i>	Pearly-vented Tody-tyrant	LC	-	-
<i>Leioptila annectens</i>	Rufous-backed Sibia	LC	-	-
<i>Hylexetastes perrotii</i>	Red-billed Woodcreeper	LC	-	-
<i>Hylopezus nattereri</i>	Speckle-breasted Antpitta	LC	-	-
<i>Tunchiornis ochraceiceps</i>	Tawny-crowned Greenlet	LC	-	-
<i>Willisornis poecilinotus</i>	Common Scale-backed Antbird	LC	-	-
<i>Hypsipetes leucocephalus</i>	Black Bulbul	LC	-	-
<i>Icterus dominicensis</i>	Cuban Oriole	LC	-	-
<i>Icterus pyrrhopterus</i>	Variable Oriole	LC	-	-
<i>Illadopsis rufipennis</i>	Pale-breasted Illadopsis	LC	-	-
<i>Indicator variegatus</i>	Scaly-throated Honeyguide	LC	-	-
<i>Irena puella</i>	Asian Fairy-bluebird	LC	-	-
<i>Xiphorhynchus fuscus</i>	Lesser Woodcreeper	LC	-	-
<i>Lepidocolaptes squamatus</i>	Scaled Woodcreeper	LC	-	-
<i>Lepidocolaptes affinis</i>	Northern Spot-crowned Woodcreeper	LC	-	-
<i>Leptasthenura fuliginiceps</i>	Brown-capped Tit-spinetail	LC	-	-
<i>Sylviorthorhynchus yanacensis</i>	Tawny Tit-spinetail	LC	-	-
<i>Leptopogon superciliaris</i>	Slaty-capped Flycatcher	LC	-	-
<i>Limnothlypis swainsonii</i>	Swainson's Warbler	LC	-	-
<i>Lochmias nematura</i>	Streamcreeper	LC	-	-
<i>Chlorophoneus multicolor</i>	Many-coloured Bush-shrike	LC	-	-
<i>Malimbus nitens</i>	Blue-billed Malimbe	LC	-	-
<i>Mecocerculus stictopterus</i>	White-banded Tyrannulet	LC	-	-
<i>Megalaima viridis</i>	White-cheeked Barbet	LC	-	-
<i>Melanochlora sultanea</i>	Sultan Tit	LC	-	-
<i>Pericrocotus flammeus</i>	Scarlet Minivet	LC	-	-
<i>Dendropicos griseocephalus</i>	Olive Woodpecker	LC	-	-
<i>Microbates collaris</i>	Collared Gnatwren	LC	-	-
<i>Microspingus erythrophrys</i>	Rusty-browed Warbling-finch	LC	-	-
<i>Mimus polyglottos</i>	Northern Mockingbird	LC	-	-
<i>Mionectes rufiventris</i>	Grey-hooded Flycatcher	LC	-	-
<i>Muscicapa adusta</i>	African Dusky Flycatcher	LC	-	-
<i>Myiarchus sagrae</i>	La Sagra's Flycatcher	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Myiarchus tyrannulus</i>	Brown-crested Flycatcher	LC	-	-
<i>Myiopagis viridicata</i>	Greenish Elaenia	LC	-	-
<i>Myrmornis torquata</i>	Southern Wing-banded Antbird	LC	-	-
<i>Isleria guttata</i>	Rufous-bellied Antwren	LC	-	-
<i>Myrmotherula longipennis</i>	Long-winged Antwren	LC	-	-
<i>Cyanomitra olivacea</i>	Olive Sunbird	LC	-	-
<i>Neocossyphus rufus</i>	Red-tailed Ant-thrush	LC	-	-
<i>Nicator gularis</i>	Eastern Nicator	LC	-	-
<i>Ochthoeca leucophrys</i>	White-browed Chat-tyrant	LC	-	-
<i>Ochthoeca oenanthoides</i>	D'Orbigny's Chat-tyrant	LC	-	-
<i>Onychorhynchus coronatus</i>	Amazonian Royal Flycatcher	LC	-	-
<i>Oriolus chlorocephalus</i>	Green-headed Oriole	LC	-	-
<i>Pachyramphus castaneus</i>	Chestnut-crowned Becard	LC	-	-
<i>Pachyramphus validus</i>	Crested Becard	LC	-	-
<i>Pachyramphus versicolor</i>	Barred Becard	LC	-	-
<i>Pachyramphus viridis</i>	Green-backed Becard	LC	-	-
<i>Setophaga americana</i>	Northern Parula	LC	-	-
<i>Machlolophus xanthogenys</i>	Black-lored Tit	LC	-	-
<i>Pernostola rufifrons</i>	Black-headed Antbird	LC	-	-
<i>Pericrocotus flammeus</i>	Scarlet Minivet	LC	-	-
<i>Pheucticus aureoventris</i>	Black-backed Grosbeak	LC	-	-
<i>Philydor atricapillus</i>	Black-capped Foliage-gleaner	LC	-	-
<i>Anabacerthia lichtensteini</i>	Ochre-breasted Foliage-gleaner	LC	-	-
<i>Phyllastrephus cabanisi</i>	Cabanis's Greenbul	LC	-	-
<i>Phyllastrephus debilis</i>	Lowland Tiny Greenbul	LC	-	-
<i>Phyllastrephus flavostriatus</i>	Yellow-streaked Greenbul	LC	-	-
<i>Phyllastrephus xavieri</i>	Xavier's Greenbul	LC	-	-
<i>Phylloscopus occipitalis</i>	Western Crowned Leaf-warbler	LC	-	-
<i>Phylloscopus reguloides</i>	Blyth's Leaf-warbler	LC	-	-
<i>Phylloscopus ruficapilla</i>	Yellow-throated Woodland-warbler	LC	-	-
<i>Phylloscopus trochiloides</i>	Greenish Warbler	LC	-	-
<i>Piaya cayana</i>	Common Squirrel-cuckoo	LC	-	-
<i>Picumnus cirratus</i>	White-barred Piculet	LC	-	-
<i>Picumnus pygmaeus</i>	Spotted Piculet	LC	-	-
<i>Picumnus temminckii</i>	Ochre-collared Piculet	LC	-	-
<i>Piranga flava</i>	Red Tanager	LC	-	-
<i>Pithys albifrons</i>	White-plumed Antbird	LC	-	-
<i>Ploceus bicolor</i>	Dark-backed Weaver	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Pogoniulus leucomystax</i>	Moustached Green Tinkerbird	LC	-	-
<i>Pogonocichla stellata</i>	White-starred Robin	LC	-	-
<i>Polioptila lembeyei</i>	Cuban Gnatcatcher	LC	-	-
<i>Pomatorhinus horsfieldii</i>	Indian Scimitar-babbler	LC	-	-
<i>Priotelus temnurus</i>	Cuban Trogon	LC	-	-
<i>Pselliophorus tibialis</i>	Yellow-thighed Finch	LC	-	-
<i>Pyriglena leucoptera</i>	White-shouldered Fire-eye	LC	-	-
<i>Quiscalus niger</i>	Greater Antillean Grackle	LC	-	-
<i>Rhipidura albicollis</i>	White-throated Fantail	LC	-	-
<i>Dumetia atriceps</i>	Dark-fronted Babbler	LC	-	-
<i>Rhynchocyclus olivaceus</i>	Eastern Olivaceous Flatbill	LC	-	-
<i>Coccyzus merlini</i>	Cuban Lizard-cuckoo	LC	-	-
<i>Sclerurus scansor</i>	Rufous-breasted Leaf-tosser	LC	-	-
<i>Scytalopus speluncae</i>	Mouse-colored Tapaculo	LC	-	-
<i>Schoeniparus castaneiceps</i>	Rufous-winged Fulvetta	LC	-	-
<i>Seiurus aurocapillus</i>	Ovenbird	LC	-	-
<i>Parkesia motacilla</i>	Louisiana Waterthrush	LC	-	-
<i>Parkesia noveboracensis</i>	Northern Waterthrush	LC	-	-
<i>Setophaga ruticilla</i>	American Redstart	LC	-	-
<i>Sheppardia sharpei</i>	Sharpe's Akalat	LC	-	-
<i>Sitta castanea</i>	Indian Nuthatch	LC	-	-
<i>Sitta frontalis</i>	Velvet-fronted Nuthatch	LC	-	-
<i>Sphyrapicus varius</i>	Yellow-bellied Sapsucker	LC	-	-
<i>Spindalis zena</i>	Western Spindalis	LC	-	-
<i>Spinus magellanicus</i>	Hooded Siskin	LC	-	-
<i>Stactolaema leucotis</i>	White-eared Barbet	LC	-	-
<i>Synallaxis azarae</i>	Azara's Spinetail	LC	-	-
<i>Synallaxis cinerascens</i>	Grey-bellied Spinetail	LC	-	-
<i>Synallaxis ruficapilla</i>	Rufous-capped Spinetail	LC	-	-
<i>Tachyphonus coronatus</i>	Ruby-crowned Tanager	LC	-	-
<i>Tangara cyanoventris</i>	Gilt-edged Tanager	LC	-	-
<i>Tangara icterocephala</i>	Silver-throated Tanager	LC	-	-
<i>Tangara schrankii</i>	Green-and-gold Tanager	LC	-	-
<i>Tangara seledon</i>	Green-headed Tanager	LC	-	-
<i>Tchagra australis</i>	Brown-crowned Tchagra	LC	-	-
<i>Terenotriccus erythrurus</i>	Ruddy-tailed Flycatcher	LC	-	-
<i>Terpsiphone batesi</i>	Bates's Paradise-flycatcher	LC	-	-
<i>Terpsiphone rufiventer</i>	Red-bellied Paradise-flycatcher	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat	Works
<i>Terpsiphone viridis</i>	African Paradise-flycatcher	LC	-	-
<i>Thalurania glaucopis</i>	Violet-capped Woodnymph	LC	-	-
<i>Thamnophilus ambiguus</i>	Sooretama Slaty Antshrike	LC	-	-
<i>Thamnophilus caerulescens</i>	Variable Antshrike	LC	-	-
<i>Thamnophilus murinus</i>	Mouse-colored Antshrike	LC	-	-
<i>Thraupis cyanocephala</i>	Blue-capped Tanager	LC	-	-
<i>Tangara episcopus</i>	Blue-grey Tanager	LC	-	-
<i>Tiaris olivacea</i>	Yellow-faced Grassquit	LC	-	-
<i>Tityra seminae</i>	Masked Tityra	LC	-	-
<i>Lophoceros camurus</i>	Dwarf Hornbill	LC	-	-
<i>Todus multicolor</i>	Cuban Tody	LC	-	-
<i>Troglodytes aedon</i>	House Wren	LC	-	-
<i>Trogon surrucura</i>	Southern Surucua Trogon	LC	-	-
<i>Trogon viridis</i>	Green-backed Trogon	LC	-	-
<i>Turdus abyssinicus</i>	Abyssinian Thrush	LC	-	-
<i>Turdus nigriceps</i>	Andean Slaty Thrush	LC	-	-
<i>Turdus plumbeus</i>	Northern Red-legged Thrush	LC	-	-
<i>Tyrannus caudifasciatus</i>	Loggerhead Kingbird	LC	-	-
<i>Tyrannus crassirostris</i>	Thick-billed Kingbird	LC	-	-
<i>Ochetorhynchus andaecola</i>	Rock Earthcreeper	LC	-	-
<i>Vermivora cyanoptera</i>	Blue-winged Warbler	LC	-	-
<i>Vireo altiloquus</i>	Black-whiskered Vireo	LC	-	-
<i>Vireo flavifrons</i>	Yellow-throated Vireo	LC	-	-
<i>Vireo griseus</i>	White-eyed Vireo	LC	-	-
<i>Vireo gundlachii</i>	Cuban Vireo	LC	-	-
<i>Vireo leucophrys</i>	Brown-capped Vireo	LC	-	-
<i>Vireo philadelphicus</i>	Philadelphia Vireo	LC	-	-
<i>Setophaga citrina</i>	Hooded Warbler	LC	-	-
<i>Xiphidiopicus percussus</i>	Cuban Green Woodpecker	LC	-	-
<i>Xiphocolaptes major</i>	Great Rufous Woodcreeper	LC	-	-
<i>Xiphorhynchus elegans</i>	Elegant Woodcreeper	LC	-	-
<i>Xiphorhynchus guttatus</i>	Buff-throated Woodcreeper	LC	-	-
<i>Xiphorhynchus obsoletus</i>	Striped Woodcreeper	LC	-	-
<i>Xiphorhynchus ocellatus</i>	Ocellated Woodcreeper	LC	-	-
<i>Xiphorhynchus spixii</i>	Spix's Woodcreeper	LC	-	-
<i>Xiphorhynchus pardalotus</i>	Chestnut-rumped Woodcreeper	LC	-	-
<i>Zenaida asiatica</i>	White-winged Dove	LC	-	-
<i>Zosterops ceylonensis</i>	Sri Lanka White-eye	LC	-	-

to be continued...

Species	English Name	IUCN	Habitat		Works
<i>Zosterops palpebrosus</i>	Indian White-eye	LC	-	-	
<i>Zosterops senegalensis</i>	African Yellow White-eye	LC	-	-	