



Wildlife Consumption Dynamics: Unveiling Conduru Park in Southern Bahia, Brazil

Joanison Vicente dos Santos Teixeira^{1,2*}, Wesley Duarte da Rocha²,
Jefferson Eduardo Silveira Miranda^{1,2} and Alexandre Schiavetti^{2,3}

ABSTRACT

The current investigation aimed to identify the wild animals utilized as a food source in five locations within the Serra do Conduru State Park region, Bahia, Brazil. The field survey was conducted from June 2016 to July 2017, involving semi-structured interviews and informal conversations with 45 hunters. The composition of species used for sustenance in the five locations was characterized through permutation multivariate analyses of variance. Generalized linear models were constructed to evaluate whether socio-demographic variables among hunters influenced the number of captured species. A total of 40 species was indicated like used for consumption in the region of the PESC. The species are classified to: one genus of Amphibia (one families and 1 order); 10 species and one genus of Birds (five families and five orders); 27 species and 2 genus of Mammals (19 families and 7 orders); and 2 species of Reptilia (two families and one order). *Dicotyles tajacu*, *Dasyurus novemcinctus*, and *Cuniculus paca* emerged as the most targeted species for food. Hunters who still reside within the conservation unit capture a greater number of wild animals. Older hunters and those with smaller family sizes hunt a broader range of species. The rifle and domestic dogs are the predominant techniques employed in the region. The findings underscore the persistence of illegal hunting practices in the Serra do Conduru State Park region. This emphasizes the necessity for measures directed at the conservation of hunted species, particularly those identified as being under some degree of threat in nature.

Keywords: Atlantic Forest; Hunted species; Ethnozoology.

1 Programa de Pós-graduação em Ecologia e Conservação da Biodiversidade, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16, Bairro Salobrinho, CEP 45662-900, Ilhéus, BA, Brazil.

2 Departamento de Ciências Agrárias e Ambientais, Laboratório de Etnoconservação e Áreas Protegidas (LECAP), Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16, Bairro Salobrinho, CEP 45662-900, Ilhéus, BA, Brazil.

3 Investigador asociado, CESIMAR/CENPAT, Puerto Madryn, Chubut, Argentina..

Corresponding author ✉. E-mail address: JVST (joanison1990@gmail.com)

SIGNIFICANCE STATEMENT

This study identifies the wild animal species most hunted and used as a food resource in the Serra do Conduru State Park (PESC) region, Bahia, Brazil and how this illegal activity can contribute to impact the reduction of wild animal species populations, including endangered species. Our results show the possible factors and/or motivations associated with the continuity of hunting in the region and discuss the need for actions aimed at the conservation of hunted species, such as the creation of extensive education programs for the dissemination of information on fauna conservation.

INTRODUCTION

The hunting of wild animals represents one of the oldest relationships between humans and biodiversity that is known (Alves and Souto 2010) and researchers have confirmed the importance of hunting, in various regions around the world, especially in tropical forests (Bodmer *et al.* 1996; Cullen Jr *et al.* 2001; Ojasti 1997, 2000; Peres and Nascimento 2006; Robinson and Redford 1991; Silva *et al.* 2022; Van Vliet *et al.* 2023). In many places, organized hunting of wild animals can bring considerable benefits to biodiversity conservation (Leader-Williams *et al.* 1996; Lewis and Alpert 1997; Lewis and Jackson 2005). In others, as hunted species are part of the diet of rural and urban populations or traditional communities (Castilho *et al.* 2018; Mesquita and Barreto 2015; Van Vliet *et al.* 2011, 2015, 2017) and are among the main sources of animal protein for traditional communities (e.g., indigenous, riverside communities; Alves *et al.* 2016; Faria and Malvasio 2018; Teixeira *et al.* 2020) being fundamental for the subsistence of the human population, especially those isolated (Fernandes-Ferreira and Alves 2014; Figueira *et al.* 2003).

The reasons for the growth in hunting activity in tropical forests are diverse, including increased human population, expansion of the highway network, the use of modern hunting instruments, occupation history, the preference for game meat, availability of game substitutes, socioeconomic conditions, the degree of dependence on hunting as a food source, and cultural conditions (Bennet and Robinson 2000; Bodmer *et al.* 1996; Constantino 2016; Constantino *et al.* 2008; Jerozolimski and Peres 2003; Schenck *et al.* 2006; Van Vliet and Mbazza 2011; Wright *et al.* 2007). However, although hunting is responsible for supplementing the diet of various traditional populations (Fernandes-Ferreira and Alves 2014; Silva-Neto 2016), the demand for game meat in the tropical region by traditional, urban and rural populations has been putting pressure on wild species, making it difficult to recover their numbers (Alves *et al.* 2009a; Alves *et al.* 2016; Pérez 2009; Ramos *et al.* 2016; Sousa and SrbeK Araújo 2017) and provoked several local extinctions (Canale *et al.* 2012).

Human consumption of hunted animals can lead to the extinction and/or population decline of various species, as well as a reduction in the mean body mass of animal populations as a result of selecting larger animals, diminishing future productivity of hunted populations (Alves and Rosa 2007a; Fernandes-Ferreira and Alves 2014, 2017; Ramos *et al.* 2016; Thiollay 2005; Thoisy *et al.* 2005). It should be highlighted that the effects of hunting are intensified by the loss and fragmentation of habitat, which increases the possibility of hunters accessing previously inaccessible ar-

reas, besides diminishing the occupation area of the species (Ramos *et al.* 2016). In contrast, some authors argue that hunting wild animals can be even more impactful than habitat loss (Cullen-Jr. *et al.* 2000).

Some communities have a great desire to hunt for consume meat, and this is a serious problem to Protected Areas (PA) (Castilho *et al.* 2018). A lot of factors can contribute for hunt around Protected Areas, as the household poverty and food security, and it is necessary to invest in environmental enforcement and assistance programs to communities (van Velden *et al.* 2020). To some organisms, like mammals, the PAs in tropical regions can be their diversity affected at 20% due to hunting, and aspects as poverty and inaccessibility to domestic meat increase this impacts (Benítez-López *et al.* 2019). In Brazil, hunt is strongly associated with absence of protein resources of domestic animals, and have a great importance to poor families, furthermore the cultural questions and socioeconomic factors have influence too (Alves *et al.* 2009a). This strengthens the necessity to study and understand the relations of social, cultural and economic aspects of hunting for food around of protected areas.

In the south of Bahia some Protected Areas are affected by illegal hunting, with people hunting a lot of species within and around these areas to consumption, and this situation include endangered species, which is very dangerous to these areas (Castilho *et al.* 2019). Some conflicts with the local fauna in the region are a problem to the species living in the Protected Areas of south Bahia, as hunt to consumption and the damage caused to some mammalian's species in the crops (Santos *et al.* 2020). The indiscriminate hunt of mammals in the region can provoke pressure in the local populations, and new politics need be established (Ribeiro, Schiavetti 2009).

In view of the negative impacts that hunting of wild animals bring, so that it is possible to preserve them, it is first necessary to know which species are most targeted by humans, how the catches are made and for what reasons the consumption of wild animals in the region of interest is an activity carried out by the locals (Alves and Rosa 2007b; Alves *et al.* 2009a; Alves *et al.* 2016). Furthermore, understanding the repercussions of hunting on species diversity, as highlighted by Oliveira and Calouro (2019), is crucial for developing targeted management alternatives. These alternatives should take into account both the local population and conservation efforts, particularly for species at risk of extinction. Therefore, the present study has as main objective to identify the species of wild animals most hunted for consumption in the region of the Serra do Conduru State Park (PESC), Bahia, Brazil. With this, it also seeks to quantify

the main techniques used to capture the species, the socioeconomic aspects associated with the continuity of hunting in the region. Finally, the study discusses the impacts of hunting on species conservation in the region. Another uses derived from the hunt were not considered in this work, because the focus is the hunt for food.

MATERIAL AND METHODS

Study area

The Serra do Conduru State Park ($39^{\circ} 05'32''$ W / $14^{\circ}26'17''$ S) (Figure 1) is a Protected Area (PA) preserving an area of 92.75 Km², which covers the municipalities of Ilhéus, Uruçuca, and Itacaré. This PA was created by State Decree no. 6,227 of 21st February 1997 to counterbalance the building of a highway (BA 001) (Bahia 2005). It is located within the Environmental Protection Area Itacaré-Serra Grande Coast (Bahia 2005). Both PAs include among their objectives the conservation of the remnants of Atlantic Forest and of the biodiversity of South Bahia (Bahia 2005). The PESC is categorized like an Integral Protection Area, with possibility to scientific

research, environmental education, leisure and environmental tourism, wherein to practice this activities depends to management authorization, and it is not allowed to live in the area (Brasil 2000). According to Schiavetti *et al.* (2012), illegal access to the PESC was indicated by the administration as a serious problem. The present study was carried out in five communities surrounding the PESC: District of Serra Grande (SG), District of Taboquinhas (DT), Nova Vida Settlement (NV), Camboinha Settlement (CS) and Community of Tesouras (CT) (Figure 1).

The SG ($39^{\circ}02'24''$ W / $14^{\circ}27'56''$ S) is a district of the municipality of Uruçuca, located in the south of Bahia. It has an estimated population of 3,585 inhabitants, 74% living in the urban area and 26% in the rural area (Instituto Ynamata 2008). The DT ($39^{\circ}10'21''$ W / $14^{\circ}21'31''$ S) is located 28 km from the Itacaré city hall (Instituto Ynamata 2008). Prior to tourism, its economy was based on the cultivation of cocoa, but with the decline of the crop and the advance of tourism the district began placing more value on its natural resources (Bahia 2005; Instituto Ynamata 2008). The AC ($39^{\circ} 2'45.06''$ W / $14^{\circ}20'45''$ S), located between the districts of Serra Grande and Itacaré, is made up of small rural producers dedicated



Figure 1. Location of the study area and the communities in which the fieldwork study was carried out with the hunters: Camboinha Settlement (CS), Community of Tesouras (CT), Nova Vida Settlement (NV), District of Serra Grande (SG) and District of Taboquinhas (DT).

to agriculture and local handicrafts (Bahia 2005; Instituto Ynamata 2008). The NV (39°09'04" W / 14°27'38" S) it is composed of 44 families of small farmers who were expropriated during the implementation of the PESC (Ribeiro and Schiavetti 2009). The CT (39° 5'24.21" W / 14°25'59" S) is composed of eight families who, due to slow landholding regularization, still reside within the PESC (Ribeiro and Schiavetti 2009; Sema 2005). In general, all families living around and into the PESC are not traditional population, but they are land squatters, farmers or farm workers (Bahia 2005).

Data collection

Data collection was carried out between June 2016 and July 2017. Only farmers that had also performed hunting activities in the region for at least two years, or that had already hunted for a minimum period of two years during their lives, participated in the study. The participants were interviewed only once. The hunters were selected using the criteria of "native specialists", who are those who recognize themselves as, and are recognized by the community as, culturally competent (Hays 1976), and through the snowball technique (Huntington 2000). These two techniques made it possible to interview a great number of the hunters identified in the region. Sampling was intentional and not random, whereby the interviewees were pre-defined (Albuquerque and Paiva 2004). All people were questioned about de hunting in the region of the PESC.

The information was obtained through open and semi-structured interviews, complemented by informal conversations (Huntington 2000) with questions related to the socio-demographic parameters of the interviewees (gender, origin, age, number of children, level of education, number of people in family, time living in the region and profession), the hunting strategies used (the people was free to cite the strategies) and the main common names of animals hunted and recognized by them. The "freelist" technique was used to record the names of species hunted for food (Albuquerque *et al.* 2008). This methodology considers that the more times the species appears on the list, the greater the cultural importance given to that species (Albuquerque *et al.* 2008; Quinlan 2005). To overcome the limitations of free listing were used: "nonspecific prompting", when the participants were questioned about other individuals or nonspecific groups, in order to remember more species; and "reading back", when the list cited by participants is read again slowly (Brewer 2002; Albuquerque *et al.* 2008).

To respect the rights of intellectual property, the following protocol were adopted in the field: we introduced ourselves before the interview, explaining the

nature and objectives of the study; and we then asked for permission to record the information. Before being interviewed, each participant received a Consent Form (TCLE) and a form authorizing the use of images, according to the rules established by Resolution no. 196 of the National Health Council of 10/1996 and approved by the Ethics Committee for Research with Human Beings (CEP) of the State University of Santa Cruz (UESC) (Certificate of Presentation of Ethical Appreciation: 61683516.2.0000.5526; Approval number of the project: 2.227.546).

Data analysis

The results were qualitatively analyzed according to the union model of diverse individual competencies, according to which, all the information referring to the researched subject is considered (Freitas *et al.* 1996). We considered the number of species cited by hunters like species richness, so we know the richness of each community. Species accumulation curves were constructed to verify whether the number of interviews was significant in relation to species of hunting importance that can be found in the study area. The number of species hunted was plotted as a function of the number of respondents, with 10,000 randomizations performed to generate a confidence interval (Colwell and Coddington 1994). We used the IUCN Red List of Threatened Species (IUCN 2022), Brazilian National List of Endangered Species (Brasil 2022) and Bahia List of Endangered Species (Bahia 2017) do analyze three scale of the conservation status of species hunted in the PESC region: global, national and regional.

Using a permutation multivariate analyses of variance (PERMANOVA, Anderson 2001), we evaluated differences between species cited at the five locations. We used Jaccard distance measures, and 5,000 permutations were generated. PERMANOVA is a permutation Understanding Analysis of Variance (ANOVA), which was developed to test the simultaneous response of one or more variables to one or more factors. PERMANOVA uses the "Adonis" procedure in the vegan package for R (Oksanen *et al.* 2013). We also used non-metric multidimensional scaling (NMDS) to represent the results of the PERMANOVA analyses. In the NMDS, we also used the Jaccard for the ordination of wild animal species composition used as a food source at the five locations. We realized a rarefaction curve with the iNEXT package (Chao *et al.* 2014; Hsieh *et al.* 2016), using the number of species and the communities to observe the difference between the diversity of communities. To observe the differences into species richness in the five communities studied, we use the Venn Diagram to indicate the number of species in each community, using the packages "ggVen-

nDiagram” (Gao *et al.* 2021).

Generalized linear models (GLMs) were built to assess whether the socio-demographic variables of the hunters affect the number of species cited to the hunters. The number of mentioned species were used as the response variable, and age of the hunters and socio-demographic were used as the explanatory variables. The models were subjected to an analysis of residuals to test the adequacy of the error distribution (Crawley 2012). The minimum adequate model (MAM) was obtained by extracting non-significant terms ($p < 0.05$) from the full model; when significant differences were observed between habitats, the data were submitted to contrast analysis by aggregating levels (Crawley 2012). If the level of aggregation was not significant and did not alter the deviance explained by the null model, the levels were pooled together (contrast analyses). All statistical analyses were conducted with R software (R Development Core Team 2022). We considered in all tests only the species cited to used like food by hunters, because this was the objective of the work.

RESULTS

A total of 45 hunters (44 men and one woman) from the five locations in the study area were interviewed. The number of interviewees in each location was: NV – 14; AC – 10; CT – 8; SG – 8; and DT – 5. Most of the interviewees were aged between 58 and 67 ($n = 11$ people; mean = 62.82; $sd = 2.82$), followed by 38 to 47 ($n = 9$ people; mean = 43.33; $sd = 3.60$), 68 to 77 ($n = 8$ people; mean = 70.75; $sd = 2.37$), 28 to 37 ($n = 6$ people; mean = 33.33; $sd = 3.14$), 48 to 57 ($n = 5$ people; mean = 53.40; $sd = 2.30$), 18 to 27 ($n = 3$ people; mean = 23.00; $sd = 4.58$), 78 to 87 ($n = 2$ people; mean = 82.00; $sd = 5.65$), and 88 to 97 (only one person with 90 years old). Most hunters are local residents born in the region (42.2%) or born in another location (57.8%), but who have lived there for several years. Regarding education, most of the interviewees completed only the first period of elementary school (1st to 4th grade; 42.2%), followed by those who admitted never having studied (37.8%). In terms of occupational activity, the majority are farmers (60%) and extractivists (22.2%) and, considering the number of children, most have three children (26,7%) or more than seven children (17,8%). Regarding the size of the family, most live alone (22,2%) or with another person (complementary material).

A total of 40 species was indicated like used for

consumption in the region of the PESC. The species are classified to: one genus of Amphibia (one families and 1 order); 10 species and one genus of Birds (five families and five orders); 27 species and 2 genus of Mammals (19 families and 7 orders); and 2 species of Reptilia (two families and one order). The species accumulation curves demonstrates complete stabilization (Figure 2), reaching the asymptote in approximately five interviews, indicating sampling efficiency in data collection.

The people indicated only one amphibian hunted for food, the frog (*Leptodactylus* sp.). Among the birds, the little tinamou (*Crypturellus soui*), the solitary tinamou (*Tinamus solitarius*) and the rusty-margined guan (*Penelope superciliaris*), had the highest number of citations. Among the mammals, the lowland paca (*Cuniculus paca*) is the species most indicated for hunting for food, followed by the collared peccary (*Dicotyles tajacu*) and the long-nosed armadillo (*Dasyppus novemcinctus*). The tegus (*Salvator merianae*) and the Boa (*Boa constrictor*) were the only species of reptile used for food (Table 1).

People indicated five species hunted for food and present in the three red lists, indicated here respectively with IUCN, Brazilian (BR) and Bahia (BA) status of conservation in parentheses: one bird, the *Crax blumenbachii* (IUCN = Endangered; BR = Endangered; BA = Critically Endangered); four mammals, the *Bradypus torquatus* (IUCN = Vulnerable; BR = Vulnerable; BA = Vulnerable), the *Sapajus xanthosternos* (IUCN = Critically Endangered; BR = Endangered; BA = Endangered), the *Callicebus melanochir* (IUCN = Vulnerable; BR = Vulnerable; BA = Vulnerable), and the *Chaetomys subspinosus* (IUCN = Vulnerable; BR = Vulnerable; BA = Vulnerable). So, was indicated *Penelope superciliaris* (Birds: Cracidae), present like “Near Threatened” in IUCN red list, *Tinamus solitarius* (Birds: Tinamidae) in “Near Threatened” and “Endangered” categories at IUCN and Bahia red list respectively, *Sylvilagus brasiliensis* (Mammalia: Leporidae) in the “Endangered” category of IUCN red list.

Hunters mentioned four hunting strategies: rifle hunting; hunt with a dog; trapping; and hunting with bait. CT was the only place where all 45 hunters used the four hunting strategies. Rifle hunting, the most common technique in the region, is used by 35 hunters; followed by hunting with dogs, traps and bait, used by 27, 21 and 11 interviewees, respectively. Most of the time, hunters use a combination of two or more hunting techniques, such as rifle hunting accompanied by dogs.




Figure 2. Species accumulation curve for the number of species hunted based on the number of respondents in the region of the Serra do Conduru State Park, Bahia, Brazil. The shaded area represents the confidence intervals of 95% based on 10,000 randomizations.




Figure 3. Non-metric multidimensional scaling (NMDS) analyses showing the dissimilarity of the composition of wild animal species used as a food source at the five locations studied in the region of the Serra do Conduru State Park, Bahia, Brazil. Each symbol represents a different location. Camboinha Settlement (CS), Comunidade Tesouras (CT), Nova Vida Settlement (NV), District of Serra Grande (SG), and District of Taboquinhas (DT). A significant difference in species composition captured for food ($p < 0.05$) among the five locations was observed in the PERMANOVA analysis. Data recorded between June/16 and June/17.

Table 1. List of wild animal species captured by the hunters in the region of the Serra do Conduru State Park, Bahia, Brazil. N^o: number of citations to hunt for food; IUCN: category in IUCN Red List; BR: category in Brazilian list of threatened species; SEMA: category in list of threatened species of Bahia.

CLASS	Order	Family	Scientific name	Vernacular name	N ^o	IUCN	BR	BA	
AMPHIBIA	Anura	Leptodactylidae	<i>Leptodactylus sp.</i>	Caçote	1	-	-	-	
BIRDS	Columbiformes	Columbidae	<i>Patagioenas cayennensis</i>	Pale-vented Pigeon	3	LC	-	-	
		Cracidae	<i>Penelope superciliaris</i>	Rusty-margined Guan	7	NT	-	-	
	Galliformes	Cracidae	<i>Ortalis araucuan</i>	East brasilian chachalaca	6	LC	-	-	
		Cracidae	<i>Crax blumenbachii</i>	Red-billed Curassow	4	EN	EN	CR	
	Passeriformes	Icteridae	<i>Psarocolius decumanus</i>	Crested Oropendola	1	LC	-	-	
	Piciformes	Ramphastidae	<i>Ramphastos sp.</i>	Toucan	1	-	-	-	
	Tinamiformes	Tinamidae	<i>Crypturellus soui</i>	Little Tinamou	8	LC	-	-	
		Tinamidae	<i>Tinamus solitarius</i>	Solitary tinamou	8	NT	-	EN	
		Tinamidae	<i>Rhynchotus rufescens</i>	Red-winged Tinamou	6	LC	-	-	
		Tinamidae	<i>Crypturellus parvirostris</i>	Small-billed Tinamou	6	LC	-	-	
MAMMALIA	Carnivora	Canidae	<i>Cerdocyon thous</i>	Crab-eating fox	21	LC	-	-	
		Felidae	<i>Leopardus sp.</i>	Leopard	2	-	-	-	
		Mustelidae	<i>Lontra longicaudis</i>	Neotropical otter	4	NT	-	VU	
		Mustelidae	<i>Eira barbara</i>	Tayra irara	5	LC	-	-	
		Procyonidae	<i>Nasua nasua</i>	South american coati	6	LC	-	-	
		Procyonidae	<i>Procyon cancrivorus</i>	Crab-eating raccoon	3	LC	-	-	
		Procyonidae	<i>Potos flavus</i>	Kinkajou	17	LC	-	-	
		Cetartiodactyla	Cervidae	<i>Mazama americana</i>	Red brocket deer	5	DD	-	-
			Cervidae	<i>Subulo gouazoubira</i>	Gray brocket deer	8	LC	-	-
Tayassuidae	<i>Dicotyles tajacu</i>		Collared peccary	38	LC	-	-		

CLASS	Order	Family	Scientific name	Vernacular name	Nº	IUCN	BR	BA
MAMMALIA	Cingulata	Dasypodidae	<i>Dasypus novemcinctus</i>	Nine-banded armadillo	26	LC	-	-
		Chlamyphoridae	<i>Euphractus sexcinctus</i>	Six-banded armadillo	22	LC	-	-
		Chlamyphoridae	<i>Cabassous tatouay</i>	Greater naked tailed armadillo	9	LC	-	-
	Didelphimorphia	Didelphidae	<i>Didelphis aurita</i>	Black-eared opossum	12	LC	-	-
	Lagomorpha	Leporidae	<i>Sylvilagus brasiliensis</i>	Tapiti	1	EN	-	-
	Pilosa	Bradypodidae	<i>Bradypus variegatus</i>	Brown-throated three-toed	1	LC	-	-
		Bradypodidae	<i>Bradypus torquatus</i>	Maned three-toed sloth	1	VU	VU	VU
		Myrmecophagidae	<i>Tamandua tetradactyla</i>	Southern tamandua	14	LC	-	-
		Cebidae	<i>Sapajus xanthosternos</i>	Buff-headed Capuchin	1	CR	EN	EN
		Pitheciidae	<i>Callicebus melanochir</i>	Coastal Black-handed Titi	2	VU	VU	VU
		Caviidae	<i>Hydrochoerus hydrochaeris</i>	Capybara	12	LC	-	-
		Cricetidae	<i>Oligoryzomys sp.</i>	Rodents	2	-	-	-
	Rodentia	Cricetidae	<i>Nectomys squamipes</i>	Water-rat	3	LC	-	-
		Cuniculidae	<i>Cuniculus paca</i>	Spotted paca	40	LC	-	-
		Dasyproctidae	<i>Dasyprocta leporina</i>	Agouti	4	LC	-	-
		Erethizontidae	<i>Coendou (Sphiggurus) insidiosus</i>	Bahia hairy dwarf porcupine	8	LC	-	-
		Erethizontidae	<i>Chaetomys subspinosus</i>	Bristle-spined Rat	5	VU	VU	VU
Boidae		<i>Boa constrictor</i>	Boa constrictor	12	LC	-	-	
REPTILIA	Squamata	Teiidae	<i>Salvator merianae</i>	Tegu	29	LC	-	-

The five locations were dissimilar (Permanova $r^2 = 0.18$; $p = 0.04$) in relation to species composition captured for food (Figure 3), and the rarefaction demonstrated that non stabilization curve for each community, with more divert to SG, NV, CS, CT and DT respectively (Figure 4). Ten species were cited exclusively at one location (Figure 5), three species were cited at two of the five locations, 11 were shared by three of the communities, five used for food were common to four of the communities and 12 species, such as the *C. paca* and the *D. novemcinctus* were of common use in the diet at five locations.

Some sociodemographic parameters are associated with the number of species used for food. The age of the hunters is positively related to the number of species hunted ($p = 0.02$; Table 2; Figure 6), hunters of a larger age group pointed a greater number of species. However, the families size is related to the number of species ($p = 0.01$; Table 2; Figure 7), that is, larger families capture fewer species. The average number of species hunted did not differ between sites (Deviance $4.40 = 71.53$; $p > 0.05$), but was significantly higher in the CT (Deviance $1.43 = 56.433$; $p = 0.01$; Figure 8). All 41 species used for food resource were cited by hunters CT, while 29, 23, 21 and 20 species are, respectively, food sources for hunters SG, NV, DT and AC. The other sociodemographic parameters analyzed were not related to the number of species used for feeding ($p > 0.05$; Table 2).

DISCUSSION

Our results demonstrate that hunting is a practice that still occurs in the region of the Serra do Conduru State Park. A great part of the wild species of the Atlantic Forest, mainly mammals, is exploited as a source of protein, which, according to hunters, is the main reason for hunting in the region. However, it should be noted that the slaughter of wild animals is not limited solely to the need for subsistence, but it is not our focus in this work. Through our results, it is evident the need for measures aimed at the conservation of hunted species, especially those that are under some degree of threat.

Although in different cultures women play an important role in hunting activities as support or hunting trips (Pereira and Schiavetti 2010; Ribeiro and Schiavetti 2009; Santos *et al.* 2018), we already expected the absence or low number of women identified as hunters. In the locations studied, only one woman identified herself as a hunter, demonstrating that this activity is predominantly performed by men. In other regions of Brazil occur the prevalence of men hunting too, like in Catimbau National Park, in Pernambuco state (Marques *et al.* 2022), National Forest Restinga of Cabedelo, Benjamim Maranhão Botanical Garden,

Engenho Gargaú Private Natural Heritage Reserve, Guaribas Biological Reserve (Barbosa *et al.* 2022) and in the north region of Brazil (Ramos *et al.* 2020).

The age group of the interviewees indicates that hunting is practiced by people of advanced age, which may be associated with the local culture, where the elderly are hunters. This is possibly due to the greater experience of these people with regard to hunting equipment and strategies for capturing these animals due to empirical knowledge about the behavioral characteristics of animals (Ayres and Ayres 1979). In other regions of northeastern Brazil the same happens, with older men involving with the hunting. Dantas-Aguiar *et al.* (2011) found more hunter with 30 to 40 year old and 52 to 62 year old in the community of Campos Formoso, Bahia. In the Catimbau National Park, Pernambuco, was found the same result, with hunters older, into 41 and 62 years old (Marques *et al.* 2022). A study carried out in four protected areas of Paraíba, the hunters were between 26 and 45 years old (Barbosa *et al.* 2022). Possibly the new generations are more access to education and information, and this result in the less interesting of younger in the hunt. So, this needs to evaluate in the news works. In some cases, the age influence on education level, and younger have the most education level (Barbosa *et al.* 2022). Usually hunters have less education level (Ramos *et al.* 2020; Marques *et al.* 2022), and the same result was found in our work. The relationship between the low level of education of hunters in the PESC region and hunting seems to be a common pattern observed in several other studies (Ribeiro and Schiavetti 2009; Castilho *et al.* 2018).

The hunted species play an important role in providing proteins to local families following a traditionally known pattern (Alves *et al.* 2009b, 2012; Barboza *et al.* 2016; Melo *et al.* 2014; Pereira and Schiavetti, 2010). Furthermore, the preference for mammals follows a pattern similar to several studies carried out in several neotropical biomes (Alves *et al.* 2009b; Mesquita and Barreto 2015; van Vliet *et al.* 2017), in the state of Bahia (Alves 2012; Dantas-Aguiar *et al.* 2011) and Southern Bahia (Castilho *et al.* 2018; Pereira and Schiavetti, 2010; Ribeiro and Schiavetti, 2009; Teixeira *et al.*, 2020). This preference is mainly due to the degree of difficulty of capture and the cost of hunting, flavor and yield of meat supplied by the animal (Cullen *et al.* 2001; Damaceno *et al.* 2019; Pereira and Schiavetti 2010).

Mammals are historically the animals most affected by hunting, due to their medium size and abundant population (Martínez and Garcia 2015; Trinca and Ferrari 2006; Ramos *et al.* 2020). Despite the dietary importance of hunting for the rural population, overexploitation of these species has serious consequences for the ecosystem (Alves and Souto 2010).




Figure 4. Rarefaction with the species diversity and sample coverage based on the hunter’s answers of five communities around the Serra do Conduru State Park, Bahia, Brazil. CS: Camboinha Settlement; CT: Community of Tesouras; DT: Districti of Taboquinhas; NV: Nova Vista Settlement; SG: Districti of Serra Grande.

In tropical forests, the abundance of wildlife is directly correlated with hunting patterns more than with other factors such as forest type, habitat size or protected area status (Pérez 2009). In the south of Bahia, Cas-

sano *et al.* (2012) found no traces of *Cuniculus paca*, *Mazama americana* or *Subulo gouazoubira* for example, this suggesting that hunting may have made these species disappear in some regions of the Bahia.




Figure 5. Venn diagram demonstrating the number of species in the communities around the Serra do Conduru State Park, Bahia, Brazil. CS: Camboinha Settlement; CT: Community of Tesouras; DT: Districti of Taboquinhas; NV: Nova Vista Settlement; SG: Districti of Serra Grande.

Table 2. Results of deviance of the minimal adequate model showing the effects of explanatory variables (socio-demographic parameters) on the number of species hunted (response variable). The error distribution used in the model was Quasi-Poisson. * represents statistical difference ($p < 0.05$).

Response variable	Explanatory variables	df	Deviance / <i>F</i>	<i>P</i>
Number of species hunted	Locality	4	71.530	0.03*
	Age	1	41.239	0.02*
	Gender	1	15.335	0.10
	Education level	4	36.585	0.20
	Profession	7	63.327	0.17
	Number of children	1	3.185	0.45
	Origin	1	4.220	0.38
	Time living in the region	1	2.461	0.50
	Family size	1	46.480	0.01*




Figure 6. Effect of hunters' ages on the number of species hunted for food in the region of the Serra do Conduru State Park, Bahia, Brazil. The error distribution used in the model was Quasi-Poisson.

As well as the concern regarding species directly affected by hunting, studies on the consequences of defaunation indicate that the ecology of forests used for hunting is severely disturbed (Harrisson 2011). The structure and dynamic of tropical forests are compromised due to the scarcity or extinction of herbivores, carnivores, and frugivores (Dirzo and Miranda 1991). The effects are related to the predation and dispersion

of seeds, herbivory, and increased seedling density and imbalances in the food chain of the ecosystems, such as increased density of small mammals due to a lack of predators and competitors (Alves and Souto 2010; Terborgh *et al.* 2001).

The number of citations indicated the food preference, and the *C. paca*, *D. novemcinctus* and *D. tajacu* were the most used species in local consump-




Figure 7. Effect of family size on the number of species hunted for food in the region of the Serra do Conduru State Park, Bahia, Brazil. The error distribution used in the model was Quasi-Poisson.




Figure 8. The average of species hunted for food observed for the five localities (CS - Camboinha Settlement, NV - Nova Vida Settlement, CT - Communitie Tesouras, SG - District of Serra Grande and DT - District of Taboquinhas) in the region of the Serra do Conduru State Park, Bahia, Brazil. The vertical bars correspond to the standard error (\pm se). Different letters above the columns represent statistically different means ($p < 0.05$)

tion. Other studies carried out in Bahia (Flesher and Laufer 2013), in South Bahia (Castilho *et al.* 2018;

Pereira and Schiavetti 2010), and in Brazil (Cullen Jr. *et al.* 2001; Fernandes-Ferreira and Alves 2014) also

indicate the lowland paca and armadillo species as the most consumed species and the most appreciated meats. Historical studies indicate that the lowland paca has suffered from hunting since the olden days (Cardim 1925; Varnhagen 1860; Silva 1898), hunted in various regions of Brazil (Canale *et al.* 2012; Freitas *et al.* 2005; Peres and Nascimento 2006) and indicated as the vertebrate species the most preferred mammal for hunting within the national territory (Fernandes-Ferreira and Alves 2014).

The *D. novemcinctus*, is valued in the region for its flavor and its ease of capture in comparison to other prey. The valorization of this species was also registered in several regions of Brazil (Hanazaki *et al.* 2009; Pereira and Schiavetti 2010; Rocha-Mendes 2005). According to the hunters in our study, the collared peccary, *D. tajacu*, is found throughout the region, making it easy to capture. According to Desbiez *et al.* (2011), the species has a gregarious habitat, also making it easier to capture and even kill more than one at a time. For Freitas *et al.* (2005), *D. tajacu* is widely pursued for eating as it has a high biomass, guaranteeing greater food resources, besides having a large percentage of body fat, a characteristic appreciated by the hunters. In the same region, Santos *et al.* (2020) identified the collared peccary as the most killed species as a result of the invasion, consumption, and/or destruction of agricultural crops, which is characterized by further motivation to slaughter the collared peccary.

The tegu, *Salvator merianae*, identified as the lizard of greatest dietary importance in the region of the PESC, was also considered by Alves *et al.* (2012) as the reptile of greatest hunting importance in a study carried out in two municipalities in Paraíba. Teixeira *et al.* (2020) indicated the tegus as the reptile most used for zootherapy in the region of the PESC. Fitzgerald (1994) states that lizards of this genus are also commonly hunted in Argentina, Paraguay, and parts of Bolivia.

The inclusion of hunted species on lists of endangered species represents a challenge to find forms of exploitation that minimize the impact on hunted species, and for this it is necessary to understand the context involving hunting practices (Alves 2012). The fact that endangered species are used as food source by hunters in the PESC region alerts us to the necessity for research that seeks to understand the degree of threat that these species are subjected to and the need for conservation strategies to minimize and revert such threats. Thus, it is necessary to understand the multidimensional context of hunting in the region, considering cultural and socioeconomic aspects, so that there is a reduction in pressure from hunting (Alves *et al.* 2016). To reduce the pressure of poaching is necessary considering cultural factors to guar-

antee the sustainability regional hunting, actions as educational programs, news components in environmental legislation, combating sport hunting and limitations on the use of hunting technologies, as firearms (Alves *et al.* 2009a).

Understand the hunting strategies is important to reduce the risk of many species, because some strategies are less selective and can increase the chance to kill endangered species, as strategies to trap with firearms and use of dogs (Oliveira and Calouro 2019; Santos *et al.* 2022; Yudha *et al.* 2022; Pattiselanno *et al.* 2023). Hunting strategies are fundamental to obtaining greater success in the search and killing of prey, and, consequently, reduce the expenditure of energy in the performance of hunts (Blasco *et al.* 2010; Costamagno *et al.* 2006). For Alves *et al.* (2009a), the variety of hunting strategies reflects the necessity to access the abundance of hunted animals living in different habitats. This diversity and choice of hunting strategy was also observed in the studied area. The hunters argue as to the greatest success in diversity and abundance of captured species when using different hunting techniques; for example, the use of a rifle with a dog, or the use of traps in areas when carrying out an active hunt with the support of a rifle and a domestic dog. The strategy is dependence of the prey, and hunters use the more efficient to kill more animals, generally they use firearms and dogs (Souza *et al.* 2022a), so the actively strategies offer more risk to the animals (Souza *et al.* 2022b).

The number of species indicated at the five locations may be related to the frequency of occurrence and the abundance of species in each location. According to Alves *et al.* (2009a), the hunting pattern of vertebrates for food can be influenced by availability and abundance of species in the region. Community Tesouras is within the PESC, an environmental protected area of integral protection, where the forest fragments are preserved to a greater degree and there is possibly a greater abundance and occurrence of species within its limits, which makes it possible for the hunters to have more hunting options. All the hunters from CT are extractivists who depend on natural resources to guarantee their survival and, possibly, invest more time in hunting activities as a way of acquiring a source of protein. It is worth highlighting that as an integral CU, hunting or any direct use of the natural resources is prohibited in the PESC (Brasil 2000).

The low species diversity in DT occur probably because this community is located far from the PESC. It is possible see the communities more proximity to the PESC present more species hunted. The PAs are more efficient to protected biodiversity and commonly are more richness and abundance of species (Gray *et al.* 2016). Non-protected areas are more exposed to

humans use for many reasons as economy, culture or recreation (Avigliano *et al.* 2019), and this may have influenced in our result, demonstrating more species cited by hunters in areas more close to PESC. So, the future works need evaluated this effect, and analyze the relationship between distance and biodiversity in this areas with methods to study the occurrence of fauna around this communities.

The positive relation between age and number of species cited by hunters was expected. Other works founded the same, with older hunter indicating more species (Barbosa *et al.* 2022; Oliveira and Calouro 2019). In some cases, the older hunter needs learning younger how the species behaving, and ecological aspects (Oliveira and Calouro 2019). With this, the tradition and knowledge are transferred for youngers, and de community maintains the tradition (Barbosa *et al.* 2022).

Larger families may prioritize the quality of the prey over quantity, resulting in a lower number of indicated species by hunters. Consequently, hunters with larger family sizes tend to prefer pursuing larger animals such as *Cuniculus paca*, *Dasypus novemcinctus*, *Dicotyles tajacu*, and *Euphractus sexcinctus*, as these species provide a greater quantity of meat. However, this preference poses a potential risk to endangered mammal species in the PESC region.

Several actions can be implemented aiming at the conservation of species involving the local population, in particular hunters, as the creation of extensive education programs for the dissemination of information on the conservation of endangered species, on the rules and legislation that regulate game activity. It is important to create channels of communication between academic institutions and government agencies with the population (Alves 2012), strengthening relations and allowing the greater involvement of populations in important decisions regarding the conservation of fauna in their communities. In this sense, it is important to argue about the need to include community participation in decisions involving the PESC management plan. In addition, research aimed at estimating the pressure of hunting and wild animal trafficking in the region should be encouraged.

CONCLUSION

Our study demonstrate that is necessary measures aimed at conservation of hunted species, especially for those found under some degree of threat. Its necessary understanding more reasons to hunting at the area of PESC and work mainly with families more dependent to the hunt. This will guide new proposals in the PESC. Families with older people and less scholar instructions need attention, because this may be a factor to increase the hunt. Work with news gen-

erations would be fundamental to better results about hunt, and this younger generation could contribute to reduce the hunt in the PESC.

As a strategic management plan involving the local population, particularly the hunters, creation of broad education programs for the dissemination of information on conservation of endangered species and the rules and legislation regulating hunting activity is recommended. Within conservation measures, it is necessary to consider the dietary importance of hunting for local families. The effects of hunting on the wild fauna are not easy to measure; further research is needed to estimate the pressure of hunting on the region and assess the impact of the activity on the local fauna. Furthermore, the environmental organs responsible for the conservation of protected areas should guarantee efficient monitoring, since hunting has been shown to be present in the studied areas.

ACKNOWLEDGMENT

We would like to thank the hunters from the five locations in the Serra do Conduru State Park for their trust and contributions during the research; the Post-Graduate Program in Zoology (PPGZOO) of the State University of Santa Cruz (UESC); the Coordination for the Improvement of Higher Education Personnel - Brazil (CAPES) - Finance Code 001, for the concession of a grant; CNPq for the productivity grant of the latter author; and the Iuri Dias, for map elaboration.

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

CONTRIBUTION STATEMENT

Conceived the idea presented: J.V.S.T.; A.S. Carried out the study: J.V.S.T. Carried out the data analysis: J.V.D.T.; W.D.R.; J.E.S.M. Wrote the first draft of the manuscript: J.V.S.T. Review and final writing of the manuscript: J.V.S.T.; A.S.; W.D.R.; J.E.S.M.

REFERENCES

- Albuquerque UP, Paiva RF (2004) **Métodos e Técnicas na Pesquisa Etnobotânica** Recife: Editora Livro Rápido/NUPEEA.
- Albuquerque UP, Lucena RP, Alencar NL (2008) **Métodos e técnicas para a Pesquisa Etnobotânica**. 2. ed. Recife: NUPEEA.
- Alves RRN (2012) **Relationships between fauna and people and the role of ethnozoology in animal conservation**. *Ethnobiology and Conservation* 1(2). doi: [10.15451/ec2012-8-1.2-1-69](https://doi.org/10.15451/ec2012-8-1.2-1-69).
- Alves RRN, Feijó A, Barboza RRD, Souto WMS, Fernandes-Ferreira H, Cordeiro-Estrela P, Langguth A (2016) **Game mammals of the Caatinga biome**. *Ethnobiology and Conservation*, 5(5). doi: [10.15451/ec2016-7-5.5-1-51](https://doi.org/10.15451/ec2016-7-5.5-1-51).
- Alves RRN, Mendonça LET, Confessor MVA, Vieira WLS, Lopez LCS (2009a) **Hunting strategies used in the semi-arid region of northeastern Brazil**. *Journal of Ethnobiology and Ethnomedicine*, 5(12). doi: [10.1186/1746-4269-5-12](https://doi.org/10.1186/1746-4269-5-12).
- Alves RRN, Neto NAL, Brooks SE, Albuquerque UP (2009b) **Commercialization of animal-derived remedies as complementary medicine in the semi-arid region of Northeastern Brazil**. *Journal of Ethnopharmacology*, 124(3): 600-608. doi: [10.1016/j.jep.2009.04.049](https://doi.org/10.1016/j.jep.2009.04.049).
- Alves RRN, Rosa IL (2007a) **Zootherapy goes to town: The use of animal-based remedies in urban areas of NE and N Brazil**. *Journal of Ethnopharmacology*, 113(3): 541-555. doi: [10.1016/j.jep.2007.07.015](https://doi.org/10.1016/j.jep.2007.07.015).
- Alves RRN, Rosa IL (2007b) **Biodiversity, traditional medicine and public health: where do they meet?** *Journal of Ethnobiology and Ethnomedicine*, 3(14). doi: [10.1186/1746-4269-3-14](https://doi.org/10.1186/1746-4269-3-14).
- Alves RRN, Souto WMS (2010) **Etnozoologia: conceitos, considerações históricas e importância**. In: Alves RRN, Souto WMS, Mourão JS. (Eds). *A Etnozoologia no Brasil: Importância, Status atual e Perspectivas*. Volume 7. 1 edition. Recife, PE, Brazil. NUPEEA.19-40.
- Alves RRN, Gonçalves MBR, Vieira WLS (2012) **Caça, uso e conservação de vertebrados no semiárido Brasileiro**. *Tropical Conservation Science*, 5: 396-416.
- Anderson MJ (2001) **A new method for non-parametric multivariate analysis of variance**. *Austral Ecology* 26: 32-46.
- Avigliano E, Rosso JJ, Lijtmae D, Ondarza P, Piacentini L, Izquierdo M, Cirigliano A, Romano G, Nuñez Bustos E, Porta A, Mabragna E, Grassi E, Palermo J, Bukowski B, Tubaro P, Schenone N (2019) **Biodiversity and threats in non-protected areas: A multidisciplinary and multi-taxa approach focused on the Atlantic Forest**. *Heliyon*, 5(8): e02292. doi: [10.1016/j.heliyon.2019.e02292](https://doi.org/10.1016/j.heliyon.2019.e02292).
- Ayres JM, Ayres C (1979) **Aspectos da caça no alto Rio Aripuanã**. *Acta Amazonica*, 9:287-298.
- Bahia (2017) Portaria nº 37 de 15 de agosto de 2017. **Torna pública a lista oficial das espécies da fauna ameaçadas de extinção do estado da Bahia**. Diário Oficial do Estado da Bahia. Access 19 october 2023.
- Bahia (2005) Secretaria Estadual de Meio Ambiente e Recursos Hídricos. **Plano de Manejo do Parque Estadual da Serra do Conduru**. Bahia.
- Barbosa JAA, Aguiar JO, Alves RRN (2022) **Hunting and wildlife use in protected areas of the Atlantic rainforest, northeastern Brazil, Desenvolvimento e Meio Ambiente**, 60: 249-270. doi: [10.5380/dma.v60i0.74388](https://doi.org/10.5380/dma.v60i0.74388).
- Barboza RR, Lopes S, Souto W, Fernandes-Ferreira H, Alves RRN (2016) **The role of game mammals as bushmeat in the Caatinga, northeast Brazil**. *Ecology and Society*, 21(2). doi: [10.5751/ES-08358-210202](https://doi.org/10.5751/ES-08358-210202).
- Bennett EL, Robinson JG (2000) **Hunting of Wildlife in tropical forests: Implications for biodiversity and forest peoples**. The World Bank Environment Department. *Biodiversity Series – Impact Studies*, N.76.
- Benítez-López A, Santini L, Schipper AM, Busana M, Huijbregts, AJ (2019) **Intact but empty forests? Patterns of hunting-induced mammal defaunation in the tropics**. *PLoS Biol*, 17(5):e3000247. doi: [10.1371/journal.pbio.3000247](https://doi.org/10.1371/journal.pbio.3000247).
- Blasco R, Fernández Peris J, Rosell J (2010) **Several different strategies for obtaining animal resources in the late Middle Pleistocene, The case of level XII at Bolomor Cave (Valencia, Spain)**. *Comptes Rendus Palevol*, 9(4): 171–184. doi: [10.1016/j.crpv.2010.05.004](https://doi.org/10.1016/j.crpv.2010.05.004).
- Bodmer RE, Puertas PE, Garcia JE, Dias DR, Reyes C (1996) **Game animals, palms and people of the flooded forests: management considerations for the Pacaya - Samiria National Reserve, Peru**. *New York Botanical Garden Press*, 13:217-231.
- Brasil (2000) Ministério do Meio Ambiente. **Lei no 9.985 de 18 de julho de 2000. Regulamenta o**

art. 225, § 1º, incisos I, II, III e VII da Constituição Federal, institui o Sistema Nacional de Unidades de Conservação da Natureza e dá outras providências. Diário Oficial da República Federativa do Brasil, Brasília, DF, July 18, 2000.

Brasil (2022) Ministério do Meio Ambiente. Portaria MMA nº 148, de 7 de junho de 2022. **Altera os Anexos da Portaria nº 443, de 17 de dezembro de 2014, da Portaria nº 444, de 17 de dezembro de 2014, e da Portaria nº 445, de 17 de dezembro de 2014, referentes à atualização da Lista Nacional de Espécies Ameaçadas de Extinção.** Diário Oficial da União, Brasília, 8 junho 2022.

Brewer DD (2002) **Supplementary interviewing techniques to maximize output in free listing tasks.** *Field Methods, Washington*, 14:108-118. doi: [10.1177/1525822X02014001007](https://doi.org/10.1177/1525822X02014001007).

Cassano CR, Barlow J, Pardini R (2012) **Large Mammals in na Agroforestry Mosaic in the Brazilian Atlantic Forest.** *Biotropica* 0(0):1-8. doi: [10.1111/j.1744-7429.2012.00870.x](https://doi.org/10.1111/j.1744-7429.2012.00870.x).

Canale GR, Peres AC, Guidorizzi CE, Gatto CAF, Kierulff MCM (2012) **Pervasive defaunation of forest remnants in a tropical biodiversity hotspot.** *PloS one*, 7(8): e41671. doi: [10.1371/journal.pone.0041671](https://doi.org/10.1371/journal.pone.0041671).

Cardim F (1925) **Tratado da terra e gente do Brasil.** Rio de Janeiro, J. Leite e Cia.

Castilho LC, Vleeschouwer KM, Milner-Gulland EJ, Schiavetti A (2018) **Attitudes and Behaviors of Rural Residents Toward Different Motivations for Hunting and Deforestation in Protected Areas of the Northeastern Atlantic Forest, Brazil.** *Tropical Conservation Science*, 11(1). doi: [10.1177/1940082917753507](https://doi.org/10.1177/1940082917753507).

Castilho L, De Vleeschouwer K, Milner-Gulland E, Schiavetti A (2019) **Hunting of mammal species in protected areas of the southern Bahian Atlantic Forest, Brazil.** *Oryx*, 53(4): 687-697. doi: [10.1017/S0030605317001247](https://doi.org/10.1017/S0030605317001247).

Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM (2014) **Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies.** *Ecological Monographs*, 84: 45-67. doi: [10.1890/13-0133.1](https://doi.org/10.1890/13-0133.1).

Colwell R, Coddington J (1994) **Estimating terrestrial biodiversity through extrapolation.** *Phil. Trans. Roy. Soc. Lond. B* 345(1311):101-118.

Constantino PAL (2016) **Deforestation and hunting effects on wildlife across Amazonian in-**

igenous lands. *Ecology and Society*, 21(2). doi: [10.5751/ES-08323-210203](https://doi.org/10.5751/ES-08323-210203).

Constantino PAL, Fortini LB, Kaxinawa AMK, Kaxinawa ES, Kaxinawa AP, Kaxinawa LS, Kaninawa JM, Kaxinawa JP (2008) **Indigenous collaborative research for wildlife management in Amazonia: the case of the Kaxinawá, Acre, Brazil.** *Biological Conservation*, 141(11): 2718-2719. doi: [10.1016/j.biocon.2008.08.008](https://doi.org/10.1016/j.biocon.2008.08.008).

Costamagno S, Liliane M, Cédric B, Bernard V, Bruno, M (2006) **Les Pradelles (Marillac-le-Franc, France), a mousterian reindeer hunting camp?** *Journal of Anthropological Archaeology*, 25: 466-484. doi: [10.1016/j.jaa.2006.03.008](https://doi.org/10.1016/j.jaa.2006.03.008).

Crawley MJ (2012) **The R Book, Second Edition.** Willey, New York.

Cullen-Jr L, Bodmer ER, Valladares-Padua C (2001) **Ecological consequences of hunting in Atlantic forest patches, São Paulo, Brazil.** *Oryx* 35:137-144. doi: [10.1046/j.1365-3008.2001.00163.x](https://doi.org/10.1046/j.1365-3008.2001.00163.x).

Cullen-Jr L, Bodmer RE, Valladares Pádua C (2000) **Effects of hunting in habitat fragments of the Atlantic forests, Brazil.** *Biological Conservation*, 95(1): 49-56. doi: [10.1016/S0006-3207\(00\)00011-2](https://doi.org/10.1016/S0006-3207(00)00011-2).

Damaceno AB, Ortega GP, Turci LCB (2019) **Uso da caça de subsistência no assentamento Santa Luzia, Cruzeiro do Sul, Acre.** *PUBVET*, 13(2):1-8. doi: [10.31533/pubvet.v13n2a262.1-8](https://doi.org/10.31533/pubvet.v13n2a262.1-8).

Dantas-Aguiar PR, Barreto RM, Santos-Fita D, Santos EB (2011) **Hunting Activities and Wild Fauna Use: A Profile of Queixo D'antas Community, Campo Formoso, Bahia, Brazil.** *Bioremediation, Biodiversity and Bioavailability*, 5(1):34-43.

Desbiez JAL, Keuroghlian UP, Bodmer RE (2011) **Invasive species and bushmeat hunting contributing to wildlife conservation, the case of feral pigs in a Neotropical wetland.** *Oryx*, 45(1): 78-83. doi: [10.1017/S0030605310001304](https://doi.org/10.1017/S0030605310001304).

Dirzo R, Miranda AM (1991) **Altered patterns of herbivory and diversity in the forest understory: a case study of the possible consequences of contemporary defaunation.** In: Wiley J. Plant-animal interactions: evolutionary ecology in tropical and temperate regions. New York.

Faria VA, Malvásio A (2018) **Aspectos sobre a caça, comercialização e consumo de quelônios na região do Corredor Ecológico Araguaia Bananal no Estado do Tocantins.** *Revista Ouricuri*, 8(2): 026-048. doi: [10.29327/ouricuri.v8.i2.a3](https://doi.org/10.29327/ouricuri.v8.i2.a3).

- Fernandes-Ferreira H, Alves RRN (2014) **Legislação e mídia envolvendo a caça de animais silvestres no Brasil, uma perspectiva histórica e socioambiental.** *Revista Gaia Scientia*, 8(1):1-7.
- Fernandes-Ferreira H, Alves RRN (2017) **The researches on the hunting in Brazil: a brief overview.** *Ethnobiology and Conservation*, 6(6). doi: [10.15451/ec2017-07-6.6-1-7](https://doi.org/10.15451/ec2017-07-6.6-1-7).
- Figueira MLOA, Carrer CRO, Silva-Neto PB (2003) **Weight gain and evolution of a wild white-lipped peccaries under extensive and semi-extensive systems, on a Savanna area.** *Revista Brasileira de Zootecnia*, 32:191-199. doi: [10.1590/s1516-35982003000100024](https://doi.org/10.1590/s1516-35982003000100024).
- Fitzgerald LA (1994) **Tupinambis lizards and people: a sustainable use approach to conservation and development.** *Conservation Biology*, 8:12-15.
- Flesher KM, Laufer J (2013) **Protecting wildlife in a heavily hunted biodiversity hotspot: a case study from the Atlantic Forest of Bahia, Brazil.** *Tropical Conservation Science*, 6:181-200. doi: [10.1177/194008291300600202](https://doi.org/10.1177/194008291300600202).
- Freitas H, Cunha jr. M, Moscarola J. (1996) **The content analysis: qualitative practical application in marketing.** Angra dos Reis/Brazil: Proceedings of ANPAD.
- Freitas FO, Moreira JR, Freitas JZF (2005) **Tradição cultural como diferenciador da dieta a base animal de duas etnias indígenas.** 1 ed. Embrapa Recursos Genéticos e Biotecnologia, Brasília, 9 p. (Comunicado Técnico 126).
- Gao CH, Yu G, Cai P (2021) **VennDiagram: An Intuitive, Easy-to-Use, and Highly Customizable R Package to Generate Venn Diagram.** *Frontiers in Genetics*, 12:706907. doi: [10.3389/fgene.2021.706907](https://doi.org/10.3389/fgene.2021.706907).
- Gray CL, Hill SLL, Newbold T, Hudson LN, Börger L, Contu S, Hoskins AJ, Ferrier S, Purvis A, Scharlemann JPW (2016) **Local biodiversity is higher inside than outside terrestrial protected areas worldwide.** *Nature Communications*, 7:12306. doi: [10.1038/ncomms12306](https://doi.org/10.1038/ncomms12306).
- Hanazaki N, Alves RRN, Begossi A (2009) **Hunting and use of terrestrial fauna used by Caicarás from the Atlantic Forest coast (Brazil).** *Journal of Ethnobiology and Ethnomedicine*, 5(1): 1-36. doi: [10.1186/1746-4269-5-36](https://doi.org/10.1186/1746-4269-5-36).
- Harrisson RD (2011) **Emptying the Forest: Hunting and the Extirpation of Wildlife from Tropical Nature Reserves.** *BioScience*, 61:919-924. doi: [10.1525/bio.2011.61.11.11](https://doi.org/10.1525/bio.2011.61.11.11).
- Hays TE (1976) **An Empirical Method for the Identification of Covert Categories in Ethnobiology.** *American Ethnologist* 3:489-507.
- Hsieh TC, Ma KH, Chao A (2016) **iNEXT: An R package for interpolation and extrapolation of species diversity (Hill numbers).** *Methods in Ecology and Evolution*, 7:1451-1456. doi: [10.1111/2041-210X.12613](https://doi.org/10.1111/2041-210X.12613).
- Huntington HP (2000) **Using Traditional Ecological Knowledge in Science: Methods and Applications.** *Ecological Applications*, 10:1270-1274. doi: [10.1890/1051-0761\(2000\)010\[1270:UTEKIS\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[1270:UTEKIS]2.0.CO;2).
- Instituto Ynamata (2008) **Instituto Floresta Viva. Diagnóstico Participativo de Serra Grande.** Serra Grande. 132p.
- IUCN. 2022. **The IUCN Red List of Threatened Species.** Version 2022-2. <https://www.iucnredlist.org>. Accessed on 19 october 2023.
- Jerolimski A, Peres CA (2003) **Bringing home the biggest bacon: a cross-site analysis of the structure of hunter-kill profiles in Neotropical forests.** *Biological Conservation*, 111: 415-425. doi: [10.1016/S0006-3207\(02\)00310-5](https://doi.org/10.1016/S0006-3207(02)00310-5).
- Leader-Williams N, Kayera JA, Overton GL. (Eds) (1996) **Tourist Hunting in Tanzania.** IUCN Species Survival Commission. Occasional Paper No. 14.
- Lewis DM, Alpert P (1997) **Trophy hunting and wildlife conservation in Zambia.** *Conservation Biology*, 11(1):59-68.
- Lewis D, Jackson J (2005) **Safari hunting and conservation on communal land in southern Africa.** In: Woodroffe R, Thirgood SJ, Rabinowitz A. (Eds) *People and Wildlife: Conflict or Coexistence?* Cambridge University Press, Cambridge, pp. 239-252.
- Marques AK, da Silva Novato T, de Albuquerque UP, Soldati GT (2022) **Can Socioeconomic Variables Influence Bird Hunting Activity in the Brazil's Semi-Arid Region?** *Human Ecology*, 50:515-530. doi: [10.1007/s10745-022-00330-8](https://doi.org/10.1007/s10745-022-00330-8).
- Martínez D, García D (2015) **Disentangling habitat use by frugivorous birds: Constant interactive effects of forest cover and fruit availability.** *Basic and Applied Ecology*, 16:460-468. doi: [10.1016/j.baae.2015.04.012](https://doi.org/10.1016/j.baae.2015.04.012).
- Melo RS, Silva O, Souto A, Alves R, Schiel N (2014) **The role of mammals in local com-**

- munities living in conservation areas in the Northeast of Brazil: an ethnozoological approach. *Tropical Conservation Science*, 7:423-439. doi: [10.1177/194008291400700305](https://doi.org/10.1177/194008291400700305).
- Mesquita GP, Barreto LN (2015) Evaluation of mammals hunting in indigenous and rural localities in Eastern Brazilian Amazon. *Ethnobiology and Conservation*, 4:114. doi: [10.15451/ec2015-1-4.2-1-14](https://doi.org/10.15451/ec2015-1-4.2-1-14).
- Ojasti J (1997) **Wildlife utilization in Latin America: current situation and prospects for sustainable management**. Food and Agriculture Organization of the United Nations, Rome.
- Ojasti J, Dallmeier F (2000) **Manejo de Fauna Silvestre Neotropical**. 1 ed. Smithsonian Institution/MAB Biodiversity Program, Washington D.C.
- Oksanen J et al. (2013) **vegan: Community Ecology Package**. <<https://cran.r-project.org/web/packages/vegan/index.html>>, Accessed in 06 October 2017.
- Oliveira MA, Calouro AM (2019) **Hunting agreements as a strategy for the conservation of species: the case of the Cazumbá-Iracema Extractive Reserve, state of Acre, Brazil**. *Oecologia Australis*, 23(2): 357-366.
- Pattiselanno, F., Janice, K. F., Krockenberger, A., Arobaya, A. Y. S., Sheil, D. (2023). **Hunting in Indonesian New Guinea: dogs, conservation and culture**. *Ethnobiology and Conservation*, 12: 1-11.
- Pereira JPR, Schiavetti A (2010) **Conhecimentos e usos da fauna cinegética pelos caçadores indígenas “Tupinambá de Olivença” (Bahia)**. *Biota Neotropica*, 10:175-183.
- Peres CA, Nascimento H (2006) **Impact of game hunting by the Kayapo of south-eastern Amazonia, Implications for wildlife conservation in tropical forest indigenous reserves**. *Biodiversity and Conservation*, 15:2627-2653. doi: [10.1007/s10531-005-5406-9](https://doi.org/10.1007/s10531-005-5406-9).
- Pérez JM (2009) **Parasites, Pests, and Pets in a Global World: New Perspectives and Challenges**. *Journal of Exotic Pet Medicine*, 18:248-253. doi: [10.1053/j.jepm.2009.09.003](https://doi.org/10.1053/j.jepm.2009.09.003).
- Quinlan M (2005) **Considerations for collectiong freelists in the field: Examples from ethnobotany**. *Field Methods*, 17(3): 219-234. doi: [10.1177/1525822X05277460](https://doi.org/10.1177/1525822X05277460).
- R Core Team (2022). **R: A language and environment for statistical computing**. R Foundation for Statistical Computing, Vienna, Austria. URL Available in <https://www.R-project.org>.
- Ramos CGS, Santos RB, Santos RWC, Oliveira MA (2020) **Hunting in a community of waste pickers of recyclable materials in Rondônia, Brazil**. *Revista Brasileira de Ciências da Amazônia*, 9: 4-15. doi: [10.47209/2317-5729.v.9.n.3.p.4-15](https://doi.org/10.47209/2317-5729.v.9.n.3.p.4-15).
- Ramos RM, Pezzuti JC, Vieira EM (2016) **Age structure of the Vulnerable whitelipped peccary *Tayassu pecari* in areas under different levels of hunting pressure in the Amazon Forest**. *Oryx*, 50:56-62. doi: [10.1017/S0030605314000350](https://doi.org/10.1017/S0030605314000350).
- Ribeiro GC, Schiavetti A (2009) **Conocimiento, Creencias y Utilización de la mastofauna por los pobladores del Parque Estatal de la Sierra de Conduru, Bahia, Brasil**. In: Costa-Neto EM, Santos-Fita D, Clavijo MV (Eds) *Manual de Etnozoología: Una guía teórico-práctica para investigar la interconexión del ser humano con los animales*. Tun-dra Ediciones, Valencia, España.
- Robinson JG, Redford KH (1991) **Neotropical wildlife use and conservation**. 1 ed. University of Chicago Press, Chicago.
- Rocha-Mendes F, Mikich SB, Bianconi GV, Pedro WA (2005) **Mamíferos do município de Fênix, Paraná, etnozootaxia e conservação**. *Revista Brasileira de Zoologia*, 22(4):991-1002. doi: [10.1590/S0101-81752005000400027](https://doi.org/10.1590/S0101-81752005000400027).
- Santos, SL, De la Fuente M F, Alves RRN (2022) **Patterns associated with hunting with dogs in a semiarid region of northeastern Brazil**. *Journal of Ethnobiology and Ethnomedicine* 18(1): 1-13.
- Santos JS, Teixeira JVS, Guanaes DHA, Rocha WD, Schiavetti A (2020) **Conflicts between humans and wild animals in and surrounding protected area (Bahia, Brazil): an ethnozoological approach**. *Ethnobiology and Conservation* 9(5). doi: [10.15451/10.15451/ec2020-02-9.05-1-22](https://doi.org/10.15451/10.15451/ec2020-02-9.05-1-22).
- Santos GP, Perilli ML, Cullem Jr L, Valladares-Padua C, Uezu A (2018) **Influência do Entorno de uma Unidade de Conservação sobre a Pressão de Caça: RPPN Estação Veracel como Estudo de Caso**. *Biodiversidade Brasileira*, 8(2): 219-231.
- Schenck M, Effa EN, Starkey M, Wikie D, Abernethy K, Telfer P, Godoy R, Treves A (2006) **Why People Eat Bushmeat: results from two-choice, taste tests in Gabon, Central Africa**. *Human Ecology*, 34:433-445. doi: [10.1007/s10745-006-9025-1](https://doi.org/10.1007/s10745-006-9025-1).
- Schiavetti A, Magro TC, Santos MS (2012) **Implementação das unidades de conservação do corredor central da Mata Atlântica no estado da Bahia: desafios e limites**. *Revista Árvore*, 36:611-623. doi: [10.1590/S0100-67622012000400004](https://doi.org/10.1590/S0100-67622012000400004).

- Silva AB, Pereyra PE, El Bizri HR, Souto WMS, Barboza RSL (2022). **Patterns of wildlife hunting and trade by local communities in eastern Amazonian floodplains.** *Ethnobiology and Conservation*, 11: 1-19
- Silva H (1898) **A Caça no Brasil**. Rio de Janeiro, Oficinas da Livraria Moderna.
- Silva-Neto AA, Ferreira PB, Torres RA, Texeira RHF, Duarte JMB, Barbosa AC, Vargas RC, Garcia JE (2016) **Diagnostic Cytochrome b gene profiles for the identification of paca (*Cuniculus paca*) bushmeat: implications for the monitoring of illegal hunting and wildlife trade.** *Brazilian Journal of Biology*, 76(1):55-58. doi: [10.1590/1519-6984.10814](https://doi.org/10.1590/1519-6984.10814).
- Sousa JAC, Srbek-Araujo AC (2017) **Are we headed towards the defaunation of the last large Atlantic Forest remnants? Poaching activities in one of the largest remnants of the Tabuleiro forests in southeastern Brazil.** *Environmental monitoring and assessment*, 189:197-206. doi: [10.1007/s10661-017-5854-1](https://doi.org/10.1007/s10661-017-5854-1).
- Souza JM, Landim AS, Ferreira FS (2022a) **A caça e fatores que influenciam o uso de espécies cinegéticas: uma revisão.** *Ethnoscintia*, 7(3):36-61. doi: [10.18542/ethnoscintia.v7i3.12075](https://doi.org/10.18542/ethnoscintia.v7i3.12075).
- Souza JM, Lins Neto EMF, Ferreira FS (2022b) **Influence of the sociodemographic profile of hunters on the knowledge and use of faunistic resources.** *Journal of Ethnobiology and Ethnomedicine*, 18(38). doi: [10.1186/s13002-022-00538-4](https://doi.org/10.1186/s13002-022-00538-4).
- Teixeira JVS, Santos JS, Guanaes DHA, Rocha WD, Schiavetti A (2020) **Uses of wild vertebrates in traditional medicine by farmers in the region surrounding the Serra do Conduru State Park (Bahia, Brazil).** *Biota Neotropica*, 20(1): e20190793. doi: [10.1590/1676-0611-BN-2019-0793](https://doi.org/10.1590/1676-0611-BN-2019-0793).
- Terborgh J, Lopez L, Nuñez P, Rao M, Shahabuddin G, Orihuela G, Riveros M, Ascanio R, Adler GH, Lambert TD, Balbas L (2001) **Ecological meltdown in predator free forest fragments.** *Science*, 294:1923-1926. doi: [10.1126/science.1064397](https://doi.org/10.1126/science.1064397).
- Thiollay J (2005) **Effects of hunting on guianan forest game birds.** *Biodiversity and Conservation*, 14:1121-1135. doi: [10.1007/s10531-004-8412-4](https://doi.org/10.1007/s10531-004-8412-4).
- Thoisy B, Renoux F, Juliot C (2005) **Hunting in northern French Guiana and its impacts on primate communities.** *Oryx*, 39:149-157. doi: [10.1017/S0030605305000384](https://doi.org/10.1017/S0030605305000384).
- Trinca CT, Ferrari SF (2006) **Caça em assenta-**
- mento rural na Amazônia matogrossense.** In: Jacobi P, Ferreira LC. (Orgs) *Diálogos em ambiente e sociedade no Brasil.* Annablume, São Paulo, p. 155-167.
- van Velden JL, Wilson K, Lindsey PA, McCallum H, Moyo BHZ, Biggs D (2020) **Bushmeat hunting and consumption is a pervasive issue in African savannahs: insights from four protected areas in Malawi.** *Biodiversity and Conservation*, 29:1443-1464. doi: [10.1007/s10531-020-01944-4](https://doi.org/10.1007/s10531-020-01944-4).
- Van Vliet N, Rovero F, Muhindo J, Nyumu J, Mbangale E, Nziavake, S, Cerutti P, Nasi R, Quintero S (2023) **Comparison of local ecological knowledge versus camera trapping to establish terrestrial wildlife baselines in community hunting territories within the Yangambi landscape in the Democratic Republic of Congo.** *Ethnobiology and Conservation*, 12: 1-14.
- Van Vliet N, Moreno J, Gomez J, Zhou W, Fa JE, Golden C, Alves RRA, Nasi R (2017) **Bushmeat and human health: Assessing the Evidence in tropical and sub-tropical forests.** *Ethnobiology and Conservation*, 6(3). doi: [10.15451/ec2017-04-6.3-1-45](https://doi.org/10.15451/ec2017-04-6.3-1-45).
- Van Vliet N, Mbazza P (2011) **Recognizing the multiple reasons for bushmeat consumption in urban areas: a necessary step toward the sustainable use of wildlife for food in Central Africa.** *Human Dimensions of Wildlife*, 16(1): 45-54. doi: [10.1080/10871209.2010.523924](https://doi.org/10.1080/10871209.2010.523924).
- Van Vliet N, Fa JE, Nasi R (2015) **Managing hunting under uncertainty: from one-off ecological indicators to resilience approaches in assessing the sustainability of bushmeat hunting.** *Ecology and Society*, 20(3). doi: [10.5751/ES-07669-200307](https://doi.org/10.5751/ES-07669-200307).
- Varnhagen FA (1860) **A caça no Brasil, ou, Manual do caçador, em toda a América tropical, acompanhado de um glossário dos termos usuas de caça.** Rio de Janeiro. Em casa de E. & H. Laemmert.
- Wright SJ, Hernandez A, Condit R (2007) **The bushmeat harvest alters seedling banks by favoring lianas, large seeds, and seeds dispersed by bats, birds and wind.** *Biotropica*, Washington, 39:363-371. doi: [10.1111/j.1744-7429.2007.00289.x](https://doi.org/10.1111/j.1744-7429.2007.00289.x).
- Yudha AP, Kusri MD, Arida E (2022) **Chasing for water monitors using dogs in West Java, Indonesia: a recreational hunting or pest control?** *Ethnobiology and Conservation*, 11:1-10.

Received: 01 June 2023
Accepted: 29 July 2023
Published: 05 January 2024
Editor: Rômulo Alves