



## Knowledge, use and traditional management of wildlife in the community of Zoquital, Morelos, Mexico

Mónica Bello-Román<sup>1</sup>, Alejandro García-Flores<sup>1\*</sup>, José Manuel Pino Moreno<sup>2</sup>

### ABSTRACT

Through their traditional productive activities, peasant and indigenous communities have generated knowledge on the use and management of wildlife to satisfy their needs for food, medicine or even to complement their basic food basket through the marketing of the species. The objective of this research was to analyze the knowledge, use and traditional management of wildlife in a rural community in the southwestern part of the state of Morelos, Mexico. The methodology included participant observation, semistructured interviews and guided tours and methods for obtaining and analyzing quantitative data “such as analysis of variance, a Mann–Whitney U test and  $X^2$  test”. The interviewees recognized a total of 57 species of wildlife, of which 22 are used as food, medicine, ornament, amulets, furs and agricultural tools in four traditional production units. In addition, eight species were reported in the oral tradition. Four hunting techniques and five weapons were documented, with the shotgun being the most commonly used. Two factors regulate hunting in the community: the closed season and religion. A total of 62,454 kg of useful biomass was recorded, with mammals being the group that contributed the most kilograms, followed by reptiles and birds. There were no significant differences in species consumption, across months or rainy and dry seasons due to factors regulating hunting. Based on the total value index of the species, *Odocoileus virginianus* was the most important for the inhabitants of the community.

**Keywords:** Subsistence Hunting, Traditional Ecological Knowledge, Ethnozoology, Uses, *Odocoileus Virginianus*.

<sup>1</sup> Biological Research Center, Universidad Autónoma del Estado de Morelos. Avenida Universidad 1001, Col. Chamilpa, C.P. 62209. Cuernavaca, Morelos, Mexico.

<sup>2</sup> Institute of Biology, Universidad Nacional Autónoma de México. Tercer Circuito Exterior s/n, Ciudad Universitaria, C.P. 04510, Coyoacán, Ciudad de México, Mexico.

\* Corresponding author ✉. E-mail address: MBR ([monica.bello@uaem.mx](mailto:monica.bello@uaem.mx)), AGF\* ([alejandrogarcia@uaem.mx](mailto:alejandrogarcia@uaem.mx)), JMPM ([jpino@ib.unam.mx](mailto:jpino@ib.unam.mx)).

## SIGNIFICANCE STATEMENT

Peasant communities possess traditional knowledge of the wildlife of the place where they live, which allows them to complement their basic needs, such as food and medicine, or to obtain a monetary income through the use and management of the species. The present study records the useful species in a rural community in the southwest of the state of Morelos, Mexico, where studies on wildlife are scarce and where in spite of community rules and prohibitions by the religion they profess, they still conserve knowledge about hunting techniques and tools. Family dynamics are related to the search for better income outside the community, and the registry of the species hunted in five family units over one year shows that the number of hunters has decreased; however, the inhabitants recognize the deer *Odocoileus virginianus* as the species with the highest value for food, economic and cultural use.

## INTRODUCTION

Wildlife is the main source of food self-sufficiency for the inhabitants of diverse regions of Mesoamerica, including Mexico, and they obtain tangible or intangible benefits from it as medicine, ornament, companion, amulet or climate indicators (Quijano and Calmé 2002; Montiel 2010; Monroy and García 2013; Gotoo 2016; Contreras and Yanguéz 2017; Dardón and Retana 2017; Asprilla and Díaz 2020; Zarazúa *et al.* 2020).

Indigenous and peasant groups, coexisting with wildlife in their immediate environment, have generated multiple dynamic interactions that vary according to each territory, historical period and culture of these societies. As part of these relationships, these traditional groups have accumulated deep traditional ecological knowledge about the animals with which they have interacted over time. This knowledge is characterized by being dynamic, holistic, adaptive and transmitted through spoken language in a generational way. It is derived from the systematic observation of nature, producing detailed and in-depth empirical information on the biological resources with which they are interrelated, such as use values, sexual differentiation, distribution, development and feeding habits of animals, as well as knowledge associated with classification, nomenclature and identification (Boege 2008; Toledo and Barrera 2008; Lira-Torres *et al.* 2014; Alves and Souto 2015; Velarde and Cruz 2015; Tamburini 2016).

However, territorial fragmentation caused by land use change and the lack of economic resources promote population movements in search of better living conditions, which are the factors that threaten this traditional knowledge (Monroy *et al.* 2011). Displacements of people from one place of residence to another imply relocation and loss of their territory of origin but encourage the search for identity to act and adapt to the new territory where they arrive (Pries 1999; Arévalo 2016), modifying their strategies and traditional knowledge to achieve their social welfare (UNESCO 2009); an example of this is the peasant community of Zoquital, Amacuzac, Morelos State,

Mexico that was created approximately 80 years ago by inhabitants of the state of Guerrero. The traditional knowledge recorded in this research is approached from the perspective of ethnozoology, a branch of ethnobiology that studies traditional zoological knowledge (CZT), i.e., how human groups conceive, classify and relate to animals (Costa-Neto *et al.* 2009). The objectives were: 1. To characterize the species with the value of use and exchange value recognized by the inhabitants. 2. To document hunting techniques, weapons, and strategies for regulating wildlife, and 3. To quantify the biomass of the species consumed by the local community.

## MATERIAL AND METHODS


### Study area

The peasant community of El Zoquital, municipality of Amacuzac, Morelos is located at coordinates 18° 32' 08"LN and 99° 21' 53"LO and at an altitude of 1117 masl (Figure 1). The climate is warm subhumid with summer rains, the average annual temperature and precipitation are 23.6°C and 1022.7 mm, respectively (Taboada *et al.* 2009), and the vegetation type is low deciduous forest and oak forest (Miranda and Hernández-X 2014). The total population is 152 inhabitants, and the degree of marginalization of the community is low (INEGI 2020).

### Qualitative research techniques

Initially, an approach was made to the community through previous visits to contact the local authority, who was informed about the objectives of the present work and the activities and techniques to be carried out, as suggested by the Code of Ethics of the Latin American Society of Ethnobiology (Cano-Contreras *et al.* 2016).

From December 2020 to December 2021, using the "snowball" technique or chain sampling (Taylor and Bogdan 1987; Castillo and Peña 2015), open and semistructured interviews were applied to 33 family units, representing 91.6% of the total households con-



**Figure 1.** Location of the study area.

sidering the disposition of local collaborators and that they were older than 18 years old. At all times, the health protocols indicated by the municipal assistant were followed for the safety of the families and the interviewer (wearing masks, keeping a distance, not kissing or hugging, conducting the interview in an open-air space, and avoiding the visit if any symptoms such as fever or cough were present). The information was recorded in a field logbook and included the local name of wildlife, its distribution in the territory, categories of use, forms of appropriation, beliefs, stories or legends.

In addition, the wildlife hunting events within the traditional productive units were recognized to obtain the number of individuals and weight of the species. A form was prepared to record the species hunted, the destination of the animals, the productive unit where they were obtained and their weight. To obtain the biomass, a Famsa mechanical scale with a capacity of 5 kg was used, and a Pretul Roman scale with a capacity of 50 kg was used for larger species. This form was completed during each visit to the study area with the information provided by four households that agreed to participate in the research. Wildlife referred to in the interviews was identified through guided walks and the use of field

guides for each taxonomic group (Peterson and Chaffiff 1989; Urbina and Morales 1994; Aranda 2013; Montalbán and Aréchaga 2014; Gaviño 2015; Pino *et al.* 2018a; Pino *et al.* 2018b). Scientific names were corroborated from databases such as The Reptile Database (<https://www.reptile-database.org>), Avibase- The World Bird Database (<https://avibase.bsc-eoc.org>) and ASM Mammal Diversity Database (<https://www.mammaldiversity.org>). The review of the conservation status of the species was carried out based on NOM- 059-SEMARNAT- 2010 (SEMARNAT 2010) and the International Union for Conservation of Nature (IUCN 2021).

## Analysis of the information

The information obtained from the interviews was systematized in a Microsoft Excel 360 database and analyzed using descriptive statistics.

To determine if there were significant differences in consumption between wildlife species across the months recorded, an analysis of variance was applied, and between the rainy and dry seasons with respect to biomass consumed, a Mann–Whitney U test was applied for independent samples. The normality and homoscedasticity of the data were previously checked.

ked (Infante and Zárate 2010). Finally, an  $X^2$  test was used to determine whether there were differences between wildlife capture sites. The analyses were processed in the R program (R Core Team 2021).

Finally, with the information provided by the local collaborators and the record of the biomass consumed during the study period, the total value of each wildlife species was calculated, which measures the importance of these species for a given community, using the formula proposed by Reyes-García *et al.* (2006):

$$Ve = CVe + PVe + EVe$$

where  $Ve$  = total value of species  $e$ ,  $CVe$  = cultural value of the species,  $PEve$  = practical value of the species, and  $EEve$  = economic value of the species. The  $CVe$  of each species was obtained from the following formula:

$$CVe = Uce \cdot Ice \cdot \sum IUce$$

where  $CVe$  = Cultural value of the species;  $Uce$  = total reported uses for species  $e$ /total uses of the species recorded in the present study;  $Ice$  = number of households that listed species  $e$  as useful/total households interviewed;  $UIce$  = number of households that mentioned each use of species  $e$ /total households interviewed.

The  $PEve$  of the species was obtained using the formula:

$$PEve = Upe \cdot Ipe \cdot DUpe$$

where  $PEve$  = practical value of the species;  $Upe$  = number of different uses observed for species  $e$  during field observations/total uses of the species recorded in this study;  $Ipe$  = number of times the species was brought to the household/total number of households that participated in the recording;  $DUpe$  = duration of each use/total number of field records. Reyes-García *et al.* (2006) assigned a duration of one day for species with food use and seven days for those used in medicine. In the present study, 15 days were considered for agricultural use and 30 days for ornamental and fur use.

The  $EEve$  of the species was obtained from the number of times the species was brought to the household multiplied by the price of the species. The price was referred to by the interviewees. For species for which the cost was not indicated, it was obtained by multiplying the average number of hours of capture by the price of the minimum wage for the community (\$172.87 M/N or 9.08 USD).

## RESULTS AND DISCUSSION

### Description of the reporting group

A total of 81.8% ( $n = 27$ ) of the family units were nuclear (consisting of father, mother and chil-

dren), and 18.2% ( $n = 6$ ) were extended (consisting of mothers-in-law, daughters-in-law, husbands and children). The average number of members per family unit was  $3.7 \pm 2.5$ , with an age range of 1 to 79 years, and the average length of residence in the community was 30.2 years. The distribution of activities within the family units allows them to complement their income and improve their wellbeing; thus, men are dedicated to rainfed agriculture, backyard livestock, commerce, salaried work and hunting as the main activities they carry out. Although hunting was mentioned by 22% of the households, only 16% continue to carry out this activity. Women take care of the household, collect plants for firewood or medicine, help with planting and take care of the livestock when the father of the family leaves the community and participate in the preparation of the hunted species; therefore, the form of family organization in the Zoquital includes the use of the labor force of its members: children, adolescents and adults (Aguado 1993; Solo de Zaldivar 1993; Román-Montes de Oca 2017); therefore, men and women possess and preserve knowledge about the wildlife species of their community. Ayala *et al.* (2019) call the distribution of these activities peasant specialties, where each member has specific tasks that are taught from the age of seven or eight, an age that coincides with the present research. According to local collaborators, boys participate by accompanying their parents or siblings in field activities, learning and recognizing animal species, while girls are instructed in food preparation by their mothers when the father of the family arrives home with them; in this way and as reported by Ríos (2020), girls and boys are integrated into the knowledge and wisdom of the family unit.

### Knowledge of species

The inhabitants of the community recognized 57 species of wildlife corresponding to 2 invertebrates, 1 amphibian, 14 reptiles, 22 birds and 18 mammals, belonging to 19 orders and 37 families (Table 1). The total number of species represents 16.33% of those reported for the Sierra de Huautla Biosphere Reserve (REBIOSH), 8.11% for the state of Morelos and 1.58% nationally (CONANP 2005; Sarukhán *et al.* 2017; Bustos and Castro 2020; Castro and Bustos 2020; Guerrero *et al.* 2020; Urbina 2020). According to the IUCN, 85.18% of the species (48.55 species) are in the status of least concern, while NOM-059- SEMARNAT-2010 records 14. 82%, (8 species) in some category of risk (see Table 1).

**Tabela 1:** Taxonomic relationship of wild animals recognized by the inhabitants of the community of Zoquital, Amacuzac, Morelos State, Mexico.

Order	Family	Common name	Use	Habitat	UICN	NOM-059-SEMARNAT-2010
Scientific name						
<b>Insects</b>						
Hemíptera	Pentatomidae					
	<i>Edessa</i> spp.	“Jumiles”	Al	Mo	LC	
Hymenoptera	Apidae					
	<i>Apis mellifera</i> Linnaeus, 1758	Bees	Al	Mo, Te	LC	
<b>Amphibians</b>						
Anura	Hylidae					
	<i>Agalychnis dacnicolor</i> (Cope, 1864)	Green frog		Ba	LC	
<b>Reptiles</b>						
Testudines	Kinosternidae					
	<i>Kinosternon integrum</i> Le Conte, 1854	Tortoise	Me	Ba	LC	Pr
Squamata	Iguanidae					
	<i>Ctenosaura pectinata</i> (Wiegmann, 1834)	Black iguana	Al, Me, Ve	Mo, Ba, Cu	LC	A
	Phrynosomatidae					
	<i>Phrynosoma taurus</i> Bocourt, 1870	Chameleon		Mo	LC	A
	<i>Sceloporus horridus</i> Wiegmann, 1834	“Chintete”		Mo	LC	
	Scincidae					
	<i>Plestiodon brevirostris</i> (Günter, 1860)	“Eslaboncillo”		Mo	LC	
	Teiidae					
	<i>Aspidoscelis</i> sp.	Lizard		Mo, Cu		
	Helodermatidae					
	<i>Heloderma horridum</i> (Wiegmann, 1829)	Scorpión		Mo	LC	A
Serpentes	Boidae					
	<i>Boa constrictor</i> Linnaeus, 1758	“Mazacuata”, “malacoa”, “coa”		Mo, Ba, Cu		A
	Colubridae					
	<i>Drymarchon menalunurus</i> (Duméril, Bibron & Duméril, 1854)	“Tilcuate”		Ba, Mo	LC	
	<i>Leptophis diplotropis</i> (Günter, 1872)	Green snake		Ba, Mo	LC	A
	<i>Masticophis mentovarius</i> (Duméril, Bibron & Duméril, 1854)	Ash snake		Mo	LC	
	<i>Oxibelis aeneus</i> Wagler, 1824	“Bejuquilla”		Mo	LC	

Order	Family	Common name	Use	Habitat	UICN	NOM-059-SEMARNAT-2010
Scientific name						
Elapidae	<i>Micrurus</i> sp.	“Coralillo”		Mo		
Viperidae	<i>Crotalus culminatus</i> Klauber, 1952	Rattlesnake	Me, Pe, Ve	Mo, Ba, Cu	NE	
<b>Birds</b>						
Galliformes						
Cracidae	<i>Ortalis poliocephala</i> (Wagler, 1830)	“Chachalaca”, “paita”	Al	Mo	LC	
Odontophoridae	<i>Philortyx phasiatus</i> (Gould, 1844)	“Codorniz, churrunda”	Al	Mo, Cu	LC	
Columbiformes						
Columbidae	<i>Columbina inca</i> (Lesson, 1847)	Turtle doves	Al	Mo, Cu, Ba, Pa	LC	
	<i>Leptotila verreauxi</i> Bonaparte, 1855	Pigeon	Al	Mo, Cu, Ba, Pa	LC	
	<i>Zenaida macroura</i> (Linnaeus, 1758)	“Huilotá”	Al	Mo, Cu, Ba, Pa	LC	
	<i>Zenaida asiatica</i> (Linnaeus, 1758)	White wing dove	Al	Mo, Cu, Ba, Pa	LC	
Cuculiformes						
Cuculidae	<i>Crotophaga sulcirostris</i> Swainson, 1827	“Garrapatero”, “tíjolo”		Mo, Cu, Ba, Pa	LC	
	<i>Geococcyx velox</i> (Wagner, 1836)	“Correcaminos”		Mo	LC	
Accipitriformes						
Cathartidae	<i>Coragyps atratus</i> (Bechstein, 1793)	Buzzard	Am	Mo, Cu	LC	
	<i>Cathartes aura</i> (Linnaeus, 1758)	“Aura”		Mo, Cu	LC	
Accipitridae	<i>Accipiter striatus</i> Vieillot, 1808	Hawk		Mo, Cu, Pa	LC	
	<i>Buteo jamaicensis</i> (Gmelin, JF, 1788)	Eagle		Mo, Cu, Pa	LC	
Strigiformes						
Strigidae	<i>Glaucidium brasilianum</i> (Gmelin, 1788)	“Cuacuana”		Mo	LC	
Coraciiformes						
Momotidae	<i>Momotus mexicanus</i> (Swainson, 1827)	“Pájaro bobo”		Mo	LC	

Order	Family	Common name	Use	Habitat	UICN	NOM-059-SEMARNAT-2010
<b>Order</b>						
<b>Family</b>						
<b>Scientific name</b>						
<hr/>						
Piciformes						
Picidae						
	<i>Melanerpes chrysogenys</i> (Vigors, 1839)	“La chica”, “carpintero”		Mo	LC	
Falconiformes						
Falconidae						
	<i>Herpetotheres cachinans</i> (Linnaeus, 1758)	“Guaco”		Mo	LC	
	<i>Caracara cheriway</i> (von Jacquin, 1784)	“Quebranta-huesos”		Mo	LC	
Passeriformes						
Corvidae						
	<i>Corvus corax</i> Linnaeus, 1758	Raven		Mo, Cu	LC	
Turdidae						
	<i>Turdus rufopalliatus</i> Lafresnaye, 1840	“Primavera”		Mo	LC	
Icteridae						
	<i>Icterus pustulatus</i> (Wagler, 1829)	“Calandrias”		Mo	LC	
	<i>Quiscalus mexicanus</i> (Gmelin, JF, 1788)	“Imes”, “zanate”		Mo	LC	
Tyrannidae						
	<i>Pitangus sulfuratus</i> (Linnaeus, 1766)	“Luis”		Mo	LC	
<hr/>						
<b>Mammals</b>						
<hr/>						
Artiodactyla						
Cervidae						
	<i>Odocoileus virginianus</i> (Zimmerman, 1780)	Whitetail deer, deer “cuernicabra”	Al, Me, Or, Esa, Pe, Ve	Mo	LC	
Tayassuidae						
	<i>Pecari tajacu</i> (Linnaeus, 1758)	Wild pig		Mo	LC	
Carnivora						
Canidae						
	<i>Canis latrans</i> Say, 1823	Coyote	Me, Am	Mo	LC	
	<i>Urocyon cinereoargenteus</i> (Schreber, 1775)	Fox		Mo	LC	
Felidae						
	<i>Herpailurus yagouaroundi</i> (Geoffroy Saint-Hilaire, 1803)	“Onza”		Mo	LC	A
	<i>Leopardus wiedii</i> (Schinz, 1821)	Margay		Mo	NT	P
	<i>Lynx rufus</i> (von Schreber, 1777)	Wild cat		Mo	LC	
Mephitidae						
	<i>Conepatus leuconotus</i> (Liechtenstein, 1832)	White-bandedskunk o “cadena”	Me, Ve	Mo	LC	
	<i>Spilogale gracilis</i> (Linnaeus, 1758)	Spottedskunk	Me, Ve	Mo	LC	
Procyonidae						
	<i>Nasua narica</i> (Linnaeus, 1775)	Badger	Al, Me, Ve	Mo	LC	
	<i>Procyon lotor</i> (Linnaeus, 1758)	Raccoon	Al	Mo	LC	

Order	Family	Common name	Use	Habitat	IUCN	NOM-059-SEMARNAT-2010
	Scientific name					
Cingulata						
	Dasypodidae					
	<i>Dasypus novemcinctus</i> Linnaeus, 1758	Armadillo	Al, Me, Or	Mo, Cu	LC	
Didelphimorphia						
	Didelphidae					
	<i>Didelphis virginiana</i> Kerr, 1792	Opossum	Me	Mo, Pa	LC	
Lagomorpha						
	Leporidae					
	<i>Lepus</i> sp.	Hare		Mo, Cu		
	<i>Silvylagus cunicularius</i> (Waterhouse, 1848)	Rabbit	Al, Or, Am	Mo, Cu	LC	
Rodentia						
	Muridae					
	<i>Peromyscus</i> sp.	Field rats		Mo		
	Sciuridae					
	<i>Sciurus aureogaster</i> Cuvier, 1829	Squirrel	Al	Mo, Cu	LC	
	<i>Otospermophilus variegatus</i> (Erxleben, 1777)	“Cuinique”		Mo, Cu	LC	

Use: Al= food, Me= medicinal, Or= ornamental, Am= amulet, Pe= fur, EsA= agricultural tool, Ve= sale; Ba= ravine, Mo= bush, Cu= crops, Pa= yard; IUCN (International Union for Conservation of Nature): LC= least concern, NE= not evaluated, NT= near threatened; NOM- 059- SEMARNAT- 2010: A= threatened, P= endangered, Pr= protected.

## Use of fauna

The community harvests 22 species of wildlife for different uses, which in order of importance are food (68%), medicine (45%), ornaments (18%), amulets (9%), furs (9%) and agricultural tools (4.5%). The number of species with use and sale is low when compared to other localities in the country such as Aguascalientes (Amador and De la Riva 2016), Campeche (Méndez and Montiel 2007) Chiapas (Gotoo 2016), Guerrero (Zavala *et al.* 2018), Morelos (García 2008; Monroy *et al.* 2008; Monroy and García 2013; García *et al.* 2020), Quintana Roo (Quijano and Calmé 2002) and the Tehuacán-Cuicatlán valley (Zarazúa *et al.* 2020), this may be due to two reasons: 1) 74% of the interviewees are neighborhood and when facing new conditions in their environment they have had to appropriate their territory (Arévalo 2016) by obtaining species through learning acquired locally or transmitted by parents and/or grandparents; and 2) the lack of economic resources within the community propitiates the father of the family to seek better job opportunities such as salaried work in the municipal capital or abroad, leaving the housewife in charge of the family and who does not go hunting, modifying the temporary or permanent organization of the family unit, a characteristic habit for Morelos farmers (Guzmán and León 2014).

Mammals were the taxonomic group that contributed the most species with use, followed by birds and reptiles. Similar results are reported by Asprilla and Díaz (2020) in Colombia, Tamburini (2016) in Argentina, García *et al.* (2020) in Morelos, Mexico and Contreras and Yáñez (2017) in Panama.

Food use was the most important, as it was highly mentioned by the inhabitants of the community, as in other regions of the country (Guerra *et al.* 2004; Monroy *et al.* 2008; Monroy-Vilchis *et al.* 2011; Barrasa 2012; Estrada *et al.* 2018). The part used for consumption is the meat, and the most common forms of preparation are roasted, fried or prepared in marinade, sauce and barbecue depending on family tastes. The insects *Edessa* spp. are consumed roasted or in sauce, and the honey of the bee *A. mellifera* is used for honey.

In addition, several authors report the current use of wildlife by indigenous and peasant communities to counter ailments and diseases related to the digestive, circulatory, respiratory, integumentary and cultural affiliation systems (Monroy and García 2013; Ramírez-Mella *et al.* 2016; Zavala 2018; García *et al.* 2019a; Valle *et al.* 2021). In the study area, reptiles are reported to use *Kinosternon integrum* to remove the "chipil" from children; the meat and blood of *Ctenosaura pectinata* to cure eyesight and anemia, as well as the skin of *Crotalus culminatus* that is placed on

the affected part to counteract the effect of the scorpion bite or stop nosebleeds, regarding mammals, the use of the burnt tail of *Didelphis virginiana* combined with other plants to cure empacho and its meat consumed together with that of *Spilogale gracilis* to cure asthma; the ground carapace of *Dasyppus novemcinctus* mixed with fat relieves sores caused by saddles in horses and the fat of *Nasua narica*, *D. novemcinctus*, *Canis latrans* and *Odocoileus virginianus* is used to relieve muscular pains while the consumption of meat of *S. gracilis* and *Conepatus leuconotus* prevents the appearance of pimples on the skin.

The paws of *Sylvilagus cunicularius* and the tusks of *C. latrans* were considered an amulet because they grant "good luck" to people; this protection is also mentioned by Monroy and García (2013) and Estrada *et al.* (2018). In addition, the head of *Coragyps atratus* is smeared on the nose of dogs to "make them crawl" and help in the capture of prey.

The skin of *C. culminatus*, the tail of *S. cunicularius*, the carapace of *D. novemcinctus* and the skin and antlers of *O. virginianus* have ornamental uses, while the bones of *O. virginianus* are used to trample the cob.

Forty-five percent of the species presented multiple uses; of these, *O. virginianus* had 6 uses, followed by *C. culminatus*, *C. pectinata*, *D. novemcinctus*, *S. cunicularius* and *N. narica*, with three uses each, as has also been reported for the community of Bonifacio García (García *et al.* 2020).

Species sold in the community or outside with family or friends were *Edessa* spp., *C. pectinata*, *C. culminatus*, *S. cunicularius*, *N. narica*, *M. macroura*, *S. gracilis*, *O. virginianus* and *A. mellifera* honey. Prices are assigned by the hunter and depend on the size of the organism; for example, *S. gracilis* is priced at \$100.00 per whole individual, while *O. virginianus* ranges from \$450.00 to \$600.00 per kilo or up to \$5,000.00 per whole animal. The price of *O. virginianus* meat is high compared to other places such as Campeche, Puebla or Tabasco, where the price ranges from \$85.00 to \$300.00 per kilo (Hernández-López *et al.* 2013; Estrada *et al.* 2018; Retana and Padilla 2018); however, as reported by Monroy-Vilchis *et al.* (2011), the flavor and quality of the meat and the time invested to capture the species are characteristics that families take into consideration when selling individuals or parts of individuals.

## Species reported in oral tradition

Species that due to their characteristics, habits, song or behavior were included in anecdotes transmitted from grandparents to parents or among the inhabitants and that are part of the intangible zoological heritage of indigenous and peasant communities

(Vargas 2009) are the following:

(a) **Reptiles**

*Drymarchon melanurus* "chicotea" (hits people with its tail) if it is found nearby"; this story is also reported for the Sierra de Montenegro, Morelos (Reyna 2013). *Phrynosoma taurus* is considered "the king of all animals because it carries a little crown on its head", while *Plestiodon brevirostris* "can cause death if grabbed". *C. culminatus* is referred to in the Bible, and interviewees report that "it will only bite from the heel down".

(b) **Ominous birds**

The song of *Crotophaga sulcirostris* announces rain, while the song of *Glaucidium brasilianum* and *Herpetotheres cachinans* portends misfortune or death for someone in the community, a belief documented in other regions of Morelos (Monroy and García 2013; García *et al.* 2019b; García-Flores *et al.* 2022), Guerrero (Zavala *et al.* 2018) and Aguascalientes, where nocturnal raptors are considered witches and to scare them away, one must shout profanities (Amador and De la Riva 2016).

(c) **Mammals**

"La piedra del venado" tells the story of a deer named "chalchihue" that was difficult to hunt because the hunters' shotguns would not shoot in front of it. It is said that one day they "cured" one of the guns and with it they managed to kill it. Later, when they cut up this animal, they found in the stomach a ball of hair, also known as "stone", which was put in a glass with water or blood and took with him to the hunt, and thanks to her, they caught deer easily. In southeastern Mexico, the "piedra" is called "Tunich Kéej" and serves as an amulet to hunt deer, but in the course of the day, it must be returned to the bush because otherwise it can be harmful to whoever possesses it (Quijano and Calmé 2002; Retana and Padilla 2018; Ríos 2020).

The inhabitants of Zoquital further related that when the cry of the fox *Urocyon cinereoargenteus* is heard, it is because "they announce that a girl from the community is going to leave with her boyfriend". The situation is different in the state of Guerrero, where the appearance of this species in front of deer hunters portends a bad hunt (Zavala *et al.* 2018).

## Hunting techniques

Hunting techniques in order of importance by frequency of mention were muzzleloading, champering, spying, and lamping. To obtain *O. virginianus*, the "arriada" is used, by which friends and/or family members gather and head toward the bush and once they arrive at the selected place, the work is divided: those who carry guns are located in a specific place while the rest of the hunters advance in the opposite direction and with the support of dogs, they begin to make noise to direct the deer to where those who carry guns are waiting and once the animal is identified, they proceed to hunt it. This method allows the division of labor, coexistence and participation of all members of the family unit from the organization of the tools to the preparation of the species. This technique is also known as "acorralar" (García 2008), "búsqueda" (Bardales *et al.* 2017) or batida (Plata 2017; Retana and Padilla 2018). In addition, it has been reported in other states of Mexico, such as Guerrero, Morelos and Yucatán (Montiel 2010; Velarde and Cruz 2015; López *et al.* 2018).

The "campeada" refers to the direct encounter with wildlife while performing activities such as planting, grazing or collecting firewood, mushrooms or edible and/or medicinal plants; it is also known as opportunistic (Lira-Torres *et al.* 2014), surprise or excursion (Retana and Padilla 2018), chance encounter (Bardales *et al.* 2017) or diurnal routes (Gotoo 2016). This is common for hunting *C. pectinata*, *S. cunicularius*, *C. culminatus* and/or *C. leuconotus*.

The spying technique consists of identifying places where it is known that animals will arrive for food or water; they then place themselves in nearby trees (e.g., *Ficus* sp. amates) and wait for the animal to approach the site. This is mainly used for hunting *O. virginianus* and *N. narica* and is also reported in central Morelos (García 2008), southeastern Mexico (Quijano and Calmé 2002; Gotoo 2016; Retana and Padilla 2018) and Peru (Bardales *et al.* 2017). This suggests that the inhabitants of the communities possess knowledge about the feeding habits of local wildlife, which gives them an advantage in gaining easier access to prey by recognizing the spaces used by the species to meet their biological needs.

The "lampareada" is recognized by the inhabitants as "campeada de noche" (night hunting), and species such as *D. novemcinctus*, *O. virginianus* and *N. narica* are sought. The "lampareo" starts at dusk and can last until the following day depending on the species and has also been reported for communities in Chiapas (Gotoo 2016), Guerrero (López *et al.* 2018) and Tabasco (Hernández-López *et al.* 2013).

The average time spent capturing animals depends on the number of participants, distance and accessibi-

lity. For *Edessa* spp. For example, an average of nine hours is needed, from 8 to 18 hours for *O. virginianus* and from approximately 10 minutes to 3 hours for the rest of the animals.

## Hunting weapons

The weapons used are shotguns, "dogs," slingshots, machetes, and "cuaxtlera"; the latter is a type of hand-made shotgun that uses gunpowder and ixtle blocks. Although the use of the shotgun implies the purchase of ammunition and/or cartridges, it allows for obtaining species easily and quickly, and such characteristics make this instrument widely used in communities in the state of Morelos (García 2008; Velarde and Cruz 2015), Campeche, Chiapas, Guerrero, Oaxaca, Puebla and Veracruz, (Lira-Torres *et al.* 2014; Gotoo 2016; Estrada *et al.* 2018; López *et al.* 2018; Retana and Padilla 2018) and Argentina, Panama and Peru (Tamburini 2016; Bardales *et al.* 2017; Contreras and Yanguez 2017).

Dogs accompany hunters during the development of their activities when they go alone or in groups, and their function is to follow the trail of the species and chase wounded prey until they are located; therefore, they provide benefit not only by guarding the home but also by obtaining wildlife meat useful for the inhabitants of the community. In Campeche, depending on their performance, they can be considered leader or master dogs or else, support or secretary dogs (Plata 2017).

## Regulation of species harvesting


The hunting season is from December 21 to February 21. However, when species such as *N. narica*, *D. virginiana* or *C. latrans* threaten crops or backyard animals, they are hunted regardless of the closed season. The rest of the animals are captured when they are accidentally encountered while tending crops, tending livestock or collecting firewood. In the municipality of Pasto, Colombia (Martínez-Ceballos 2014), the productive activities carried out by the inhabitants allow them to cover their food needs, so the consumption of wildlife is null; however, in the Zoquital, despite being dedicated to agricultural activities, if a species is obtained, the family decides whether to consume it, give it to another family or sell it to complement their basic food basket, which indicates the importance of this resource in food and the economy.

In addition, 40% of the interviewees belong to the Israelite religion, who follow the "law of food" reported and explained in the Bible in the book Leviticus 1, 11; it indicates the type of animals they are allowed to consume according to certain characteristics, for example, "the animal with a split hoof, cleft in two

nails and that a ruminating. You will eat those who have fins and scales either live well in the sea or in the rivers, in the case of four-legged insects, will you be able to eat those that, in addition to their four legs, have legs to jump on the ground. Animals that do not have these characteristics are considered "unclean" according to their beliefs and therefore are not hunted in the community. However use them and about hunting techniques; in this way, the traditional knowledge about the species with use continues to be maintained in the Zoquital in comparison to the Chenes region in Campeche, where religion prohibits the use of wildlife as a means of healing and therefore there is a loss of traditional medicinal knowledge (Dardón and Retana 2017).

## Species captured


Seventy captured individuals corresponding to 11 species were recorded. The total biomass extracted was 116.34 kg, and the biomass consumed was 62.454 kg. Of the reptiles, 11,525 kg were consumed, while birds and mammals contributed 1,804 kg and 49,125 kg, respectively. The total biomass consumed during one year was lower compared to the Chaco Seco in Argentina, where species consumption was 1,483 kg (Tamburini 2016). This difference may be because in that region, only mammals were hunted, whose biomass is higher compared to other vertebrate groups, while in the Zoquital, the number of species consumed was five mammals, four birds and two reptiles. Lira-Torres *et al.* (2014) reported 1 900 kg of biomass extracted from mammals in a community in the Zoque jungle, while Montiel (2010) reported 4.8 tons in localities of the Yucatan Peninsula from the same group of wild vertebrates, which reflects that communities consider other taxonomic groups according to the region and habitat where they are located to supplement their basic needs. According to the statistical analysis, there were no significant differences in consumption between species ( $H = 3.12$ ,  $p = 0.7539$ ). The species that contributed the most biomass was *N. narica* at 24.575 kg, followed by *O. virginianus* (19.5 kg) and *C. pectinata* (7.925 kg) (Figure 2). Mammals by size and weight were the taxonomic group that contributed the most biomass in the Zoquital, which has also been reported in Tabasco (Hernández-López *et al.* 2013), Colombia (Martínez-Ceballos 2014) and Peru (Bardales *et al.* 2017; Costa *et al.* 2018). Although there were no significant differences in species consumption across months ( $H = 5.86$ ,  $p = 0.4012$ ), March was the month in which the most biomass was obtained (16.925 kg), of which 88% came from *N. narica*; on the other hand, no species were caught in April (Figure 3). Likewise, there were no significant differences in consumption between the rainy and dry



**Figure 2.** Wildlife biomass consumed by species for the inhabitants of the community of Zoquital, Amacuzac, Morelos State, Mexico.

seasons ( $W = 4658.00$ ,  $p = 0.4012$ ) characteristic of the low deciduous forest, which suggests that the capture of organisms occurs occasionally while activities are carried out in the traditional productive units regardless of the months or times of the year or because

they caused some damage to the crops. According to Tamburini (2016), hunting is practiced for three main objectives: obtaining meat for food, for causing damage and to obtain income from them, which coincides in the Zoquital community, since 40% of the



**Figure 3.** Wildlife biomass consumed per month for the inhabitants of the community of Zoquital, Amacuzac, Morelos State, Mexico.

species captured were obtained for food while 60% had an exchange value through sale (54%) or as a gift (6%).

## Hunting areas

Informants identified four traditional productive units where wildlife species are distributed: the forest, the ravine, crops or plots of land, and backyards or family gardens. Species can be found in more than one site (see Table 1) because of their ability to move in search of food, perch or to reproduce; these characteristics are considered by hunters when looking for a particular species.

Of the sites identified, during the field work, the capture of the species was recorded in the bush, which contributed 50.924 kg of biomass, followed by the cornfield with 10.8 kg and the pasture with 0.7 kg. The bush is the main space where appropriation takes place, as it is the natural environment inhabited by the species. López *et al.* (2018) identified 11 hunting spaces in Guerrero, which indicates that the inhabitants travel long distances and possess knowledge of their environment that allows them to obtain bush meat (Aldana *et al.* 2016; Tamburini *et al.* 2016; Estrada *et al.* 2018; López *et al.* 2018; García *et al.* 2020).

## Total value of species

Of the 22 species harvested by the community, *O. virginianus* is the most important (see Table 2). According to Martínez-Ceballos (2014), conservation processes, environmental education and income derived from productive activities influence the low values of species use, which coincides with the present work because those in charge of hunting animals and seeking income to meet family expenses are men who migrate, and therefore, hunting decreases. For example, the price referred to by the interviewees for *O. virginianus* was approximately \$5,000.00 per animal, compared to \$32.41 for the birds, which was calculated based on the average time invested in capturing them. The practical and economic value was obtained from the species that were recorded during the study period, and only the animals referred to by the participating families were taken into account; however, *O. virginianus* was the most culturally, practically and economically important species for the inhabitants of Zoquital, as has been documented for communities in central, southern and southeastern Mexico (Velarde and Cruz 2015; Ramírez-Mella *et al.* 2016; Estrada *et al.* 2018; López *et al.* 2018; Retana and Padilla 2018; García *et al.* 2020; Montiel 2010).

## CONCLUSION

The relationship that the inhabitants of the Zoquital community have with wildlife has allowed them to meet their basic needs through direct use or by generating income from the sale of animals. Although the use of the species is regulated by community and religious rules, the inhabitants of the community still maintain their traditional knowledge about hunting techniques and weapons as well as the recognition of spaces where the animals carry out their activities, allowing them to obtain the species in diverse productive units such as the bush, yard, pasture or cornfield, the latter being the space where the wild fauna that competes for the crops is hunted. In addition to the information provided by the local collaborators, studies of the species used in the community should be carried out, which will allow us to know the current situation of the populations in use, as well as to generate strategies for their management, conservation and exploitation without losing the traditional knowledge of the species in the Zoquital.

## ACKNOWLEDGMENT

The first author thanks the National Council of Science and Technology for the scholarship granted during her Master's studies, which allowed the development of this research. The authors thank the inhabitants of the community of El Zoquital for their hospitality and support during the field work.

## DATA AVAILABILITY

The data used to support the conclusions of this study are available from the corresponding author upon request.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## CONTRIBUTION STATEMENT

Conception of the idea presented: MBR, AGF.

Conduct of field work: MBR.

Conducting data analysis: MBR, AGF.

Drafting of the manuscript: MBR, AGF, JMPM.

Review and final drafting of the manuscript: MBR, AGF, JMPM.

Supervision: AGF.

**Tabela 2.** Value of wildlife species in the community of Zoquital, Amacuzac, Morelos, Mexico.

Scientific name	Common name	CVe	PVe	EVe	Ve
<i>Odocoileus virginianus</i>	Whitetail deer,deer “cuernicabra”	0.781	0.465	1000	10001.245
<i>Crotalus culminatus</i>	Rattlesnake	0.004	0.167	2400	2400.170
<i>Conepatus leuconotus</i>	White-bandedskunk o “cadena”	0.001	0.000	1500	1500.001
<i>Nasua narica</i>	Badger	0.006	0.032	1250	1250.038
<i>Ctenosaura pectinata</i>	Black iguana	0.004	0.000	750	750.004
<i>Spilogale gracilis</i>	SpottedSkunk	0.001	0.009	500	500.010
<i>Silvylagus cunicularius</i>	Rabbit	0.006	0.235	200	200.240
<i>Columbina inca</i>	Turtle doves	0.074	0.032	162.05	162.156
<i>Zenaida macroura</i>	“Huilota”	0.061	0.011	97.23	97.303
<i>Philortyx fasciatus</i>	“Codorniz, churrunda”	0.006	0.001	32.41	32.417
<i>Leptotila verreauxi</i>	Pigeon	0.004	0.001	32.41	32.415
<i>Dasypus novemcinctus</i>	Armadillo	0.019	0.000	0.000	0.019
<i>Canis latrans</i>	Coyote	0.004	0.000	0.000	0.004
<i>Edessa spp.</i>	“Jumiles”	0.001	0.000	0.000	0.001
<i>Apis mellifera</i>	Bees	0.001	0.000	0.000	0.001
<i>Zenaida asiatica</i>	White wing Dove	0.001	0.000	0.000	0.001
<i>Kinosternon integrum</i>	Tortoise	0.0006	0.000	0.000	0.0006
<i>Ortalis poliocephala</i>	“Chachalaca”, “paita”	0.0006	0.000	0.000	0.0006
<i>Coragyps atratus</i>	Buzzard	0.0002	0.000	0.000	0.0002
<i>Didelphis virginiana</i>	Opossum	0.0002	0.000	0.000	0.0002
<i>Sciurus aureogaster</i>	Squirrel	0.0002	0.000	0.000	0.0002
<i>Procyon lotor</i>	Raccoon	0.0002	0.000	0.000	0.0002

CVe= Cultural value of the species; PVe= Practical value of the species; EVe= Economic value of the species; Ve= Total value of species e.

## REFERENCES

Aguado LE (1993) **La reproducción campesina y las estrategias de sobrevivencia en el mundo rural.** *Convergencia* 1:157-187.

Aldana MNJ, Díaz PM, Feijoo MA, Quintero H (2016) **Percepciones y reconocimiento local de fauna silvestre, municipio de Alcalá, departamento del Valle del Cauca, Colombia.** *Revista Luna Azul* No. 43: 56-81.

Alves RRN, Souto WMS (2015) **Ethnozology: A Brief Introduction.** *Ethnobiology and Conservation* 4. doi: [10.15451/ec2015-1-4.1-1-13](https://doi.org/10.15451/ec2015-1-4.1-1-13)

Amador ASA, De la Riva HG (2016) **Uso tradicional de la fauna silvestre en las serranías del oc-**

**cidente del estado de Aguascalientes, México.** *Etnobiología* 14:20-36.

Aranda M (2013) **Guía de campo: huellas de los mamíferos mexicanos.** Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México.

Arévalo PML (2016) **La reubicación como proceso de desterritorialización.** *Política y Cultura* 45:153-180.

Asprilla PJ, Díaz PJM (2020) **Uso de alimentos silvestres de origen animal en comunidades rurales asociadas con bosque húmedo tropical al noroeste de Colombia.** *Interciencia* 45:76-83.

Ayala EMI, García LF, Román MOE (2019) **La apropiación de los recursos naturales silvestres y cultivados.** *Estrategia alimentaria de los pue-*

**blos originarios.** En: Román MOE (coord.) prácticas agropecuarias como estrategias de seguridad alimentaria 1 ed. Universidad Autónoma del Estado de Morelos, México, págs.73-104.

Bardales AC, Torres OL, Tirado HE, Antúnez CM (2017) **Presión de caza en mamíferos de la Reserva Nacional Matsés, al noreste de la Amazonía Peruana.** *Folia Amazonica* 26:75-84.

Barrasa GS (2012) **Conocimiento y usos tradicionales de la fauna en dos comunidades campesinas de la Reserva de la Biósfera de la Encrucijada, Chiapas.** *Etnobiología* 10:16-28.

Boege E (2008) **El patrimonio biocultural de los pueblos indígenas de México. Hacia la conservación in situ de la biodiversidad y agrobiodiversidad en los territorios indígenas.** 1 edición Instituto Nacional de Antropología e Historia, Comisión Nacional para el Desarrollo de los Pueblos Indígenas, México.

Bustos ZMG, Castro FR (2020) **Reptiles.** En: Cruz AA, Nájera CKC, López HD, Melgarejo ED, González FL, Maldonado KC, Flores CME, Fuentes VL (coords.) La biodiversidad en Morelos. Estudio de Estado 2. 1 ed. Comisión Nacional para el Uso de la Biodiversidad, México, págs. 295-301.

Cano-Contreras EJ, Medinaceli A, Sanabria O, Argueta A (2016) **Código de Ética para la investigación, la investigación-acción y la colaboración etnocientífica en América Latina.** 1 ed. Asociación Etnobiológica Mexicana, Sociedad Latinoamericana de Etnobiología, Ciudad de México.

Castillo AA, Peña MJL (2015) **Métodos de investigación social: fundamentos, técnicas y aportaciones para el entendimiento de las relaciones sociedad-vida silvestre.** En: Gallina S (ed.) Manual de técnicas del estudio de la fauna. 1 edición Instituto de Ecología, A. C. Veracruz, México págs. 189-210.

Castro FR, Bustos ZMG (2020) **Anfibios.** En: Cruz AA, Nájera CKC, López HD, Melgarejo ED, González FL, Maldonado KC, Flores CME, Fuentes VL (coords.) La biodiversidad en Morelos. Estudio de Estado 2. 1 ed. Comisión Nacional para el Uso de la Biodiversidad, México, págs. 289-294.

CONANP (2005) **Programa de Conservación y Manejo de la Reserva de la Biósfera Sierra de Huautla.** 1 edición Comisión Nacional de Áreas Naturales Protegidas, México.

Contreras M, Yanguéz A (2017) **Caracterización de los usos de vertebrados silvestres en la comunidad de Quebrada Ancha, Colón (Panamá).** *Revista Científica CENTROS* 6:18-30.

Costa-Neto EM, Vargas-Clavijo M, Santos-Fita D (2009) **Introducción.** En: Costa-Neto EM, Santos-Fita D, Vargas-Clavijo M (coords.) Manual de etnozooloía. Una guía teórico-práctica para investigar la interconexión del ser humano con los animales. 1 edición Tundra Ediciones, Valencia. págs. 15-20.

Costa JF, Ríos-Choronto R, Peña-Candia LZ, Simoes E (2018) **Aves y mamíferos silvestres usados por pobladores del Bajo Urubamba, Cusco, Perú.** *Revista Peruana de Biología* 25:463-470.

Dardón ER, Retana GO (2017) **Uso medicinal de la fauna silvestre por comunidades mayas, en la región de los Chenes, Campeche, México.** *Revista Etnobiología* 15:68-83.

Estrada PDS, Rosas ROC, Parra IF, Guerrero RJD, Tarango ALA (2018) **Valor de uso, importancia cultural y percepción sobre mamíferos silvestres medianos y grandes en la mixteca poblana.** *Acta Zoológica Mexicana* 34:1-15.

García FA (2008) **La etnozooloía como una alternativa para el desarrollo comunitario sustentable en la Reserva Estatal Sierra de Montenegro, Morelos, México.** Tesis de maestría. Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos.

García FA, Valle MR, Pino MJM, Monroy MR (2018) **Knowledge and uses of wildlife in the community of Tetelpa, municipality of Zacatepec, Morelos, México.** *Gaia Scientia* 12:19-35.

García FA, Colín BH, Monroy MR (2019a) **Uso social de mamíferos silvestres en Tepoztlán y Tlaltizapán, Morelos.** *Inventio* 15:31-36.

García FA, Farfán EE, Monroy MR, Monroy OR, Colín BH, Pino MJM (2019b) **Fauna registrada en huertos frutícolas tradicionales de Yautepec, Morelos, México.** *Tropical and Subtropical Agroecosystems* 22:359-378.

García FA, Monroy MR, Pino MJM (2020) **Estudio etnozoológico de los vertebrados silvestres de la comunidad Bonifacio García, Morelos, México.** *Revista Peruana de Biología* 27:361-374.

García FA, Bello RM, Rebollo SDB, Cosme MJL, Urbina TF, Román MOE, Colín BH (2022) **Conocimiento ecológico tradicional de las aves de San Andrés de la Cal, Tepoztlán, Morelos, México.** *Tropical and Subtropical Agroecosystems* 25:1-18.

Gaviño TG (2015) **Aves de Morelos. Estudio general, guía de campo y recopilación bibliográfica.** 1 edición Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México.

Gotoo CN (2016) **Importancia del uso de fauna**

silvestre en dos localidades costeras de Chiapas, México. Tesis de licenciatura. Universidad de Ciencias y Artes de Chiapas, Tonalá, Chiapas.

Guerra RMM, Naranjo PEJ, Limón AF, Mariaca MR (2004) **Factores que intervienen en la regulación local de la cacería de subsistencia en dos comunidades de la selva lacandona, Chiapas, México.** *Etnobiología* 4:1-18.

Guerrero JA, Ávila LG, Uriostegui JM, Rizo AA (2020) **Mamíferos.** En: Cruz AA, Nájera CKC, López HD, Melgarejo ED, González FL, Maldonado KC, Flores CME, Fuentes VL (coords) *La biodiversidad en Morelos. Estudio de Estado 2.* 1 ed. Comisión Nacional para el Uso de la Biodiversidad, México, págs. 339-347.

Guzmán GE, León LA (2014) **Peculiaridades campesinas del Morelos rural.** *Economía, Sociedad y Territorio* 14:175-200.

Hernández-López A, López-Alamilla E, Rodríguez-Ramírez A, Aquino-Bravata V (2013) **Diagnóstico del uso de la fauna silvestre en el Área de Protección de Flora y Fauna “cañón del Usumacinta”, Tenosique tabasco.** *Ra ximhai* 9:1-14.

INEGI (2020) **Censo de población.** México: Instituto Nacional de Estadística y Geografía. <https://www.inegi.org.mx/programas/ccpv/2020/default.html#Microdatos> Consultado el 20 de diciembre de 2021.

Infante GS, Zárate L (2010) **Métodos estadísticos: un enfoque multidisciplinario.** 1 edición. Trillas, México.

Lira-Torres I, Briones SM, Gómez AFR, Ojeda RD, Peláez AA (2014) **Uso y aprovechamiento de la fauna silvestre en la Selva Zoque, México.** *Acta Zoológica Mexicana* (n.s.) 30:74-90.

López GM, Bustamante GA, Vargas LS, Morales JJ, Pérez RN, Guadarrama LR, Díaz HH (2018) **Conocimiento y aprovechamiento local del venado cola blanca (*Odocoileus virginianus mexicanus*) en Iliatenco, Guerrero.** *Agroproductividad* 11:127-132.

Martínez-Ceballos M (2014) **Caracterización del conocimiento, uso y manejo tradicional de la fauna silvestre en áreas protegidas y no protegidas del corregimiento del Encano municipio de Pasto, estudio etnozoológico.** Tesis de licenciatura. Universidad de Nariño, San Juan de Pasto, Colombia.

Méndez CF, Montiel S (2007) **Diagnóstico preliminar de la fauna y flora silvestre utilizada por la población maya de dos comunidades coste-**

**ras de Campeche, México.** *Universidad y Ciencia* 23:127-139.

Miranda F, Hernández-X E (2014) **Los tipos de vegetación de México y su clasificación.** Edición conmemorativa 1963- 2013. 1 edición Fondo de Cultura Económica, México.

Monroy-Vilchis O, Cabrera L, Suárez P, Zarco-González MM, Rodríguez SC, Urios V (2008) **Uso tradicional de vertebrados silvestres en la Sierra de Nanchititla, México.** *Interciencia* 33:308-313.

Monroy MR, García FA, Monroy OR (2011) **Importancia de la fauna silvestre en regiones bajo presión inmobiliaria en la cuenca del Río Grande Amacuzac, Morelos, México.** En: Monroy R, García A, Pino MJM, Monroy R. (eds) *Etnozoología, un enfoque binacional México- Colombia.* 1 edición Universidad Autónoma del Estado de Morelos, Morelos, México, págs. 19-40.

Monroy R, García FA (2013) **La fauna silvestre con valor de uso en los huertos frutícolas tradicionales de la comunidad indígena de Xoxocotla, Morelos, México.** *Etnobiología* 11:44-52.

Montalbán HCA, Aréchaga OS (2014) **Anfibios de Morelos Región Centro-Sur.** 1 edición Facultad de Ciencias Biológicas, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos.

Montiel OS (2010) **Aprovechamiento de fauna silvestre en la península de Yucatán: usos y costumbres.** 1 edición Fornix Campeche. Gobierno del estado de Campeche, Conacyt, México.

Peterson RT, Chaliff EL (1989) **Guía de campo de las Aves de México.** 1 edición Editorial Diana, México.

Pino MJM, Reyes PH, García FA (2018a) **Edible insects in some Latin American Countries: Advantages and Prospects.** En: Ganguly A, Naskar K (eds) *Entomology, Current status and future strategies.* 1ed. Daya Publishing House, Astral International Pvt.Ltd., New Delhi, pp.171-190.

Pino MJM, García FA, Monroy MR, Barreto SSD (2018b) **The “Jumiles” (Hemiptera: Pentatomidae and Coreidae) in the State of Morelos, México: Taxonomy, Distribution and Commercialization.** En: Ganguly A, Naskar K (eds) *Entomology, Current status and future strategies.* 1ed. Daya Publishing House, Astral International Pvt.Ltd., New Delhi, pp.191-213.

Plata EE (2017) **El uso del perro (*Canis lupus familiaris*) en la cacería maya tradicional en grupo (batida): relevancia práctica y sociocul-**

tural. Tesis de maestría Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional Mérida, Yucatán.

Pries L (1999) **La migración internacional en tiempos de globalización.** Nueva Sociedad 164:56-68.

Quijano HE, Calmé S (2002) **Patrones de cacería y conservación de la fauna silvestre en una comunidad maya de Quintana-Roo, México.** *Etnobiología* 2:1-18.

R Core Team (2021) **R: A language and environment for statistical computing.** <https://www.r-project.org> Consultada el 8 de enero del 2022.

Ramírez-Mella M, Candelaria-Martínez B, Dorantes-Jiménez J, Tarango-Arámbula LA, Flota-Bañuelos C (2016) **Uso y aprovechamiento de fauna silvestre en zonas rurales de Campeche, México.** *Agroproductividad* 9:3-9.

Retana GOG, Padilla PSE (2018) **Cacería y aprovechamiento del venado cola blanca por indígenas mayas.** *Tropical and Subtropical Agroecosystems* 21:283-294.

Reyes-García V, Huanca T, Valdez V, Leonard W, Wilkie D (2006) **Cultural, practical, and economic value of wild plants: a quantitative study in the Bolivian Amazon.** *Economic Botany* 60:62-74.

Reyna RMA (2013) **Conocimiento etnoherpetológico en las comunidades de Barranca Honda y Ticumán, en la Reserva Estatal Sierra de Montenegro, Morelos, México.** Tesis de licenciatura, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos.

Ríos VA (2020) **El potencial del campesino-cazador para el monitoreo comunitario de fauna silvestre en una comunidad maya de Campeche.** Tesis de maestría, CINVESTAV Unidad Mérida, Mérida Yucatán.

Román-Montes de Oca E (2017) **La familia agrícola y sus estrategias de reproducción.** En: Guzmán-Gómez E, Madera-Pacheco JA (coords) *Estrategias e identidades productivas campesinas.* 1 edición, Asociación Mexicana de Estudios Rurales, A. C., Universidad Autónoma de Nayarit, Universidad Autónoma de Chapingo, Universidad Autónoma Metropolitana, México, págs. 91-105.

Sarukhán J, Koleff P, Carabias J, Soberon J, Dirzo R, Llorente-Bousquets J, Halffter G, González R, March I, Mohar A, Anta S, de la Maza J, Pisanty I, Urquiza HT, Ruiz GSP, García MG (2017) **Capital natural de México. Synthesis: evaluación del conoci-**

**miento y tendencias de cambio, perspectivas de sustentabilidad, capacidades humanas e institucionales.** 1 edición, Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México.

SEMARNAT (2010) **Norma Oficial Mexicana NOM-059-SEMARNAT-2010.** [https://dof.gob.mx/nota\\_detalle\\_popup.php?codigo=5173091](https://dof.gob.mx/nota_detalle_popup.php?codigo=5173091) Consultada el 20 de diciembre del 2021.

Solo de Zaldivar VB (1993) **¿De campesino a agricultor? La pequeña producción familiar en el marco del desarrollo capitalista.** *Noticiario de Historia Agraria* 5:127-159.

Taboada SM, Granjeno CAE, Guadarrama RO (2009) **Normales climatológicas (temperatura y precipitación) del estado de Morelos.** 1 edición, Facultad de Ciencias Biológicas, Universidad Autónoma del Estado de Morelos, México.

Tamburini DM (2016) **La fauna silvestre en las estrategias de reproducción social de los campesinos del Chaco Seco de la Provincia de Córdoba (Argentina).** Tesis de doctorado, Universidad Nacional de Córdoba, Córdoba, Argentina. AREA.

Taylor SJ, Bogdan R (1987) **Introducción a los métodos cualitativos de investigación.** Editorial Paidós, España.

Toledo VM, Barrera BN (2008) **La memoria biocultural. La importancia ecológica de las sabidurías tradicionales.** 1 edición Icaria editorial, España.

UICN (2021) **The UICN Red List of Threatened Species. Version 2020-3.** <https://www.iucnredlist.org/es> Consultada el 10 de enero del 2022.

UNESCO (2009) **Invertir en la diversidad cultural y el diálogo intercultural.** *Informe Mundial de la UNESCO* 7:205-235.

Urbina TF, Morales GG (1994) **Aves de Morelos de importancia económica y rapaces diurnas.** 1 edición Centro de Investigaciones Biológicas, Universidad Autónoma del Estado de Morelos, Secretaría de Educación Pública, Cuernavaca, Morelos.

Urbina TF (2020) **Aves.** En: Cruz AA, Nájera CKC, López HD, Melgarejo ED, González FL, Maldonado KC, Flores CME, Fuentes VL (coords.) *La biodiversidad en Morelos. Estudio de Estado 2.* 1 ed. Comisión Nacional para el Uso de la Biodiversidad, México, págs. 311-324.

Valle MR, García FA, Colín BH (2021) **Fauna silvestre con valor de uso en la Reserva de la Biosfera Sierra de Huautla, Morelos, México.** *Revista Peruana de Biología* 4:63-78. doi:

[10.15381/rpb.v28i4.19921](https://doi.org/10.15381/rpb.v28i4.19921).

Vargas CM (2009) **Patrimonio zoocultural: el mundo animal en las expresiones tradicionales de los pueblos**. En: Costa EM, Santos FD, Vargas CM. (eds) Manual de etnozología. Una guía teórico-práctica para investigar la interconexión del ser humano con los animales. 1 edición Ediciones Tundra, Valencia, págs. 118-144.

Velarde ES, Cruz LA (2015) **La fauna silvestre y su relación con el bienestar de tres comunidades de la Reserva de la Biósfera Sierra de Huautla**. *Etnobiología* 13:40-52.

Zarazúa CM, Chávez GM, Romero BY, Rangel LS, Moreno CAI, Alvarado RLF, Smith SE, Blancas J, del Val E, Arizmendi MC, Casas A (2020) **Use and management of wild fauna by people of the Tehuacán-Cuicatlán Valley and surrounding areas, México**. *Journal of Ethnobiology and*

*Ethnomedicine* 16:1-23.

Zavala SZ (2018) **Valoración cultural, usos y manejo de la fauna silvestre en comunidades rurales del estado de Guerrero**. Tesis de maestría. Universidad Autónoma de Guerrero, Chilpancingo de los Bravo, Guerrero.

Zavala SZ, Segura PHR, Ávila NDM, Herrera CND, Barrera CE, Sarabia RG (2018) **Valoración cultural y uso de la fauna silvestre en San Vicente de Benítez, Guerrero, México**. *Revista Etnobiología* 16:78-92.

**Received:** 19 November 2022

**Accepted:** 16 March 2023

**Published:** 28 March 2023

**Editor:** Ulysses Albuquerque