

Food Biodiversity as an Opportunity to Address the Challenge of Improving Human Diets and Food Security

Michelle Cristine Medeiros Jacob^{1*} , Alice Medeiros Souza² ,
Aline Martins de Carvalho³ , Carlos Frederico Alves de Vasconcelos Neto⁴ ,
Daniel Tregidgo⁵ , Danny Hunter⁶ , Fillipe de Oliveira Pereira⁷ ,
Guilhermo Ros Brull⁸ , Harriet V. Kunhlein⁹ , Lara Juliane Guedes da Silva¹⁰ ,
Larissa Mont'Alverne Jucá Seabra¹¹ , Mariana de Paula Drewinski¹² ,
Nelson Menolli Jr¹³ , Patricia Carignano Torres¹⁴ , Pedro Mayor¹⁵ ,
Priscila F. M. Lopes¹⁶ , Rafael Ricardo Vasconcelos da Silva¹⁷ ,
Sávio Marcelino Gomes¹⁸  and Juliana Kelly da Silva-Maia¹⁹ 

ABSTRACT

Scientists have warned for several years that food systems have become major drivers of environmental degradation, malnutrition, and food insecurity. In this paper, we present arguments from specialists that suggest that in the transition to more sustainable food systems, biodiversity and food security can be mutually supportive rather than conflicting goals. We have divided the opinions of these scientists into two “Big Topics”. First, they examine the synergies and challenges of the intersection of biodiversity and food security. In the second section, they explain how various forms of food biodiversity, such as mushrooms, terrestrial wild animals, aquatic animals, algae, and wild plants, can contribute to food security. Finally, we present three main pathways that, according to these experts, could guide the transition toward biodiversity and food security in food systems.

Keywords: Food Biodiversity, Food Security, Nutrition, Wild Foods, Sustainable Development.

1 LabNutrir, Nutrition Department, Federal University of Rio Grande do Norte, Av. Senador Salgado Filho, s/n, Lagoa Nova, 59078-970, Natal, RN, Brazil.

2 LabNutrir, Federal University of Rio Grande do Norte.

3 Department of Nutrition, School of Public Health, University of Sao Paulo, Brazil.

4 Grupo de Pesquisa em Ecologia de Vertebrados Terrestres (ECOVERT), Instituto de Desenvolvimento Sustentável Mamirauá (IDSMA), Rua Estrada do Bexiga, 2585, Bairro Fonte Boa, Tefé, AM.

5 Grupo de Pesquisa em Ecologia de Vertebrados Terrestres, Instituto de Desenvolvimento Sustentável Mamirauá, Rua Estrada do Bexiga, 2585, Bairro Fonte Boa, Tefé, Amazonas, Brazil.

6 Alliance of Biodiversity International and CIAT, Via di San Domenico 1, 00153, Rome, Italy.

7 FUNGI Research Group, Academic Unit of Health, Education and Health Center, Federal University of Campina Grande, Sítio Olho D'água da Bica, s/n, 58175-000, Cuité, Brazil.

8 Zerca y Lejos, Calle de Sambara, 128, 28027, Madrid, Spain.

9 Centre for Indigenous Peoples' Nutrition and Environment (CINE), and School of Human Nutrition, McGill University, Montreal, Canada.

10 Nutrition Department, Federal University of Rio Grande do Norte, Av. Senador Salgado Filho, s/n, Lagoa Nova, 59078-970, Natal, RN, Brazil.

11 Nutrition Department, Federal University of Rio Grande do Norte, Av. Senador Salgado Filho, s/n, Lagoa Nova, 59078-970, Natal, RN, Brazil.

12 IFungiLab, Instituto Federal de Educação, Ciência e Tecnologia de São Paulo (IFSP), Câmpus Sao Paulo (SPO), Departamento de Ciências da Natureza e Matemática (DCM), Subárea de Biologia (SAB), Rua Pedro Vicente 625, São Paulo, SP, 01109-010, Brazil. Instituto de Pesquisas Ambientais (IPA), Núcleo de Pós-graduação Stricto Sensu, Pós-graduação em Biodiversidade Vegetal e Meio Ambiente, Av. Miguel Stéfano 3687, 04301-902, São Paulo, SP, Brazil.

13 IFungiLab, Instituto Federal de Educação, Ciência e Tecnologia de São Paulo (IFSP), Câmpus Sao Paulo (SPO), Departamento de Ciências da Natureza e Matemática (DCM), Subárea de Biologia (SAB), Rua Pedro Vicente 625, São Paulo, SP, 01109-010, Brazil. Instituto de Pesquisas Ambientais (IPA), Núcleo de Pós-graduação Stricto Sensu, Pós-graduação em Biodiversidade Vegetal e Meio Ambiente, Av. Miguel Stéfano 3687, 04301-902, São Paulo, SP, Brazil.

14 Graduate Program in Complex Systems Modeling, School of Arts, Sciences and Humanities, University of São Paulo, Rua Arlindo Bettio 1000, São Paulo, SP, Brazil.

15 Departament de Sanitat i Anatomia Animals, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain.

16 Fishing ecology, management and economics group, Department of Ecology, Universidade Federal do Rio Grande do Norte, Natal, RN, Brazil.

17 Laboratory of Biocultural Ecology, Conservation and Evolution (LECEB), Campus of Engineering and Agricultural Sciences, Federal University of Alagoas, BR-104, km 85, s/n, 57100-000, Rio Largo, AL, Brazil.

18 Department of Nutrition, Federal University of Paraíba, Street Tabelaio Stanislaw Eloy, s/n, Castelo Branco, 58050-585, João Pessoa, PB, Brazil.

19 Nutrition Postgraduate Program, Nutrition Department, Federal University of Rio Grande do Norte, Av. Senador Salgado Filho, s/n, Lagoa Nova, 59078-970, Natal, RN, Brazil.

* Corresponding author ✉. E-mail address: MCMJ (michelle.jacob@ufrn.br), AMS (almedeiros.s@gmail.com), AMC (alinenutri@usp.br), CFAVN (fredvasconcelosnt@gmail.com), DT (dantregidgo@gmail.com), DH (d.hunter@cgiar.org), FOP (fillipe.oliveira@professor.ufcg.edu.br), GRB (guillermo.ros.brull@gmail.com), HVK (harriet.kuhnlein@mcgill.ca), LJGS (lara.guedes.123@ufrn.edu.br), LMJS (larissa.seabra@ufrn.br), MPD (maridrewinski@gmail.com), NMJ (menolljr@yahoo.com.br), PCT (patriciactorres@gmail.com), PM (pedrogines.mayor@uab.cat), PFML (priscila.lopes@ufrn.br), RRVS (rafaelrvsilva@gmail.com), SMG (savio.gomes@academico.ufpb.br) and JKSM (juliana.maia@ufrn.br).

INTRODUCTION

The idea that biodiversity conservation and food security are incompatible is based on a narrow view of food security as solely related to food production (Fischer *et al.* 2017). Some theories (e.g., land-sparing, Malthusian and neo-Malthusian, production possibility frontier, ecological modernization theory, etc.) assume that increasing food production will automatically lead to food security (Béné *et al.* 2019; Meli *et al.* 2019). However, the case of Brazil illustrates the weakness of this logic. Data from the National Food Supply Company of Brazil (CONAB) show that in the 2022/2023 harvest, grain production reached a new record of 312.4 million tons, surpassing the previous record set in 2021/2022 by 41.5 million tons (CONAB 2022). Despite being a major producer and exporter of commodities such as meat, soy, and corn (Aragão and Contini 2021), food insecurity in Brazilian households has also reached new highs. In 2020, 9%, or 19 million people, experienced severe food insecurity (a proxy for hunger); in 2022, this number rose to 33 million, or 15.5% of the population (PENSSAN 2021,

2022). Additionally, even though Brazil is a major agricultural supplier, the prevalence of anemia among children under two years old is 10%, with the highest rate being found in the North region at 17% (ENANI 2021). Moreover, anemia is more prevalent among nontribal river dwellers (ribeirinhos) in the northern region at 59% (Torres *et al.* 2022) and even higher among indigenous peoples such as the Yanomamis at 68% (Basta and Orellana 2020). Therefore, food security is not the automatic result of food production.

According to the Food and Agriculture Organization (FAO 1996), food security is defined as a state in which people have access to sufficient, safe, and nutritious food to meet their dietary needs and food preferences while being able to maintain an active and healthy lifestyle and preserving cultural diversity and economic, social, and environmental sustainability. This concept is supported by four key pillars: availability (a reliable food supply), accessibility (the resources to obtain proper food), utilization (food quality and health conditions), and stability (permanent access to food). Therefore, based on this definition, food production (a proxy for availability) does not auto-

matically lead to food distribution to people (access), as these are distinct food security levels. Furthermore, by considering that food security is more than production, biodiversity can provide a range of opportunities to enhance access to nutritious and safe food (i.e., utilization) (El Bizri *et al.* 2020; Lachat *et al.* 2018) at all times (i.e., stability) (Tregidgo *et al.* 2020). It is difficult to quantify exactly how much of the world's biodiversity is neglected as food. However, it is widely recognized that a significant proportion of the world's biodiversity is underutilized or undervalued as food. For example, a study (Khoury *et al.* 2014) used national per capita food supply data published by the FAO to analyze trends in the richness, abundance, and composition of agricultural commodities measured in the food supply of 152 countries, comprising 98% of the world's population, between 1961 and 2009. Although there may be approximately 30,000 edible plant species on the planet (FAO 2010), according to the study, just 50 of the measured agricultural commodities (such as wheat, rice, and corn) contribute to the top 90% of calories, protein, fat, and weight in the world. The increasing reliance on a smaller number of crops to feed a growing population poses a real threat to food security (see Figure 1).

Biodiversity conservation and food security can be mutually reinforcing rather than competing goals. The Convention on Biological Diversity defines biodiversity as *the variability among living organisms from all sources including but not limited to, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystem* (UNEP 1992). There are many synergies between food security and biodiversity, as highlighted in a literature review by Blicharska *et al.* (2019) that examined the contributions of biodiversity (SDG14 and SDG15) to sustainable development goals (SDG2). These contributions can be broken down into the critical pillars of food security. At the availability level, biodiversity improves soil fertility, structure, quality, and health; provides crop pollination; and controls pests. At the utilization level, it improves dietary quality. At the stability level, it increases agricultural output and future yields, provides potential new crops, and maintains productivity in marine ecosystems. In addition, genetic diversity is important for increasing the resilience of agri-food systems to threats such as pathogens and climate change (IPCC 2019; UNEP 2020). These synergies can be achieved through a new paradigm of food production that includes sustainable agriculture, aquaculture and livestock systems, agroecological practices, the safeguarding of native species, varieties, breeds, and habitats, and ecological restoration (IPBES 2019).

Given the various opportunities and challenges that exist at the intersection of food security and biodiversity, we asked specialists in both fields how food biodiversity could help us address the challenge of food security. We surveyed specialists based primarily in the Americas, but some also conducted research in European and African countries. Two main themes emerged from their responses. The first theme focused on the synergies and challenges of the relationship between biodiversity and food security, considering the diverse research areas in which these scientists work. The second theme provided an overview of food biodiversity resources that have the potential to support the transition to sustainable diets while ensuring food security. Figure 2 summarizes the topics covered within “Big Topics” one and two.

BIG TOPIC 1: EXPLORING THE INTERSECTION OF BIODIVERSITY AND FOOD SECURITY: SYNERGIES AND CHALLENGES

This first section summarizes some of the multiple factors that should be considered when exploring the intersection of biodiversity and food security, such as seasonality and stability, indigenous and traditional knowledge, governance approaches, and the global scale. It also identifies the main challenges (e.g., climate change, the emergence of hunger, and the emergence of zoonotic diseases) that we could overcome by considering both people and nature when analyzing and proposing changes to our food systems. All opinions listed below are in alphabetical order.

Aline Martins de Carvalho e Larissa Mont’Alverne Jucá Seabra: Climate change mitigation through a local and biodiverse diet

According to the Intergovernmental Panel on Climate Change (IPCC), the food system is responsible for 11 to 19 Gt CO₂-equivalent emissions per year globally, which constitutes 21% to 37% of total anthropogenic emissions (IPCC 2019). The situation is worse in Brazil, where approximately 70% of greenhouse gas emissions (GHGE) are related to the food system (SEEG. System of Estimates of Emissions and Removals of Greenhouse Gases 2019). The high level of greenhouse gas emissions (GHGE) in Brazil can be partly attributed to the country's status as a major agroproducer and one of the world's leading producers and exporters of meat, soy, corn, and other commodities (Belik 2020). One way to mitigate climate change is through the adoption of a local and biodiverse diet.




Figure 1. Key pillars of food security, which extends beyond just food production. These components include the availability of a reliable food supply, the accessibility of resources to obtain adequate food, the utilization of food in terms of quality and health, and the stability of permanent access to food.




Figure 2. Summary of the synergies and challenges at the intersection of biodiversity and food security (Big Topic 1) and the role of strategic food biodiversity resources in enhancing food security (Big Topic 2).

Food production is the largest contributor to emissions in the food supply chain, but reducing food miles (i.e., the transportation of food) can decrease total emissions by approximately 5%, benefit small local farmers socially and financially, and reduce food waste (Crippa *et al.* 2021). A more biodiverse diet is a significant dietary change that can have environmental benefits, as the current Brazilian diet has a high en-

vironmental impact and is of poor quality. In a study optimizing nutritional recommendations for the Brazilian population, all the models showed that reducing red meat and increasing fruits and vegetables would decrease the impact of GHGE from diet by 10-27% (Verly Jr *et al.* 2022) and increase biodiversity. Currently, beef is one of the most consumed foods in the country and is responsible for 68% of GHGs (Travas-

sos *et al.* 2020). Other meats, such as goats, which are well suited to the semiarid climate in Brazil, as well as legumes, cereals, fruits, and vegetables (Gazzillo *et al.* 2019), have a much lower environmental impact. Therefore, a local and biodiverse diet can be a solution for mitigating climate change, especially in Brazil.

Danny Hunter: Incorporating Food Biodiversity into the Global Food System: Challenges and Opportunities for Advancing Human Nutrition

Food biodiversity is under threat globally because current agri-food production and consumption rely on a small number of food species. Homogenous diets, limited food access, and underdeveloped markets for trading underutilized, nutrient-rich food biodiversity contribute to the persistence of malnutrition, especially hidden hunger, poverty, and environmental degradation, including biodiversity loss and climate change. This reliance on a narrow range of food crops introduces a significant vulnerability in our food production and food systems, as large areas of monocrops continue to depend on corporate-controlled seed systems that require significant external chemical inputs. Our global agri-food system remains unsustainable and inequitable, with over 80% of global agricultural subsidies going toward industrial agriculture for cereal production and red meat, while less than 1% goes to much-needed fruit and vegetable production, which is nearly universally underconsumed (Hunter *et al.* 2020). Additionally, very little in the way of subsidies goes toward research and development of underutilized food biodiversity species. The governance and political economy of the agri-food system play a key role in the current crisis humanity is facing. To address this issue, a significant overhaul of the agri-food system is necessary, including a reorientation of subsidies toward more biodiversity-friendly food production systems, including targeted support for untapped, nutrient-rich food biodiversity. The strategic use of locally adapted, culturally appropriate nutrient-rich food biodiversity must be central to this challenge to conserve this biodiversity and offer environmental, livelihood, and health benefits to communities worldwide. Based on our work on this topic at Biodiversity International (Kennedy *et al.* 2017), we highlight three strategic pathways for how country-driven collaborations and networks employing multi-sectoral and interdisciplinary approaches can support agri-food system transformation toward biodiversity: 1) providing evidence for the nutritional value of food biodiversity; 2) influencing policy and markets to incentivize food biodiversity; and 3) creating awareness, advocacy, and action around consumer behavi-

oral change.

Daniel Tregidgo: Exploring the Role of Food Biodiversity in Ensuring Stability in Food Security

Food stability, or being food always secure, is often overlooked as one of the four key pillars of food security (Cruz-Garcia *et al.* 2016). Food instability can be caused by various shocks, such as wars, crop failures, and droughts, but is most commonly a result of seasonality, a characteristic of agricultural food systems worldwide (Vaitla *et al.* 2009). Food biodiversity, including wildmeat, fish, plants, and mushrooms, is also highly seasonal and is influenced by species phenology - e.g., Nyahongo *et al.* (2009) - and fluctuating environmental factors that impact harvestability. In areas where people rely on wild food, such as fish, lean seasons can result in severe food insecurity (Tregidgo *et al.* 2020). Food biodiversity can, however, act as an important safety net during periods of food instability (lean seasons and environmental shocks) (Karjalainen *et al.* 2010). Increased wild fish (Coomes *et al.* 2010) and wildmeat (Golden *et al.* 2019) consumption is common during agricultural lean seasons, while increased wildmeat (Brashares *et al.* 2004; Endo *et al.* 2016) and even fruit consumption during periods of fish shortages may offset reduced fish-derived nutrient intake (Tregidgo *et al.* In review). Wild foods are argued to be more resilient than many farmed foods, making them particularly important natural insurance against shocks in the context of ongoing climate change. In addition, the seasonality of food biodiversity systems can increase dietary biodiversity through the availability of different species throughout the year, potentially leading to improved micronutrient adequacy (Lachat *et al.* 2018).

Frederico Vasconcelos and Pedro Mayor: Assessing the Impact of COVID-19 on Food Security in South America and the Role of Biodiversity as a Guarantee

Latin America is known for its rich biodiversity, accounting for one third of all animal and plant species on the planet (Raven *et al.* 2020), as well as for its high agricultural production, which unfortunately also poses a major threat to its biodiversity. The current productive model, combined with other sociopolitical factors, contributes to high levels of social inequality and, as a result, serious problems with food and nutrition security (Rosen and Shapouri 2009). The global health crisis caused by COVID-19

has further exposed these pre-existing vulnerabilities. Our estimates for 2019 in South America showed that 22 million people had moderate and severe food insecurity (Clapp and Moseley 2020). For 2020, during the COVID pandemic, we saw an increase of 12.4% (+84.4 M) in the number of people facing food insecurity in urban areas (Clapp and Moseley 2020). Meanwhile, the contribution of the agricultural sector grew, and food exports increased by 5% (Vasconcelos-Neto *et al.* in press); however, the population experienced food price inflation of up to 22.5% (October 2020) (The World Bank 2022). In general, efforts to alleviate hunger should take a multisectoral approach, including policies that support agrarian reform, ensure access to education and basic health, reduce poverty and job insecurity, and regulate food prices (FAO 2020a). At the same time, subsistence hunting and fishing have helped to ensure food security in rural societies in South America. Therefore, to guarantee food security and the well-being of these societies that rely on subsistence economies, we advocate for policies that promote the conservation of biodiversity and traditional knowledge (Bodmer *et al.* 2020).

Harriet V. Kuhnlein: Indigenous Peoples' food systems contain extensive knowledge related to food biodiversity that needs documentation, protection, and promotion

There is growing evidence of the numerous benefits of indigenous peoples' food systems, including the nutrient quality and health and wellness benefits derived from their sociocultural, mental, and spiritual dimensions (Kuhnlein *et al.* 2013, 2009). These unique foods, derived from plant and animal species, are known to indigenous peoples through their biocentric connection to sustainably managed resources that historically provided adequate diets with traditional knowledge in the many ecosystems inhabited by indigenous peoples around the world. However, these food systems face numerous risks to their biodiversity from environmental threats to indigenous territories and health disparities for indigenous peoples due to poverty and marginalization, leading to a loss of food security and higher rates of obesity and undernutrition. It is important to recognize the scientific and cultural value of the biodiversity of indigenous food systems and to protect this knowledge for both its holders and the global food heritage. Indigenous peoples' food systems should be a priority for research and documentation, promotion, and protection to shift global food systems toward more sustainable diets and food security. Universities, governments, and United Nations agencies can contribute to this

effort by conducting research with indigenous peoples using free, prior, and informed consent and by empowering women and youth with intercultural food and diet education at all levels, benefiting both indigenous communities and the wider global population (FAO 2021a, 2021b).

BIG TOPIC 2: EXPLORING THE POTENTIAL OF STRATEGIC FOOD BIODIVERSITY RESOURCES FOR ENHANCING FOOD SECURITY

The FAO (2017) defines “food biodiversity” as the *diversity of plants, animals, fungi, algae, and other organisms used for food, including the genetic resources within species, between species, and provided by ecosystems*. This final section summarizes how various forms of food biodiversity, such as mushrooms, terrestrial wild animals, aquatic animals, algae, and wild plants, can contribute to food security. We have received two comments about wild meat and edible mushrooms each. In the case of these food resources, we have placed the opinions alongside each category.

EDIBLE MUSHROOMS

Fillipe de Oliveira Pereira: Edible Mushrooms Contribute to Sustainable Human Diets

Promoting food security necessarily involves facilitating access to food, diversifying the diet, and sustainably producing food. Thus, wild and cultivated species of edible mushrooms can be strategic in achieving these goals. The consumption of mushrooms has long been part of the dietary history of humanity. Traditionally, eastern regions of the planet have a higher consumption of mushrooms, but this distinction has lessened with broadened access to edible species in the West. At present, people in urban areas around the globe may find species dried, fresh, canned, or frozen (Niazi and Ghafoor 2021). Therefore, here, I present three points that support the use of edible mushrooms to meet the challenge of improving human diets and food security. First, mushrooms improve the quality of the human diet, as they are high in protein, vitamins, and minerals and low in fat. In addition, they are rich in dietary fiber and are also recognized as prebiotic and antioxidant agents (Niazi and Ghafoor 2021). Second, growing edible mushrooms contributes to a sustainable food production system (circular food chains) because people may use agronomic and urban wastes to prepare suitable substrates for mushroom cultivation (Grimm *et al.* 2021). Third, funga,

represented by wild edible mushrooms, plays a central role in conserving the traditional knowledge of peoples and the biodiversity of the planet due to the cooperative relationship of fungi with fauna and flora (Oyanedel *et al.* 2022).

Nelson Menolli Jr. e Mariana de Paula Drewinski: Unleashing the Potential of Wild Edible Mushrooms in Brazil

Wild edible mushrooms (WEM) are often overlooked as a nonwood forest product, but they are an important resource that has gained attention in recent years. While traditional use of WEM by indigenous and traditional groups in the Amazon region has been documented (Vargas-Isla *et al.* 2013), their consumption is still not widely practiced in Brazil. However, interest in foraging for fungi and mushrooms has been growing, and access to information on WEM has increased with the publication of reports, books, and species guides in Portuguese, as well as scientific outreach profiles on social media. Some mycological tourism initiatives have also been conducted, primarily in southern and southeast Brazil. There are approximately 400 species of WEM in Brazil, with approximately 80 species consistently recorded in the country (Drewinski 2023). Even in nonnative forest plantations, such as those of *Pinus* and *Eucalyptus* species, it is possible to find ectomycorrhizal mushroom species that can be used as a food resource. In addition to the species that are already known, many others remain undiscovered and could potentially become new food resources. Mushrooms are a limited, seasonal resource that represents an important phase in the life cycle of a fungus. Therefore, studies on the domestication and cultivation of wild species are important for diversifying mushroom production, *ex situ* conservation of genetic resources, and food sovereignty. WEM are related to at least 11 of the 17 goals of the 2030 Agenda for Sustainable Development (Pérez-Moreno *et al.* 2021). Due to its great ecological, sociocultural, economic, medicinal, and biotechnological significance, WEM can promote forest sustainability, mitigate greenhouse gas emissions, conserve biodiversity, reduce hunger, improve human nutrition and health, conserve traditional knowledge, empower women, and stimulate economic development (Pérez-Moreno *et al.* 2021). Despite the importance and growing interest in WEM, it is crucial to note that recognizing edible species requires a great deal of knowledge and skill. Therefore, people should not consume wild mushrooms without first ensuring their specific identification and previous edibility records.

WILD MEAT

Guillermo Ros Brull: Navigating the Controversial Benefits and Challenges of Wildmeat as a Source of Food Security

In various parts of the world, wildmeat serves as a primary source of protein (Fa *et al.* 2002) and micronutrients (Sarti *et al.* 2015) and plays a crucial role as a source of income for indigenous peoples and local communities (Coad *et al.* 2019). In addition, the process of finding, pursuing, hunting, and consuming wildmeat is closely tied to local ecological knowledge and influenced by cultural norms. However, in the post-COVID-19 context, various stakeholders (FAO 2020b) have argued for a ban on subsistence hunting to prevent future zoonotic disease spillover (Andersen *et al.* 2020) and to address the current trend of wildlife loss caused by large-scale commercial hunting for wildmeat in urban areas (Nasi *et al.* 2011). While a ban on hunting may seem to be a straightforward solution, it would threaten food security in tropical forest areas where there are no alternative food sources. Similarly, transitioning to cattle farming in these areas would require the clearing of tropical forests, leading to the loss of habitat for wildlife (Booth *et al.* 2021), an increase in the emergence of new diseases (Allen *et al.* 2017), and the erosion of indigenous and local knowledge and value systems (van Vliet and Mbazza 2011). Therefore, the sustainable use of wildmeat represents an opportunity to reconcile and achieve both sustainable development and conservation goals (Blaikie 2005). To take advantage of this opportunity, we must (1) devolve decision-making and resource control to local communities, (2) consult with local and national governments and civil society to apply international frameworks in a context-specific manner, and (3) develop conservation strategies that consider both scientific evidence and local sociocultural values and practices (Armitage 2005).

Patricia Carignano Torres: Wildmeat: An Often-Overlooked Resource to Avoid Hidden Hunger

Hunger is estimated to affect approximately 800 million people worldwide (FAO 2021c), but hidden hunger - deficiencies in micronutrients - affects billions of people. These deficiencies can lead to health impairments such as iron-deficiency anemia (IDA), which can compromise physical and cognitive development (Micronutrient Initiative 2009). Animal-source foods (ASFs) can be critical in avoiding certain micronutrient deficiencies because they have high levels of micronutrients that are more bioavailable than vege-

table sources. Wildmeat, which has not been well researched, can therefore improve nutrition and health, particularly in the context of some traditional food systems and given that billions of people cannot afford healthy diets (FAO 2021c). Wildmeat consumption has been shown to increase micronutrient intake, such as iron and zinc, in urban and peri-urban Amazonian households (Sarti *et al.* 2015). The authors argue that households not consuming wildmeat were at greater risk of anemia in the short term and other chronic health problems in the long term. In fact, the only two studies that directly investigated wildmeat consumption and health outcomes found an association with a lower prevalence of anemia. In rural Madagascar, in a context of low ASF consumption, removing access to wildmeat would increase the number of children with IDA by 29%, three times if only considering children in the poorest households (Golden *et al.* 2011). Even in a context with high fish consumption, in Brazilian Amazonia, wildmeat consumption was associated with higher hemoglobin concentrations in children from extremely poor rural households (Torres *et al.* 2022). The authors estimated that the prevalence of IDA would increase by 10% if wildmeat became unavailable. More research is needed to investigate the contribution of wildmeat to health outcomes, but the evidence thus far suggests its importance for vulnerable populations living in forested areas. Replacing wildmeat with domesticated meat is often infeasible in contexts of monetary poverty and can also be environmentally damaging due to land conversion (Nunes *et al.* 2019). Therefore, sustainable management of wildmeat offtake is crucial.

ALGAE OR SEAWEED

Lara Silva, Alice Medeiros, and Michelle Jacob: The Untapped Potential of Algae: Exploring the Benefits of this Sustainable and Nutritious Food Source

Algae are important for the health of the oceans because they are a primary source of oxygen and food for many marine species. Furthermore, algae can be an important food resource for humans. Edible algae can be found in a variety of aquatic environments, including oceans, lakes, and rivers. They are rich in nutrients and have been used as a medicine and food source in many parts of the world for centuries, as is the case for some Asian countries and islands worldwide. (Tiitii *et al.* 2022). Some common types of edible algae include kelp (*Laminaria* spp. and *Macrocystis* spp.), nori (*Porphyra* spp.), and spirulina (*Arthrospira* spp.). Algae can be consumed fresh, dried, or processed in a variety of ways, e.g.,

cooked, fried, boiled, seasoned or in algae-based products (Pérez-Lorens 2019). As the world debates the transition to sustainable food systems, edible algae could play an important role in the development of more sustainable and nutritious food sources. There are two reasons for this. First, algae can be grown with minimal inputs and have a high productivity rate due to their short life cycle (FAO 2018). Second, they are also a good source of nutrients and can be used as a source of protein and other micronutrients, such as vitamin B12 (Kwak *et al.* 2010; Rauma *et al.* 1995), a vitamin previously thought to be exclusive to animal-origin foods, as well other micronutrients (El Din and El-Sherif 2012; Ito *et al.* 2016), such as calcium and iron. Further research is necessary to gain a deeper understanding of the socioeconomic and cultural characteristics of food consumers to promote the global consumption of algae, particularly in contexts where the sustainable production or harvesting of this food resource is a viable opportunity.

FISH

Priscila Lopes: Fish Biodiversity: A Basic-Needs Grocery Package in the Water

Consumption of aquatic animals, mostly finfish, rose from 9.9 kg in the 1960s to 20.2 kg/capita/year in 2020, an annual increase of 1.4% mainly driven by upper-middle-income countries (FAO 2022). This increase in demand has been met with the support of aquaculture, as many economically relevant fish stocks show worrying signs across the globe (Pauly and Zeller 2016). Wild fish still provide 44% of the world's fish intake (FAO 2022), with marked differences between fish consumed by wealthier classes and fish consumed by the poor. While richer countries and richer people within various countries may prefer to consume a few species that are mild in taste, popular, or easy to handle (Lopes *et al.* 2017), the poor rely on a much greater diversity of species, particularly in freshwater environments (Kolding *et al.* 2019). For 3.3 billion people, aquatic animals provide approximately 20% of their protein intake, which can be as high as 50% or more in some countries and regions (FAO 2022). In places in the Brazilian Amazon, for example, fish intake can reach more than 800 g of fish/capita/day (Corrêa *et al.* 2014). For the poor living by water bodies, fish may be the most important source of animal protein and essential micronutrients (Funge-Smith and Bennett 2019; Hicks *et al.* 2019). As various threats to biodiversity relentlessly advance (e.g., unchecked adoption of the Blue Economy agenda, large river dams, and climate change), not only are species lost but risks to food security

and the sovereignty of already vulnerable populations increase (Lopes *et al.* 2021). In addition to the environmental, health, and cultural consequences, the loss of fish biodiversity means that vulnerable people are further impoverished, as they need to divert part of their income to acquire an important part of their basic-needs grocery package through conventional markets.

UNCONVENTIONAL FOOD PLANTS

Rafael Vasconcelos: Maximizing the Potential of Unconventional Food Plants for Biocultural Conservation through Commercialization

Unconventional food plants (UFP) have gained popularity worldwide, particularly in the context of transitioning to sustainable food systems. UFPs are little-used or little-known plant species, mainly outside of their natural habitats. These plants are often associated with traditional knowledge and have previously been recognized by researchers under different names, highlighting their importance in food security strategies (Simonetti *et al.* 2021). Recently, the use of these plants by renowned chefs and their promotion on various media platforms has sparked growing interest and curiosity among potential consumers. This presents a promising opportunity for businesses, particularly in the context of supporting family farming and sustainable extractivism. Therefore, it is important to consider the potential for UFP to generate income while also promoting biocultural conservation. The idea of conserving biodiversity through the commercialization of its resources has been debated for decades (Evans 1993). However, the complex relationships involved in markets for biodiversity products make it difficult to automatically associate them with sustainability or degradation, although the latter seems to be more common. When seeking to evaluate the potential for promoting the commercialization of UFP as a biocultural conservation strategy, there are several questions that need to be considered, including the following: what is the influence of the inherent features of the products, such as flavor, nutritional value, and perishability, on their marketing potential? How do ecological and management aspects, such as the conservation status of natural populations, collection site, method of production, and ease of cultivation, impact their production potential in a sustainable manner? Do social aspects, such as the production organization, market institutional relationships, land and resource tenure, and governance, contribute to the viability of commercialization? How do economic factors, such as investment capacity, ne-

gotiated amounts, and profit sharing along the value chain, impact the viability of the business?

A TRANSITION TOWARD BIODIVERSITY AND FOOD SECURITY IN FOOD SYSTEMS

In Figure 3, we summarize three main pathways that, according to these experts, could guide the transition toward biodiversity and food security in food systems.

It is important to recognize that biodiversity and food security are interconnected and can support each other and to take a holistic approach to promoting both. As we continue to prioritize and invest in education, research, and governance related to biodiversity and food security, we hope that we can take meaningful steps toward realizing and effectively utilizing the potential of biodiversity to contribute to food security in a way that benefits both nature and people. By working together and focusing our collective attention on these issues, we can create positive change and build a more sustainable and secure future for all forms of life on the planet.

FINANCIAL SUPPORT

The authors of this study would like to express gratitude for the financial support received from various organizations: (1) MPD thanks the ‘Fundação de Amparo à Pesquisa do Estado de São Paulo’ (FAPESP), Brazil for the grant #2017/25754-9; (2) NMJ thanks FAPESP for grant #2018/15677-0 and the ‘Conselho Nacional de Desenvolvimento Científico e Tecnológico’ (CNPq), Brazil for the Research Productivity grant 314236/2021-0; (3) PCT thanks Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the postdoctoral grant 001; (4) PFML thanks CNPq for the productivity grant (301515/2019-0). Additionally, this study was funded by the CNPq through a research grant to MCMJ (402334/2021-3) and a research productivity scholarship also awarded to MCMJ (306755/2021-1). The funders played no role in the study design, data collection and analysis, or the decision to publish or prepare the manuscript.




Figura 3. Strategies for Transforming Food Systems to Promote Biodiversity and Food Security in Current Food Systems.

REFERENCES

- Allen T, Murray KA, Zambrana-Torrel C, Morse SS, Rondinini C, Di Marco M, Breit N, Olival KJ, Daszak P (2017) **Global hotspots and correlates of emerging zoonotic diseases.** *Nature Communications* 8.
- Andersen KG, Rambaut A, Lipkin I, Holmes EC, Garry RF (2020) **The proximal origin of SARS-CoV-2.** *Nature medicine* 2–4.
- Aragão A, Contini E (2021) **O Agro no Brasil e no mundo: uma síntese do período de 2000 a 2020.** Embrapa SIRE, Brasília, DF. <https://www.embrapa.br/documents/10180/62618376/0+AGRO+NO+BRASIL+E+NO+MUNDO.pdf/41e20155-5cd9-f4ad-7119-945e147396cb>.
- Armitage D (2005) **Adaptive Capacity and Community-Based Natural Resource Management.** *Environmental Management* 35:703–715.
- Basta P, Orellana J (2020) **Pesquisa sobre os determinantes sociais da desnutrição de crianças indígenas de até 5 anos de idade em oito aldeias inseridas no Distrito Sanitário Especial Indígena (DSEI) Yanomami.** Fiocruz, Rio de Janeiro. <https://drive.google.com/file/d/150A1frAey8kPSYrKTTL-Mdw9Uymt1nZ7/view>.
- Belik W (2020) **Estudo sobre a cadeia de alimentos.** Instituto de Manejo e Certificação Florestal e Agrícola - Imaflora, https://www.ibirapitanga.org.br/wp-content/uploads/2020/10/EstudoCadeiaAlimentos_%C6%92_13.10.2020.pdf.
- Béné C, Oosterveer P, Lamotte L, Brouwer ID, de Haan S, Prager SD, Talsma EF, Houry CK (2019) **When food systems meet sustainability – Current narratives and implications for actions.** World Development .
- Blaikie P (2005) **Community-based natural resource management in Malawi and Botswana** *. Rural Livelihoods and Poverty Reduction Policies. Routledge, p.
- Blicharska M, Smithers RJ, Mikusiński G, Rönnbäck P, Harrison PA, Nilsson M, Sutherland WJ (2019) **Biodiversity’s contributions to sustainable development.** *Nature Sustainability* 2:1083–1093.
- Bodmer R, Mayor P, Antunez M, Fang T, Chota K, Yuyarima TA, Flores S, Cosgrove B, López N, Pizuri O, Puertas P (2020) **Wild Meat Species, Climate Change, and Indigenous Amazonians.** *Journal of Ethnobiology* 40:218–233.

- Booth H, Clark M, Milner-Gulland EJ, Amponsah-Mensah K, Antunes AP, Brittain S, Castilho LC, Campos-Silva JV, Constantino P de AL, Li Y, Mandoloma L, Nneji LM, Iponga DM, Moyo B, McNamara J, Rakotonarivo OS, Shi J, Tagne CTK, van Velden J, Williams DR (2021) **Investigating the risks of removing wild meat from global food systems.** *Current Biology* 31:1788-1797.e3.
- Brashares JS, Arcese P, Sam MK, Coppolillo PB, Sinclair ARE, Balmford A (2004) **Bushmeat hunting, wildlife declines, and fish supply in West Africa.** *Science* (New York, N.Y.) 306:1180–1183.
- Clapp J, Moseley W (2020) **This food crisis is different COVID 19 and the fragility of the neoliberal food security order.** *Journal of Peasant Studies*.
- Coad L, Fa J, Abernethy K, van Vliet N, Santamaria C, Wilkie D, El Bizri H, Ingram D, Cawthorn D-M, Nasi R (2019) **Toward a sustainable, participatory and inclusive wild meat sector.** Center for International Forestry Research (CIFOR), Bogor, Indonesia. <https://www.cifor.org/knowledge/publication/7046>.
- CONAB (2022) **Acompanhamento da safra brasileira: Primeiro levantamento de grãos 2022/23.**
- Coomes OT, Takasaki Y, Abizaïd C, Barham BL (2010) **Floodplain fisheries as natural insurance for the rural poor in tropical forest environments: evidence from Amazonia.** *Fisheries Management and Ecology* 17:513–521.
- Corrêa MA de A, Kahn JR, Freitas CE de C (2014) **Perverse incentives in fishery management: The case of the defeso in the Brazilian Amazon.** *Ecological Economics* 106:186–194.
- Crippa M, Solazzo E, Guizzardi D, Monforti-Ferrario F, Tubiello F, Leip A (2021) **Food systems are responsible for a third of global anthropogenic GHG emissions.** *Nature Food* 2:1–12.
- Cruz-Garcia GS, Sachet E, Vanegas M, Piispanen K (2016) **Are the major imperatives of food security missing in ecosystem services research?** *Ecosystem Services* 19:19–31.
- Drewinski MP (2023) **Cogumelos comestíveis da Mata Atlântica: diversidade e viabilidade de cultivo.** PhD Thesis, Instituto de Pesquisas Ambientais, São Paulo, BR.
- El Bizri HR, Morcatty TQ, Valsecchi J, Mayor P, Ribeiro JES, Vasconcelos Neto CFA, Oliveira JS, Furtado KM, Ferreira UC, Miranda CFS, Silva CH, Lopes VL, Lopes GP, Florindo CCF, Chagas RC, Nijman V, Fa JE (2020) **Urban wild meat consumption and trade in central Amazonia.** *Conservation Biology: The Journal of the Society for Conservation Biology* 34:438–448.
- El Din NGS, El-Sherif ZM (2012) **Nutritional value of some algae from the northwestern Mediterranean coast of Egypt.** *Journal of Applied Phycology* 24:613–626.
- ENANI (2021) **Biomarcadores do estado de micronutrientes: prevalências de deficiências e curvas de distribuição de micronutrientes em crianças brasileiras menores de 5 anos.** Universidade Federal do Rio de Janeiro, Rio de Janeiro. https://enani.nutricao.ufrj.br/wp-content/uploads/2021/10/Relato%CC%81rio3_ENANI-2019_Biomarcadores_vfinal-1-3.pdf.
- Endo W, Peres CA, Haugaasen T (2016) **Flood pulse dynamics affects exploitation of both aquatic and terrestrial prey by Amazonian floodplain settlements.** *Biological Conservation* 201:129–136.
- Evans M (1993) **Conservation by commercialization. Tropical forests, people and food: bio-cultural interactions and applications to development.** Hladik CM, Hladik A, Linares OF, Pagezy H, Semple A, Hadley M., Paris: Unesco, pp. 815-822.
- Fa JE, Peres CA, Meeuwig J (2002) **Bushmeat Exploitation in Tropical Forests: An Intercontinental Comparison.** *Conservation Biology* 16:232–237.
- FAO - Food and Agriculture Organization of the United Nations (1996) **Draft declaration of the world summit on food security.** FAO, Rome.
- FAO (2010) **Second Report on the State of the World's Plant Genetic Resources for Food and Agriculture.** Rome.
- FAO (2017) **Guidelines on assessing biodiverse foods in dietary intake surveys.** Rome.
- FAO (2018) **The State of World Fisheries and Aquaculture: Meeting the Sustainable Development Goals.** Rome.
- FAO (2020a) **Consumer price indices and food inflation.** Rome.
- FAO (2020b) **White Paper: Build back better in a post-COVID-19 world – Reducing future wildlife-borne spillover of disease to humans - CBFP - Congo Basin Forest Partnership.** FAO, Rome.
- FAO (2021a) **Indigenous Peoples' food systems: Insights on sustainability and resilience from**

the front line of climate change. Rome.

FAO (2021b) **The White/Wiphala Paper on Indigenous Peoples' food systems.** Rome.

FAO (2021c) **The State of Food Security and Nutrition in the World - SOFI.** FAO, Rome. [<https://www.fao.org/documents/card/en/c/cb4474en>].

FAO (2022) **The State of World Fisheries and Aquaculture 2022: toward blue transformation.** Rome.

Fischer J, Abson D, Bergsten A, French N, Dorresteijn I, Hanspach J, Hylander K, Schultner J, Senbeta F (2017) **Reframing the Food– Biodiversity Challenge.** *Trends in Ecology & Evolution* 32:335–345.

Funge-Smith S, Bennett A (2019) **A fresh look at inland fisheries and their role in food security and livelihoods.** *Fish and Fisheries* 20:1176–1195.

Garzillo J, Machado P, Louzada M, Levy R, Monteiro C (2019) **Pegadas dos alimentos e das preparações culinárias consumidos no Brasil.** Universidade de São Paulo, São Paulo.

Golden CD, Fernald LCH, Brashares JS, Rasolofoniaina BJR, Kremen C (2011) **Benefits of wildlife consumption to child nutrition in a biodiversity hotspot.** *Proceedings of the National Academy of Sciences* 108:19653–19656.

Golden CD, Vaitla B, Ravaoliny L, Vonona MA, Gasta Anjaranirina EJ, Randriamady HJ, Glahn RP, Guth SE, Fernald LCH, Myers SS (2019) **Seasonal trends of nutrient intake in rainforest communities of northeastern Madagascar.** *Public Health Nutrition* 22:2200–2209.

Grimm D, Kuenz A, Rahmann G (2021) **Integration of mushroom production into circular food chains.** *Organic Agriculture* 11:309–317.

Hicks CC, Cohen PJ, Graham NAJ, Nash KL, Allison EH, D'Lima C, Mills DJ, Roscher M, Thilsted SH, Thorne-Lyman AL, MacNeil MA (2019) **Harnessing global fisheries to tackle micronutrient deficiencies.** *Nature* 574:95–98.

Hunter D, Borelli T, Gee E (2020) **Biodiversity, Food and Nutrition: A New Agenda for Sustainable Food Systems.** Routledge, London.

IPBES - The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (2019) **The global assessment report on biodiversity and ecosystem services - summary for policy makers.** IPBES, Bonn, Germany.

IPCC - Intergovernmental Panel on Climate Change

(2019) **Summary for policymakers: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems.** <https://www.ipcc.ch/srccl/chapter/summary-for-policymakers>.

Ito F, Bernard E, Torres RA (2016) **What is for dinner? First report of human blood in the diet of the hairy-legged vampire bat *Diphylla ecaudata*.** *ACTA CHIROPTEROLOGICA* 18:509–515.

Karjalainen E, Sarjala T, Raitio H (2010) **Promoting human health through forests: overview and major challenges.** *Environmental Health and Preventive Medicine* 15:1–8.

Kennedy G, Stoian D, Hunter D, Kikulwe E, Termote C, Alders R, Burlingame B, Jamnadass R, Thilsted S (2017) **Food biodiversity for healthy, diverse diets. Mainstreaming Agrobiodiversity in Sustainable Food Systems: Scientific Foundations for an Agrobiodiversity Index.** *Biodiversity International*.

Khoury CK, Bjorkman AD, Dempewolf H, Ramirez-Villegas J, Guarino L, Jarvis A, Rieseberg LH, Struik PC (2014) **Increasing homogeneity in global food supplies and the implications for food security.** *Proceedings of the National Academy of Sciences of the United States of America* 111:4001–4006.

Kolding J, van Zwieten P, Marttin F, Funge-Smith S, Poulain F (2019) **Freshwater small pelagic fish and their fisheries in major African lakes and reservoirs in relation to food security and nutrition.** FAO, Rome.

Kuhnlein H, Erasmus B, Spigelski D, Burlingame B (2013) **Indigenous peoples' food systems and well-being: interventions and policies for healthy communities.** Centre for Indigenous Peoples' Nutrition and Environment (CINE), Quebec; FAO, Rome.

Kuhnlein HV, Erasmus B, Spigelski D (2009) **Indigenous Peoples' Food Systems: the many dimensions of culture, diversity and environment for nutrition and health.** Centre for Indigenous Peoples' Nutrition and Environment (CINE), Quebec; FAO, Rome.

Kwak CS, Lee MS, Lee HJ, Whang JY, Park SC (2010) **Dietary source of vitamin B(12) intake and vitamin B(12) status in female elderly Koreans aged 85 and older living in rural area.** *Nutrition research and practice* 4:229–234.

Lachat C, Raneri JE, Smith KW, Kolsteren P, Van

- Damme P, Verzelen K, Penafiel D, Vanhove W, Kennedy G, Hunter D, Odhiambo FO, Ntandou-Bouzitou G, De Baets B, Ratnasekera D, Ky HT, Remans R, Termote C (2018) **Dietary species richness as a measure of food biodiversity and nutritional quality of diets.** *Proceedings of the National Academy of Sciences of the United States of America* 115:127–132.
- Lopes PFM, de Freitas CT, Hallwass G, Silvano RAM, Begossi A, Campos-Silva JV (2021) **Just Aquatic Governance: The Amazon basin as fertile ground for aligning participatory conservation with social justice.** *Aquatic Conservation: Marine and Freshwater Ecosystems* 31:1190–1205.
- Lopes PFM, Mendes L, Fonseca V, Villasante S (2017) **Tourism as a driver of conflicts and changes in fisheries value chains in Marine Protected Areas.** *Journal of Environmental Management* 200:123–134.
- Meli P, Rey-Benayas J, Brancalion P (2019) **Balancing land sharing and sparing approaches to promote forest and landscape restoration in agricultural landscapes: Land approaches for forestlandscape restoration.** *Perspectives in Ecology and Conservation* 17:201–205.
- Micronutrient Initiative (2009) **Investing in the future: a united call to action on vitamin and mineral deficiencies: global report, 2009.** Micronutrient Initiative, New Delhi.
- Nasi R, Taber A, Van Vliet N (2011) **Empty forests, empty stomachs? Bushmeat and livelihoods in the Congo and Amazon Basins.** *International Forestry Review* 13:355–368.
- Niazi AR, Ghafoor A (2021) **Different ways to exploit mushrooms: A review.** *All Life* 14:450–460.
- Nunes AV, Peres CA, Constantino P de AL, Santos BA, Fischer E (2019) **Irreplaceable socioeconomic value of wild meat extraction to local food security in rural Amazonia.** *Biological Conservation* 236:171–179.
- Nyahongo JW, Holmern T, Kaltenborn BP, Røskaft E (2009) **Spatial and temporal variation in meat and fish consumption among people in the western Serengeti, Tanzania: the importance of migratory herbivores.** *Oryx* 43:258–266.
- Oyanedel R, Hinsley A, Dentinger BTM, Milner-Gulland E j., Furci G (2022) **A way forward for wild fungi in international sustainability policy.** *Conservation Letters* 15:e12882.
- Pauly D, Zeller D (2016) **Catch reconstructions reveal that global marine fisheries catches are higher than reported and declining.** *Nature Communications* 7:10244.
- PENSSAN - Rede Brasileira de Pesquisa em Soberania e SAN (2021) **I VIGISAN - Inquérito Nacional sobre Insegurança Alimentar no Contexto da Pandemia da Covid-19 no Brasil.** Fundação Friedrich Ebert, São Paulo, SP.
- PENSSAN (2022) **II VIGISAN - Inquérito Nacional sobre Insegurança Alimentar no Contexto da Pandemia da Covid-19 no Brasil.** São Paulo, SP.
- Pérez-Lorens J (2019) **Seaweed Consumption in the Americas.** *Gastronomica* 49–59.
- Pérez-Moreno J, Guerin-Laguette A, Rinaldi AC, Yu F, Verbeken A, Hernández-Santiago F, Martínez-Reyes M (2021) **Edible mycorrhizal fungi of the world: What is their role in forest sustainability, food security, biocultural conservation and climate change?** *PLANTS, PEOPLE, PLANET* 3:471–490.
- Rauma A-L, Torronen R, Hanninen O, Mykkanen H (1995) **Vitamin B-12 status of long-term adherents of a strict uncooked vegan diet ('living food diet') is compromised.** *Journal of Nutrition* 125:2511–2515.
- Raven P, Gereau R, Phillipson P, Chatelain C, Jenkins C, Ulloa C (2020) **The distribution of biodiversity richness in the tropics.** *Science Advances*.
- Rosen S, Shapouri S (2009) **Global Economic Crisis Threatens Food Security in Lower Income Countries.** <https://www.ers.usda.gov/amber-waves/2009/December/global-economic-crisis-threatens-food-security-in-lower-income-countries> Accessed December 28, 2022.
- Sarti FM, Adams C, Morsello C, van Vliet N, Schor T, Yagüe B, Tellez L, Quiceno-Mesa MP, Cruz D (2015) **Beyond protein intake: Bushmeat as source of micronutrients in the amazon.** *Ecology and Society* 20.
- SEEG - System of Estimates of Emissions and Removals of Greenhouse Gases (2019) **Análise das emissões de gases de efeito estufa e suas implicações para as metas do Brasil (1970-2018).** Observatório do Clima, Brasília/DF.
- Simonetti M, Simonetti K, Fariña L (2021) **Biodiversity as sustainability: possibility of markets for nonconventional food plants (PANC).** *Brazilian Journal of Development* 7 7:35330–35348.

The World Bank (2022) **Food exports (% of merchandise exports)**. <https://data.worldbank.org/indicator/TX.VAL.FOOD.ZS.UN>.

Tiitii U, Paul N, Burkhart S, Larson S, Swanepoel L (2022) **Traditional Knowledge and Modern Motivations for Consuming Seaweed (Limu) in Samoa**. *Sustainability* 14:6212.

Torres PC, Morsello C, Orellana JDY, Almeida O, Moraes AD, Montalván EAC, Pinto MAT, Fink MGS, Freire MP, Parry L (2022) **Wildmeat consumption and child health in Amazonia**. *Scientific Reports* 1–14.

Travassos GF, Antônio da Cunha D, Coelho AB (2020) **The environmental impact of Brazilian adults' diet**. *Journal of Cleaner Production* 272:122622.

Tregidgo D, Barlow J, Pompeu PS, Parry L (2020) **Tough fishing and severe seasonal food insecurity in Amazonian flooded forests**. *People and Nature* 2:468–482.

Tregidgo D, Piperata B, Lima J, Inglez M, Valsecchi J (In review) **Variation in dietary resilience among Amazonian populations living in a context of dramatic seasonal flooding**. *Human Ecology*.

UNEP - The United Nations Environment Programme (1992) **Convention on Biological Diversity**. Rio de Janeiro.

UNEP (2020) **Preventing the Next Pandemic: Zoonotic diseases and how to break the chain of transmission**. UNEP, Nairobi.

Vaitla B, Devereux S, Swan SH (2009) **Seasonal Hunger: A Neglected Problem with Proven Solutions**. *PLOS Medicine* 6:e1000101.

Vargas-Isla R, Ishikawa NK, Py-Daniel V (2013) **Contribuições etnomicológicas dos povos indígenas da Amazônia**. *Biota Amazônia (Biote Amazonie, Biota Amazonia, Amazonian Biota)* 3:58–65.

Vasconcelos-Neto C, Jacob M, Tregidgo D, Valle D, Monteiro F, Scofield A, Mayor P (in press) **Food insecurity increases across South America due to the COVID-19 pandemic**.

Verly Jr E, de Carvalho AM, Marchioni DML, Darmon N (2022) **The cost of eating more sustainable diets: A nutritional and environmental diet optimization study**. *Global Public Health* 17:1073–1086.

van Vliet N, Mbazza P (2011) **Recognizing the Multiple Reasons for Bushmeat Consumption in Urban Areas: A Necessary Step Toward the Sustainable Use of Wildlife for Food in Central Africa**. *Human Dimensions of Wildlife* 16:45–54.

Received: 29 December 2022

Accepted: 01 February 2023

Published: 10 February 2023

Editor: Ulysses Albuquerque