

Fishers' knowledge on the ecology, impacts and benefits of the non-native peacock bass *Cichla kelberi* in a coastal river in southeastern Brazil

Paula Araujo Catelani^{1*}; Ana Cristina Petry²; Fernando Mayer Pelicice³ and Renato Azevedo Matias Silvano⁴

ABSTRACT

Ethnoichthyology has been used to understand how humans perceive environmental changes, including species introductions. In Rio de Janeiro State, Brazil, fishers introduced juveniles of peacock bass *Cichla kelberi* in the Juturnaíba Reservoir in the 1990's. In the last years, this fish started to be caught in the São João River, downstream of the dam. This study aimed to identify whether fishers recognize the potential impact of the peacock bass over the native species, to understand the economic valuation they made to the services provided by this species, and if they include management of the non-native peacock bass in their practices. Semi-structured interviews with artisanal and sport fishers were employed between September and October 2016. A total of twenty-five interviews were carried out, 12 with artisanal and 13 with sport fishers. In general, fishers expressed knowledge about biological and behavioral aspects of the peacock bass, but opinions diverged on its impact over the native ichthyofauna. Differently from the artisanal, most sport fishers do not relate the decline in populations of native species to the occurrence of the peacock bass. Instead, they mention other anthropogenic impacts in the river basin. The later advocate the practice of fish and release in order to maintain a sustainable population of peacock bass in the study area. The ethnoichthyological knowledge gathered from fishers provide novel information of the native ichthyofauna, peacock bass introduction and other anthropogenic impacts, and might constitute a powerful tool to the development of sustainable strategies of the natural resources.

Keywords: Ethnobiology; Artisanal Fishing; Sport Fishing; Introduced Fishes; Juturnaíba Reservoir; São João River.

1 Universidade Federal do Rio de Janeiro, campus UFRJ - Macaé Professor Aloísio Teixeira, Programa de Pós-Graduação em Ciências Ambientais e Conservação, Av. São José do Barreto, 764, CEP 27965-045, Macaé, RJ, Brazil.

2 Universidade Federal do Rio de Janeiro, Instituto de Biodiversidade e Sustentabilidade, CP 119331, CEP 27910-970, Macaé, RJ, Brazil.

3 Núcleo de Estudos Ambientais, Universidade Federal do Tocantins, Rua 3, Quadra 17, CEP 77500-000 Porto Nacional, TO, Brazil.

4 Departamento de Ecologia, Universidade Federal do Rio Grande do Sul, CP 15007, CEP 91501-970, Porto Alegre, Rio Grande do Sul, Brazil.

* Corresponding author ✉. E-mail address: PAC (ktelani@gmail.com), ACP (petryanacristina@gmail.com), FMP (fmpelicice@gmail.com), RAMS (00132878@ufrgs.br)

SIGNIFICANCE STATEMENT

The assessment of the fishers' knowledge about biological invasion in a coastal dammed river in Southeast Brazil revealed that artisanal fishers are aware of the decline of native species and recognize the increase in non-native species over time. Sport fishers, on the other hand, expressed more details of reproductive biology of the non-native *Cichla kelberi*, their target species. A local sport fisher provides novel information from the first *C. kelberi* introductions in the year 2001. Sport and artisanal fishers acknowledge that by practicing fish and release they contribute to the maintenance of sustainable population of this non-native species. This study reinforces that although differing in the purpose of fisheries, users of the same invaded ecosystems may provide complementary and useful information for the development and management practices of both native and non-native species.

INTRODUCTION

The discipline of Human Ecology addresses the diversity of interactions established by human cultures with the environment and their knowledge about the natural resources, which is accumulated and transferred through the generations (Begossi *et al.* 2000; Berkes 1999). As part of Human Ecology, the Ethnobiology investigates the local or traditional ecological knowledge in order to understand how human populations perceive, classify and use the natural resources (Begossi and Garavello 1990; Berkes 1999; Porcher *et al.* 2010). Ethnobiological studies can provide a range of information because they reflect the knowledge of human communities about the classification, biology, ecology and behavior of distinct groups of animals and plants (Camacho Guerreiro *et al.* 2016; Mourão and Nordi 2002; Pinto *et al.* 2017).

Artisanal small-scale fisheries require biological knowledge about the explored fishes. The use of fisherfolk knowledge ultimately has improved the use and management of fish as an important source of animal protein along the history of mankind (Fao 2003; Magalhães *et al.* 2011; Marques 1991). Among the several unfolds of Ethnobiology, the Ethnoichthyology focuses on the knowledge, classification and use of fishes as a resource and includes aspects of fish conservation and cultural habits of people (Corneta 2008; Marques 1991; Morrill 1967; Silvano and Begossi 2002). Therefore, the perceptions of fisherfolk about the ecology and behavior of fishes may provide relevant information to guide the conservation and management of ecosystem and environments, including the potential effects of extreme climatic events on fisheries (Camacho Guerreiro *et al.* 2016; Silvano *et al.* 2008). In Brazil, the first ethnoichthyological study was published in the early 1990s (Begossi and Garavello, 1990). Since then, several studies have analyzed fisherfolk knowledge about the spatial and temporal distributions, behavior and trophic ecology of target fishes, as well as the cultural practices and traditions (e.g. Begossi *et al.* 2016; Mourão and Nordi 2002; Pinto *et al.* 2017; Ramires *et al.* 2010; Silvano and Valbo-jørgensen 2008). However, few stud-

ies have applied the local knowledge of fisherfolk to understand the extension of human impacts in rivers (Hallwass *et al.* 2013). River damming associated to the formation of reservoirs for water supply or power generation can cause environmental, social and economic impacts (Agostinho *et al.* 2016; Fearnside 1999; Petrere 1996). Among these impacts, the introduction of non-native species in reservoirs, in addition to facilitating the dispersal and establishment of these species (Johnson *et al.* 2008), has been associated with severe disturbances that vary from reductions in catches to local extinctions of native species (Latini and Petrere Junior 2004; Pelicice and Agostinho 2009). The changes in the habitat caused by the formation of reservoirs and introduction of non-native species can negatively affect the social condition of fishers, who need to adapt to different fishing strategies and target species, most of them with lower economic value (Hoeinghaus *et al.* 2009). However, little is known about the fisherfolk perception and reaction to changes in the freshwater environment and in the availability and alteration of fishing resources, including the presence of non-native fishes (Hallwass *et al.* 2013; Huntington *et al.* 2017; Silvano and Begossi 2002).

Amazonian fishes popularly known as peacock basses (English) or "tucunarés" (Brazilian Portuguese) (Teleostei: Cichlidae, *Cichla* spp.), have been introduced in several impounded rivers where they succeed in the colonization of reservoirs, experiencing rapid population increases (Espínola *et al.* 2010; Pelicice and Agostinho 2009). Species of peacock basses were recorded in at least 35% of the 71 Brazilian reservoirs evaluated by Agostinho *et al.* (2007), being the dominant species of fish in 10% of these. Drastic reductions in the diversity of native fish species have been attributed to the occurrence of peacock bass in systems where they have been introduced (Britton and Orsi 2012; Latini and Petrere Junior 2004; Pelicice *et al.* 2015; Zaret and Paine 1973). Changes in the composition of the ichthyofauna occurred after the introduction of the peacock bass and other non-native fishes and such changes can affect fishing (Rosa *et al.*, 2014). However, few studies

have been conducted to understand how fishers perceive and interact with non-native species in Brazil (Azevedo-Santos *et al.* 2010; Silvano and Begossi 2002).

Ethnoichthyology may reveal how traditional people that depend on local scale fisheries interact with environmental changes caused by humans, such as the introduction of non-native fishes (Azevedo-Santos *et al.* 2010; Carvalho 2002; Moura and Marques 2007; Moura *et al.* 2008; Pinheiro 2004). Therefore, the present study aimed to investigate how subsistence and recreational fishers (hereafter named respectively as artisanal and sport fisher) perceive and interact with the non-native species of peacock bass *Cichla kelberi*. We also aimed to investigate the perceptions of fishers about the natural resources and the invaded ecosystem in a coastal Neotropical river. In particular, we investigated whether fishers recognize potential impacts of the peacock bass, how they value this non-native species and if they have developed management practices to control or to maintain sustainable population of this fish on the studied system.

MATERIAL AND METHODS

The Juturnaíba Reservoir is located ca. 40 km upstream the São João River mouth, State of Rio de Janeiro, southeastern Brazil (22° 33' S and 42° 18' W) (Figure 1). The reservoir was built in the 1980s, it occupies an area of 43 km², provides water to the populous Região dos Lagos and serves as a recreational and sport fishing area. The São João River originates at 800 m of altitude and runs about 120 km west, until flowing into the Atlantic Ocean in a mangrove-dominated estuary in the district of Barra de São João, in the municipality of Casimiro de Abreu.

Anglers introduced *C. kelberi* in this reservoir during the 1990s (Catelani *et al.* 2020). About five years ago, this fish became frequent in the catches downstream of the reservoir in the last stretch of the São João River, which includes an estuarine area. This portion of the river was already highly modified by human intervention, due to flow regulation and previous channelization of the meandering course, which occurred between 1950 and 1985.

We conducted semi-structured interviews with fishers, and they included closed and open questions. The advantages of this approach is to gather quick and objective interviews and the confidence that important research questions have been addressed (Silvano *et al.* 2008). Two types of questionnaires were employed: one targeted to local artisanal fishers and other to sport fishers. The same person (P. A. Catelani), performed the interviews, which were carried out between September and October 2016 in three locations (Figure 1), in order to include fishers from

Juturnaíba Reservoir and the downstream last stretch of the São João River.

The questionnaires covered (i) general aspects of the fishing activity, (ii) the socio-cultural profile of fishers, (iii) their knowledge about the biology of the peacock bass, (iv) composition and economical valuation of the native and non-native fish fauna, and (v) their awareness about changes in the abundance and composition of fish fauna by anthropogenic interventions (Table 1). The popular and scientific names of the species cited by the fishers during the interview are presented in supplementary material (Additional file 1).

The snowball sampling technique (*sensu* Biernacki and Waldorf 1981) was employed after the first interview. Interviews lasted an average 20 minutes and were carried out in several places of each locality, according to the availability of each interviewee. All interviews were recorded with prior consent of the informants, and the research was previously approved by the Ethics Committee of the Federal University of Rio de Janeiro - campus Macaé, Brazil (CAAE: 58514816.9.0000.5699).

RESULTS AND DISCUSSION

Fishers profile

A total of 25 interviews were carried out with 12 traditional and 13 sport fishers. Three traditional fishers were women (Table 2), the remaining nine were men. On average, artisanal fishers were 48 ± 10 years old. Seven of them conducted their fishing activities only in the Juturnaíba Reservoir and the other five only in the last stretch of the São João River. Most fishers had incomplete elementary education and none had higher education. Almost all fishers were married (91.7%) and all of them had children. Although 75% of the fishers confirmed that their parents were fishers, only 33% of their children currently practice this activity (Table 2).

Among sport fishers, only men were interviewed. Four practice fishing only in the Juturnaíba Reservoir, four only in the last stretch of the São João River, and five in both locations (Table 2). The fishermen visit the places for leisure and are from six cities of the State of Rio de Janeiro, three cities of the State of Minas Gerais State and one of Sate of São Paulo. The average age of sport fishermen was lower than that of artisanal fishers (40 ± 13 years old), as well as the proportion of married (46%) and those who have children (54%). On average, the interviewed sport fishers have been fishing in the São João River for 12 years (± 10.8). In Juturnaíba Reservoir, most sport fishers hire guide and boat services, which include fuel and live baits. The main bait used is the tetra (*Astyanax*




Figure 1. Location of the São João River Basin, State of Rio de Janeiro, Brazil and sampling localities (1-3). (1): Juturnaíba Reservoir, Silva Jardim municipality; (2): São João River downstream of the dam, Casimiro de Abreu municipality; (3): mouth of the São João River, district of Barra de São João, Casimiro de Abreu municipality.

spp.), which was cited by five fishers. The baits are previously caught by the guides with gill nets and stored in tanks. According to these fishermen, unused baits are returned to these tanks.

Considering artisanal fisher people, although most of them have fishing as their main activity, 50% perform other activities to supplement their incomes. Only one person interviewed does not have artisanal fishing as a main occupation (he is a civil worker). Most of them (83%) have one or more family members involved in fishing activities (Table 3).

Sport fishers presented higher educational level (Table 2) and reported 10 different professions (Table 4), two of them directly related to the sport fishing activity. One interviewed owns a company that manufactures artificial baits and the other works as shop assistant in a shop specialized in fishing products.

Fishing activity

Among the five fishing gears mentioned by the artisanal fishers, gillnet was the most used, whereas line and hook with artificial or live baits were mainly used by sport fishers (Figure 2).

In general, artisanal fishers target more species than sport fishers (Figure 3). In the reservoir, the main target species of artisanal fishers were the trahira *Hoplias* aff. *malabaricus* and the peacock bass (Figure 3A), which have higher market value (between US\$ 3.15 and US\$ 4.80/kilogram). Other fishes reported cost, on average US\$ 1.60/kilogram, as the pearl cichlid *Geophagus brasiliensis*, the armored catfish *Loricariichthys castaneus*, and the singing catfish *Trachelyopterus striatulus*. In the downstream section, fishers mentioned the same freshwater species reported in Juturnaíba, in addition to some marine

Table 1. Topics and questions addressed in the questionnaires used in interviews conducted in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

Topics covered	Questions evaluated
Socio-cultural profile of fishers	Sex, age, origin, education, marital status, number of children, parents are or not fishers, children are or not fishers
General aspects of the fishing activity	Frequency and importance of fishing, other activities developed, How many family members involved in fishing, fishing area, time spent fishing, fishing methods and target species
Knowledge about the biology of the peacock bass	Origin, reproductive activity, behaviour, abundance, items consumed and predators of the peacock bass
Interaction with the peacock bass	Importance and management of the peacock bass as a resource, economical valuation of the native and non-native fish fauna
Changes caused by the peacock bass	Changes in tourism and economy, fishes before and after the introduction of the peacock bass
Changes by anthropogenic interventions	Occurrence of other non-native species, impacts perceived by interviewee




Figure 2. Fishing methods used by artisanal ($N = 12$) and sport ($N = 13$) fishers interviewed in the Juturnaíba Reservoir and in the downstream section of the São João River, between September 28 and 08 October 2016.

fishes (Figure 3B). One of these species are the sea basses *Centropomus* spp., which attain the highest market value (between US\$ 6.40 and US\$ 8.00/kilogram). The non-native redbreast tilapia *Oreochromis niloticus* was also frequently cited by the artisanal fishers. Sea catfishes were also reported, probably captured when these fishes enter the river to spawn.

The main target species of the sport fishing in the reservoir was the peacock bass, whereas the sea basses *Centropomus* spp., followed by the peacock bass, were the main species targeted in the stretch downstream of the dam (Figure 3). Even though both species are the most valued by the fishers, their preference of fishing in those areas relies mainly on the occurrence of

the peacock bass in the region.

Interaction with the peacock bass

In general, fishers had a considerable knowledge about the biology of the peacock bass, especially the artisanal ones that work in the reservoir and target this species as the main resource. The recognition that the peacock bass is not-native in the São João River was consensual, except for a young sport fisherman, who answered that this fish always occurred there. Among artisanal fishers, the most common response was that the peacock bass escaped from fish breeding tanks. According to most sport fishes, the peacock bass comes from the Amazon basin (Figure 4). When asked about the time of introduction, artisanal fishers responded that the peacock bass was introduced on average 16 years ago (± 4.2), and most

of them (83.3%) started fishing before its introduction. Only one sport fisherman that was involved in the release of the peacock bass in Juturnaíba provides the more accurate answer and stated that the introduction occurred 15 years ago. According to him, at the time of the first fish release, each sport fisher willing the peacock bass introduction contributed with US\$ 15.00 for the purchase of 5,000 specimens from the State of Mato Grosso, which were released into the reservoir.

Responses were unanimous regarding the food habit, informing that the peacock bass feeds on a wide range of aquatic animals, such as young fish, including conspecifics, and shrimp (Figure 5A). Some fishers emphasized that the peacock bass is a “heavy predator”. In general, sport fishers detailed the preys, citing the tetras, the sea bass and the peacock bass, as well as shrimp and aquatic insects.

Table 2. Profile of the 12 artisanal and 13 sport fishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

	Artisanal fishers(N= 12)	Sport fishers(N= 13)
Sex (%)		
Male	75.0	100.0
Female	25.0	0.0
Age (year)		
Minimum	33.0	23.0
Mean	48.2	40.1
Maximum	69.0	62.0
Education (%)		
None	25.0	0.0
Incomplete fundamental	58.3	7.7
Complete fundamental	0	7.7
Incomplete high school	0	7.7
Complete high school	16.7	38.5
Incomplete higher education	0	7.7
Complete higher education	0	30.8
Place of origin (%)		
Rio de Janeiro State	91.7	69.2
Minas Gerais State	0	23.1
Espírito Santo State	8.3	0.0
São Paulo State	0	7.7
Marital status (%)		
Single	8.3	54.0
Married	91.7	46.0
Children (%)		
No	0.0	46.0
Yes	100	54.0
Parents are also fishers(%)		
No	25.0	23.1
Yes	75.0	76.9
Children are also fishers(%)		
No	66.7	28.6
Yes	33.3	71.4

Table 3. Characteristics of the 12 artisanal fishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

Characteristics	Artisanal fishers(N= 12)
Fishing as main activity (%)	
No	8.3
Yes	91.7
Other activities (%)	
Guide	25.0
Cooker	16.7
Housewifely	8.3
Civil worker	8.3
None	41.7
Frequency of fishing (%)	
Daily	50.0
Weekly	50.0
Family members involved in fishing (%)	
0	16.7
1	33.3
2	16.7
3	25.0
4	8.3
Years fishing in the region	
Minimum	4.0
Mean	27.4
Maximum	48.0

Table 4: Characteristics of the 13 sport fishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

Characteristics	Sport fishers (N= 13)
Activities developed (%)	
Engineer	23.0
Bricklayer	23.0
Businessman (Artificial baits)	7.7
Shop assistant (fishing products)	7.7
Economist	7.7
Communication technician	7.7
Graphic designer	7.7
Firefighter	7.7
Civil servant	7.7
Frequency of fishing (%)	
Daily	15.4
Weekly	30.8
Fortnightly	7.7
Monthly	23.0
Bimonthly	7.7
Annually	15.4
Years fishing in the region	
Minimum	1.0
Mean	12.4
Maximum	43.0

In relation to the predators of the peacock bass, some fishers were not able to cite anyone. The main

predator mentioned was the african catfish *Clarias gariepinus*, which is also introduced in the São João




Figure 3. Target species reported by the 12 artisanal and 13 sportfishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

River, followed by the trahira (Figure 5B). One artisanal fisherman who uses harpoon as fishing gear provided an interesting response. According to him, the African catfish must be the only species that prey on the peacock bass, as juveniles, but without much success, due to the parental care exhibited by peacock bass, which remain near the nest protecting their off-

spring.

All fishers informed that peacock basses form aggregations. However, when asked at which portion of the water column the peacock bass is most commonly found, responses varied and were often related to the water temperature or to the employed fishing method. According to artisanal fishers, the peacock




Figure 4. Origin of the peacock bass according to the 12 artisanal and 13 sport fishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

bass is abundant throughout the year, but with higher catches during the summer, because of greater activity of this fish in higher water temperatures. Respective to abundance, 50% of artisanal fishers and 61.6% of sport fishers of Juturnaíba Reservoir reported that there are usually more juveniles than adults. Along the last stretch of the São João River downstream to the dam, sport fishers' perception differs, since half believe there are more juveniles than adults and the other half said that there are more adults than juveniles.

The reproductive period of the peacock bass was a less known aspect, especially among the artisanal fishers. Nearly half of them did not know when the reproductive period occurs and some said they had not seen peacock bass with eggs. Some fishers answered that the peacock bass reproduces two to three times a year, but they were unsure about the respective months. Summer, followed by spring, was the season most cited by the interviewed fishers (Figure 6).

Environmental changes associated with peacock bass introduction

Artisanal fishers, mainly those that live near Juturnaíba Reservoir, were more specific on their answers concerning alterations due to peacock bass introduction. Only three sport fishermen fished on the site before the introduction of the peacock bass.

When questioned about the tourism directed to peacock bass, artisanal fishers recognized a change in the motivation toward this activity. They reported that before the introduction of the peacock bass, local tourism was targeted to tetras, which attracted a great amount of people that arrived even in chartered buses. At that time, fishing took place on the banks of the reservoir without the need for boats, and people were able to catch a large number of tetras. According to these fishers, the current tourism focused on sport fishing is worse to the local economy than the previous tourism directed to tetras. This lower importance of tourism directed to the peacock bass was associated with low catches of this fish near the shoreline. As the fishing of peacock bass requires boats and guide services, such activity is consequently more expensive for practitioners. Even though it is more expensive and less crowded than tourism targeted to the tetra, peacock sport fishing motivates a large group of participants who agree with the practice of catch and release, aiming at maintaining the regional stock of peacock bass.

Another fact emphasized by most interviewed people irrespective to the site and modality was the change in the composition of the fish fauna after the introduction of the peacock bass and the drastic decrease in the population size of some species, mainly tetras and the toothless characin, due to predation by the peacock bass (Figure 7). The artisanal fishers of Juturnaíba Reservoir agreed that the toothless




Figure 5: Items consumed by the peacock bass fishers (A) and predators of the peacock bass (B) according to the 12 artisanal and 13 sport fishers interviewed in the interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

characin sustained the fishing activity before the introduction of the peacock bass, which were commercialized including in other nearby cities. The sport fisherman who participated in the release of peacock bass in Juturnaíba Reservoir regretted that he has contributed to the introduction of the species in the region. He is aware of the gradual reduction of the abundance of the toothless and the tetras, in addition to the establishment of the peacock bass and the control it exerts on other species the in the section of the river downstream the dam.

Fishers' responses indicated that peacock bass is currently important in Juturnaíba Reservoir, and since its introduction the species is getting more abundant than the native ones that are also fished in the region (Figure 7A). The artisanal fishers of the São João River, on the other hand, do not seem to perceive an increase on the abundance of the peacock bass, probably because this fish occurs in stretches farther away from the mouth, where, in general, the

artisanal fishing occurs (Figure 7B). As few sport fishers acted before the introduction of the peacock bass, the answers of Juturnaíba Reservoir and São João River were grouped (Figure 7C). According to these fishers, most of the species that are also targeted in their fishing practices and that were previously found in the São João River basin are disappearing. Another consensus among the interviewed people was the relative stability of the population of the trahira throughout the time, even after the introduction of peacock bass (Figure 7).

The decrease in the abundance of other species of fishes was also mentioned, among them the leporinus and the tiete tetra. However, the fishers attribute the reduction of these species to the construction of the dam, which prevented their ascendant movements to the upstream stretch of the São João River to spawn. Only two of the artisanal fishers evaluated the impoundment of the river as positive, claiming that the water level was higher compared to the former lagoon




Figure 6. Reproductive activity of peacock bass according to the 12 artisanal and 13 sport fishers interviewed in the Juturnaíba Reservoir and downstream last stretch of the São João River, between September 28 and 08 October 2016.

and that the amount of fish increased, since, according to them, they became trapped in the reservoir.

For the artisanal fishers of the São João River, two other non-native species became more frequent in fishing, the redbreast tilapia and the african catfish *C. gariepinus* (Figure 7B). In addition to changes in the abundance of some species after introduction of the peacock bass, fishers were also asked about the occurrence of other species introduced. Among the 13 cited species, the african catfish, the cachama *Colossoma macropomum* and the cichlid oscar, *Astronotus ocellatus* and the redbreast tilapia were the most cited (Table 5). Six other non-native species were also recorded in scientific samplings (Catelani *et al.* 2020).

Two interviews in Juturnaíba presented conflicting information, although provided by artisanal fishers who have lived in the region for more than 30 years and depend economically from fishing activity. According to one of them, who did not change his methods of fishing even after the introduction of the peacock bass (basically gillnet and long line), the reduction in the abundance of some species, mainly the tetras and the leporinus is due to the predation exerted by the peacock bass. For the other one who started to offer guide services after the introduction of the peacock bass and whose incomes depend mainly on this activity, the abundance of the tetras actually did not fall. According to him, this apparent reduction of the tetras results from changes on the circadian

rhythm of this species which was formerly diurnal and switched to nocturnal habits. Concerning the reduction in the abundance of the toothless, this fisherman believes it preceded the introduction of the peacock bass and is due to the wastewater treatment for public supply, that is released into the reservoir. While the guide has his economic activity directly related to the peacock bass, the other fisherman associated the reduction of the diversity and the fish stock to the peacock bass and complained that it damages the material a lot and has low catchability, since it is able to detect the net and escape, by moving away.

Even though not all fishermen perceive impacts associated with the introduction of the peacock bass, the practices of catch and release are unanimous. According to them, this practice helps to maintain populations of peacock, the fishing activity and strengthens tourism activity. In the Juturnaíba Reservoir, it is easy to verify the presence of signs encouraging the catch and release practice.

DISCUSSION

The people interviewed demonstrated awareness and biological knowledge about the non-native peacock bass in the study area. Artisanal fishers demonstrated a more detailed and a deeper comprehension of their environment, including (i) the anthropic changes experienced since the impoundment and alteration of




Figure 7. Species cited in the interviews carried out between September 28 and 08 October 2016 with artisanal fishers before and after the introduction of the peacock bass, in the Juturnaíba Reservoir (A); with artisanal fishers before and after the introduction of the peacock bass, in the São João River (B); with sport fishers before and after the introduction of the peacock bass, in the Juturnaíba Reservoir and in the São João River (C).

Table 5. Non-native species in the Juturnaíba Reservoir and downstream last stretch of the São João River which were cited by 12 artisanal and 13 sport fishers interviewed between September 28 and October 8, 2016. The “X” on the last column, indicates species recorded in experimental catches carried out by the authors or records of scientific literature.

Popular name	Scientific name	% Citation by artisanal and sport fishers (N = 25)	Recorded in scientific catches
African catfish	<i>Clarias gariepinus</i>	40	X
Cachama	<i>Colossoma macropomum</i>	24	X
Oscar	<i>Astronotus ocellatus</i>	24	X
Redbreast tilapia	<i>Oreochromis niloticus</i>	20	X
Spotted sorubim	<i>Pseudoplatystoma</i> spp.	20	
Spotted silver dollar	<i>Metynnis lippincottianus</i>	8	X
Atipa	<i>Hoplosternum littorale</i>	8	X
Arapaima	<i>Arapaima gigas</i>	4	
Dorado	<i>Salminus brasiliensis</i>	4	
Tiete tetra	<i>Brycon</i> spp.	4	
Common carp	<i>Cyprinus carpio</i>	4	
Grass carp	<i>Ctenopharyngodon idella</i>	4	

the river course throughout channelization, (ii) the decrease of native species and (iii) the introduction of that non-native ones. In fact, the effectiveness of artisanal fishing depends on this type of knowledge

(Clauzet *et al.* 2007; Marques 1991), therefore the more precise responses of artisanal fishers reflect their cultural transmission, daily fishing practice and dependence on the resources in the study area.

In general, the biological knowledge about the peacock bass was similar between artisanal and sport fishers, with exception of the reproductive aspects of the species. Answers were more similar when regarding the consumed feeding items of the peacock bass. The uncertainty in the questions about the reproductive period may be related to the partitioned spawning, characteristic of many species of Cichlidae, including the peacock bass (Gomiero *et al.* 2009). The seasonal variation in the gonadal volume of the peacock bass is less remarkable than of those single-spawning species (i.e. migratory fishes) (Vazzoler 1996), and particularly this characteristic may hamper the perception of fishers about the spawning periods of this fish. Indeed, other studies have also shown that artisanal fishers have some difficulty in recognizing fish spawning periods elsewhere (Silvano and Begossi 2002; Begossi and Silvano 2008; Silvano *et al.* 2006; Silvano and Begossi 2010). However, some fishers detailed other reproductive characteristics of peacock bass, such as the deposition of eggs on the substrate, the parental care of both sexes and the establishment and defense of a territory where eggs are laid. In fact, these characteristics are well known to peacock basses that have been reported by other fishers in ethnobiological studies (Moura and Marques 2007). Even though their difficulty of recognizing the spawning period of the peacock bass, the interviewed fishers still provided sound information on the reproductive behavior of this non-native fish.

Fishers of both modalities stated that the peacock bass is a predatory species, which feeds on juvenile fish, including conspecifics, as well as shrimp and aquatic insects. The responses given agree to those gathered in similar studies elsewhere. In the Furnas reservoir, fishers emphasized the predatory ability of the peacock bass (Azevedo-Santos *et al.* 2010). Among the several aspects and particularities of the peacock bass biology, its voracious behavior seems the main feature used by fishers to associate the possible impacts that the species may cause in an invaded environment. Ethnobiological studies (Azevedo-Santos *et al.* 2010; Moura and Marques 2007) confirmed the decrease in population and even extirpation of some native species that became prey of the introduced peacock bass in lentic systems (i.e., lakes and reservoirs). The response of a fisherman of Furnas reservoir (Paraíba do Sul River, State of Rio de Janeiro) agrees with the fishers' perception about the peacock's food habit in this study; according to him, the peacock bass is at the top of the food chain of the reservoir, where its natural predator is absent (Azevedo-Santos *et al.* 2010).

The fishers' perception about the biology of the peacock bass was also compatible with the scientific literature. The reproductive aspects, such as parental

care and territorialism (Gomiero and Braga 2004; Winemiller 2001; Winemiller *et al.* 1997), as well as information about the food habit and the decline in prey populations (Latini and Petrere Junior 2004; Pelicice and Agostinho 2009; Pelicice *et al.* 2015) were reported for the species in the studied area. Such congruencies emphasize the quality of fishers' knowledge about the species, especially those that use it as a resource.

Besides the detected awareness of the impact of peacock bass predation over fish species, most sport fishers consider the introduction of peacock bass as positive, being favorable to the maintenance of populations in the São João River. This seemingly contradiction is partially based on the importance of the peacock bass as a fishing resource in the environments where it is introduced (Azevedo-Santos *et al.* 2010; Moura and Marques 2007). For the maintenance of the peacock bass in Juturnaíba Reservoir, most of the sport fishers encourage the practice of catch and release, even if this practice is considered an environmental crime according to Law 9605 of 1998, which prohibits the release of non-native species throughout the national territory (Brasil 1998). Artisanal fishers who maintained fishing methods and targeted species even after the introduction of the peacock bass, claimed a decrease in the abundance of their fishing resources and associated such decrease mainly to the predation by this non-native species. However, those artisanal fishers who adapted their activities including guide services for sport fishers in the fishing of the peacock bass find positive the introduction of the peacock bass, even being aware of its negative impact on the native fauna. Thus, a positive socio-economic impact was verified by the presence of peacock bass in the region, as much for its value in the commerce as for the tourism of the sport fishing.

The guides associate changes in the diversity of some fish species with other anthropogenic factors, such as the dispose of sewage and waste from potable water treatment, as well as the channeling and damming of the river. In Extremoz lagoon, Northeastern Brazil, fishers associate the loss of fish diversity mostly to two major anthropogenic changes: the construction of bridges between the Doce River and the lagoon and the introduction of the peacock bass (Rosa *et al.* 2014). Due to these changes in Juturnaíba Reservoir, economically important species have become scarce in fishers' catches, such as the tetra and the toothless. As observed in this study, fishers in Neotropical rivers generally recognize the construction of dams as the main factor for the decrease of fish species (Hallwass *et al.* 2013; Hoinghaus *et al.* 2009), especially the migratory ones, which is the case of the leporinus. Furthermore, in addition to the loss of native ichthyofauna associated with the

peacock bass or impoundment, fishers reported the occurrence of an alarming number of the non-native species. Of a total of 25 species cited by fishers in this study, 13 are non-native in the São João River basin. The most cited species was the African catfish, also known to cause many impacts where it is introduced (Weyl *et al.* 2016).

CONCLUSION

In summary, the ichthyological knowledge gathered with the sport and artisanal fishers of Juturnaíba Reservoir and São João River provide novel information related to the peacock introduction and other anthropogenic impacts. The content of the responses provided by fishers agree with the information reported in other ethnobiological and ichthyological studies. By assessing the fishers' knowledge about fish biology, anthropogenic environmental modifications, and their interaction with the peacock bass, this study highlights that the fisheries management policies should necessarily include non-native fishing resources. Even when causing environmental impacts, non-native species may become more economically important than native species, especially for the sport fishing and tourism associated with this activity.

ACKNOWLEDGEMENT

The authors are grateful to Arthur de Barros Bauer and Vagner Leonardo Macedo dos Santos for their assistance in conducting the interviews and contributions that enriched the study. We also thank Mônica Pacheco de Araujo and Arthur B. Bauer for drawing Figure 1. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. P.A. Catelani also thanks CAPES for the sandwich doctorate scholarship (CSF-PVE-S 88887.127845/2016-00) and for the financial support through the Programa de Apoio à Pós-Graduação (PROAP). F.M. Pelicice and R.A.M. Silvano thank for research grant from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, respectively grants 306869/2017-9 and 303393/2019-0).

DATA AVAILABILITY

The data used to support the findings of this research is available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

CONTRIBUTION STATEMENT

Conceived of the idea: PAC, ACP, FMP, RAMS

Data analysis: PAC, ACP, FMP, RAMS

Wrote the first draft of the manuscript: PAC

Wrote the Final versions of the manuscript: PAC, ACP, FMP, RAMS

Supervision: ACP, FMP, RAMS

REFERENCES

- Agostinho AA, Gomes LC, Pelicice FM (2007) **Ecologia e manejo de recursos pesqueiros em reservatórios do Brasil**. Maringá: Eduem 501 p.
- Agostinho CS, Pelicice FM, Marques EE, Soares AB, Almeida DA (2011) **All that goes up must come down? Absence of downstream passage through a fish ladder in a large Amazonian river**. *Hydrobiologia* 675: 1–12.
- Agostinho AA, Gomes LC, Santos NCL, Ortega JCG, Pelicice FM (2016) **Fish assemblages in Neotropical reservoirs: Colonization patterns, impacts and management**. *Fisheries Research* 173: 26–36.
- Azevedo-Santos VM, Costa-Neto EM, Lima-Stripari N (2010) **Concepção dos pescadores artesanais, acerca dos recursos pesqueiros**. *Biotemas* 23(4): 135–145.
- Begossi A, Garavello JC (1990) **Notes on the Ethnoichthyology of Fishermen from the Tocantins River (Brazil)**. *Acta Amazonica* 20: 341–351.
- Begossi A, Hanazaki N, Peroni N (2000) **Knowledge and use of biodiversity in Brazilian hot spots**. *Environment, Development and Sustainability* 2: 177–193.
- Begossi A, Silvano RAM (2008) **Ecology and ethnoecology of dusky grouper garoupa, *Epinephelus marginatus* (Lowe, 1834) along the coast of Brazil**. *Journal of Ethnobiology and Ethnomedicine* 4: 4–20.
- Begossi A, Salivonchik S, Lopes PFM, Silvano RAM (2016) **Fishers' knowledge on the coast of Brazil**. *Journal of Ethnobiology and Ethnomedicine* 12:20.
- Berkes F (1999) **Sacred Ecology - Traditional Ecological Knowledge and Resource Management**. Taylor & Francis: Philadelphia, PA.
- Biernacki P, Waldorf D (1981) **Snowball sampling: Problems and techniques of chain referral sampling**. *Sociological Methods & Research* 10(2):141–63.

- Brasil (1998) **Dispõe sobre as sanções penais e administrativas derivadas de condutas e atividades lesivas ao meio ambiente, e dá outras providências.** DOU, 31:31.
- Britton JR, Orsi ML (2012) **Non-native fish in aquaculture and sport fishing in Brazil: Economic benefits versus risks to fish diversity in the upper River Paraná Basin.** *Reviews in Fish Biology and Fisheries* 22:1–11.
- Camacho Guerreiro AI, Ladle RJ, Batista VS (2016) **Riverine fisher's knowledge of extreme climatic events in the Brazilian Amazonia.** *Journal of Ethnobiology and Ethnomedicine* 12: 50.
- Carvalho AR (2002) **Conhecimento ecológico no 'varjão' do alto rio Paraná: alterações antropogênicas expressas na linguagem dos pescadores.** *Acta Scientiarum* 24(2): 581–589.
- Catelani PA, Bauer AB, Di Dario F, Pelicice FM, Petry AC (2017) **First record of pughead deformity in *Cichla kelberi* (Teleostei: Cichlidae), an invasive species in an estuarine system in south-eastern Brazil.** *Journal of Fish Biology* 90(6): 2496–2503.
- Clauzet M, Ramirez M, Begossi A (2007) **Ethnoichthyology of artisanal fishing community from Guaibim Beach, Valença (BA), Brazil.** *Neotropical Biology and Conservation* 2(3):136–154.
- Corneta CM (2008) **Etnoictiologia de Pescadores artesanais da Vila de Picinguaba, Ubatuba, São Paulo.** Master Dissertation, Campinas: Universidade Estadual de Campinas.
- Espínola LA, Minte-Vera CV, Júlio Júnior HF (2010) **Invasibility of reservoirs in the Paraná Basin, Brazil, to *Cichla kelberi* Kullander and Ferreira, 2006.** *Biological Invasions* 12: 1837–1888.
- FAO Fisheries Department (2003) **The ecosystem approach to fisheries.** FAO Technical Guidelines for Responsible Fisheries, N^o 4, Suppl. 2. Rome, 112p.
- Fearnside PM (1999) **Social Impacts of Brazil's Tucuruí Dam.** *Environmental Manage* 24:483–495.
- Gomiero LM, Braga FMS (2004) **Reproduction of species of the genus *Cichla* in a reservoir in southeastern Brazil.** *Brazilian Journal of Biology* 64(3B): 613–624.
- Gomiero LM, Villares Junior GA, Naous F (2009) **Reproduction of *Cichla kelberi* Kullander and Ferreira 2006 introduced into an artificial lake in southeastern Brazil.** *Brazilian Journal of Biology* 69(1): 175–183.
- Hallwass G, Lopes PF, Juras AA, Silvano RAM (2013) **Fishers' knowledge identifies environmental changes and fish abundance trends in impounded tropical rivers.** *Ecological Applications* 23(2): 392–407.
- Hoeinghaus DJ, Agostinho AA, Gomes LC, Pelicice FM, Okada EK, Latini JD, Kashiwaqui EAL, Wine-miller KO (2009) **Effects of river impoundment on ecosystem services of large tropical rivers: embodied energy and market value of artisanal fisheries.** *Conservation Biology* 23: 1222–1231.
- Huntington HP, Begossi A, Gearheard SF, Kersey B, Loring PA, Mustonen T, Paudel PK, Silvano RAM, Vave R (2017) **How small communities respond to environmental change: patterns from tropical to polar ecosystems.** *Ecology Society* 22(3):9.
- Johnson PTJ, Olden JD, Vander Zanden MJ (2008) **Dam invaders: Impoundments facilitate biological invasions into freshwaters.** *Frontiers in Ecology and the Environment* 6(7): 357–363.
- Latini AO, Petrere Junior M (2004) **Reduction of a native fish fauna by alien species: an example from Brazilian freshwater tropical lakes.** *Fisheries Management and Ecology, Oxford* 11(2): 71–79.
- Magalhães HF, Costa-Neto EM, Schiavetti A (2011) **Saberes pesqueiros relacionados à coleta de siris e caranguejos (Decapoda: Brachyura) no município de Conde, Estado da Bahia.** *Biota Neotropical* 11(2): 045–054.
- Marques JGW (1991) **Aspectos ecológicos na ictiologia dos pescadores do Complexo Estuarino-Lagunar Mundaú-Manguaba.** Tese de Doutorado, Campinas: Universidade Estadual de Campinas.
- Morril WT (1967) **Ethnoichthyology of the Chacha.** *Ethnology* 6: 405–417.
- Moura FBP, Marques JGW (2007) **Conhecimento de pescadores tradicionais sobre a dinâmica espaço-temporal de recursos naturais na Chapada Diamantina, Bahia.** *Biota Neotropical* 7(3): 119–126.
- Moura FBP, Marques JGW, Nogueira EMS (2008) **“Peixe sabido, que enxerga de longe”: conhecimento ictiológico tradicional na Chapada Diamantina, Bahia.** *Biotemas* 21(3): 115–123.
- Mourão JS, Nordi N (2002) **Comparações entre as taxonomias folk e científica para peixes do**

estuário do Rio Mamanguape, Paraíba, Brasil. *Interciencia* 27 (12): 664–668.

Pelicice FM, Agostinho AA (2009) **Fish fauna destruction after the introduction of a non-native predator (*Cichla kelberi*) in a Neotropical reservoir.** *Biological Invasions* 11:1789–1801.

Pelicice FM, Agostinho AA, Latini JD (2015) **Fish fauna disassembly after the introduction of a voracious predator: main drivers and the role of the invader's demography.** *Hydrobiologia* 746: 271–283.

Petrere M (1996) **Fisheries in large tropical reservoirs in South America.** *Lakes & Reservoirs Research & Management* 2:111–133.

Pinheiro L (2004) **Da ictiologia ao etnoconhecimento: saberes populares, percepção ambiental e senso de conservação em comunidade ribeirinha do rio Pirai, Joinville, Estado de Santa Catarina.** *Acta Scientiarum Biological Sciences* 26(3): 325–334.

Pinto MF, Mourão JS, Nobrega Alves RR (2017) **Animal source food consumed in two fishing communities on the northeast coast of Brazil.** *Environment, Development and Sustainability* 19(2): 679–692.

Porcher LCF, Poester G, Lopes M, Schonhofen P, Silvano RAM (2010) **Percepção dos moradores sobre os impactos ambientais e as mudanças na pesca em uma lagoa costeira do litoral sul do Brasil.** São Paulo. *Boletim Instituto de Pesca* 36(1): 61–72.

Ramires M, Clauzet M, Barrella W, Rotundo MM, Silvano RAM, Begossi A (2010) **Fishers' knowledge about fish trophic interactions in the southeastern Brazilian coast.** *Journal of Ethnobiology and Ethnomedicine* 11:19.

Rosa R, Carvalho AR, Angelini R (2014) **Integrating fishermen knowledge and scientific analysis to assess changes in fish diversity and food web structure.** *Ocean Coastal Management* 102: 258–268.

Silvano RAM, Begossi A (2002) **Ethnoichthyology and fish conservation in the Piracicaba River**

(Brazil). *Journal of Ethnobiology* 22(2): 285–306.

Silvano RAM, Begossi A (2010) **What can be learned from fishers? An integrated survey of ecological knowledge and bluefish (*Pomatomus saltatrix*) biology on the Brazilian coast.** *Hydrobiologia* 637: 3–18.

Silvano RAM, Maccord PFL, Lima RV, Begossi A (2006) **When does this spawn? Fishermen's local knowledge of migration and reproduction of Brazilian coastal fishes.** *Environmental Biology of Fishes* 76: 371–86.

Silvano RAM, Silva AL, Ceroni M, Begossi A (2008) **Contributions of ethnobiology to the conservation of tropical rivers and streams.** *Aquatic Conservation* 18: 241–260.

Silvano RAM, Valbo-Jørgensen J (2008) **Beyond fishermen's tales: contributions of fishers' local ecological knowledge to fish ecology and fisheries management.** *Environment, Development and Sustainability* 10:657–675.

Vazzoler AEAM (1996) **Biologia da reprodução de peixes teleósteos: teoria e prática.** Maringá, EDUEM; São Paulo, SBI, 169p.

Weyl OLF, Daga VS, Ellender BR, Vitulle JRS (2016) **A review of *Clarias gariepinus* invasions in Brazil and South Africa.** *Journal of Fish Biology* 89(1): 386–402.

Winemiller KO, Taphorn DC, Barbarino-Duque A (1997) **The ecology of *Cichla* (Cichlidae) in two blackwater rivers of southern Venezuela.** *Copeia* 690–696.

Winemiller KO (2001) **Ecology of peacock cichlids (*Cichla* spp.) in Venezuela.** *Journal of Aquaculture and Aquatic Sciences* 9: 93–112.

Zaret TM, Paine RT (1973) **Species introduction in a tropical lake.** *Science* 182: 449–455.

Received: 26 September 2019

Accepted: 08 September 2020

Available: 25 September 2020

Additional Files

Add File 1. Popular and scientific names of the species cited along the manuscript.

Popular name (Portuguese)	Popular name (English)	Scientific name
Bagre-africano	African catfish	<i>Clarias gariepinus</i> (Burchell 1822)
Pirarucu	Arapaima	<i>Arapaima gigas</i> (Schinz, 1822)
Cascudo-viola	Armored catfishes	<i>Loricariichthys castaneus</i> (Castelnau 1855)
Camboatá	Atipa	<i>Hoplosternum littorale</i> (Hancock, 1828)
Tambaqui	Cachama	<i>Colossoma macropomum</i> (Cuvier, 1816)
Palombeta	Castin leatherjacket	<i>Oligoplites</i> spp.
Carpa	Common carp	<i>Cyprinus carpio</i> Linnaeus, 1758
Dourado	Dorado	<i>Salminus brasiliensis</i> (Cuvier, 1816)
Carpa-capim	Grass carp	<i>Ctenopharyngodon idella</i> (Valenciennes, 1844)
Piau	Leporinus	<i>Leporinus copelandii</i> Steindachner, 1875
Tainha	Mullet	<i>Mugil curema</i> spp.
Apaiari	Oscar	<i>Astronotus ocellatus</i>
Tucunaré	Peacock bass	<i>Cichla kelberi</i> Kullander & Ferreira 2006
Acará	Pearl cichlid	<i>Geophagus brasiliensis</i> (Quoy & Gaimard 1824)
Jacundá	Pike cichlid	<i>Crenicichla lacustris</i> (Castelnau 1855)
Tilápia	Redbreast tilapia	<i>Oreochromis niloticus</i> (Linnaeus, 1758)
Robalo	Sea bass	<i>Centropomus parallelus</i> Poey 1860
Cumbaca	Singing catfish	<i>Trachelyopterus striatulus</i> (Steindachner, 1877)
Pacu	Spotted silver dollar	<i>Metynnis lippincottianus</i> (Cope, 1870)
Pintado	Spotted sorubim	<i>Pseudoplatystoma</i> spp.
Piaba	Tetra	<i>Astyanax</i> spp.
Piabanha	Tiete tetra	<i>Brycon</i> spp.
Sairú	Toothless characins	<i>Cyphocharax gilbert</i> (Quoy & Gaimard, 1824)
Bagre-branco	White sea catfish	<i>Genidens barbatus</i> (Lacepède 1803)
Traíra	Trahira	<i>Hoplias malabaricus</i> (Bloch 1794)