



## Addressing Taxonomic and Geographic Biases in ethnozoological studies on animal use in Brazil

Suellen da Silva Santos<sup>1</sup>, Hyago Kesley de Lucena Soares<sup>2</sup>, Henrique Fernandes Magalhães<sup>3</sup>, Eraldo Medeiros Costa Neto<sup>4</sup>, Fabio de Oliveira Roque<sup>5,6</sup>, Davidson Gomes Nogueira<sup>5</sup> and Reinaldo Farias Paiva de Lucena<sup>5</sup>

### ABSTRACT

This study aims to assess the current state of ethnozoological research in Brazil. Systematic review was conducted using the Web of Science and Scopus databases, encompassing all publications available until December 2022. The selected articles, written in various languages, consist of data obtained directly by the researcher, offering taxonomic descriptions of species and their respective uses. 85 articles were selected to construct a distribution map of ethnozoological studies, categorizing them based on their use. The Spearman's correlation test and the Kruskal-Wallis test, were applied to identify patterns within the data on use, number of studies and zoological class. Spearman's correlation indicated a positive relationship between the sum of uses and the number of studies conducted. The Kruskal-Wallis test revealed variations in the average number of studies and total uses across different zoological classes. The thematic classification of articles included general ethnozoology, specialized ethnozoology, general medicinal use, specific medicinal use, and hunting. Results showed a consistent increase in publications, with the Northeast region and the Caatinga biome featuring prominently. The reviewed papers encompassed 829 species, demonstrating greater representation in the food and medicinal categories. Ichthyofauna and avifauna, exhibited the highest species number. Our study highlights the utilitarian versatility of herpetofauna and mammal fauna. The classes Asterozoa, Bivalvia, Chilopoda, Echinozoa, and Hydrozoa, were identified as having below-average representation in terms of studies and use associations. In conclusion, the revision advocates for investigations with a collaborative bias among research groups from different biomes, to aid in reducing centralized studies in specific geographical areas.

**Keywords:** Traditional knowledge of fauna; Use of wild animals; Scientometrics.

1 Program in Natural Resources. Research Group on Multidisciplinary and Socioecological Studies, Institute of Biosciences, Federal University of Mato Grosso do Sul, Campo Grande, Brazil.

2 Program in Biological Sciences (Zoology), Federal University of Paraíba, João Pessoa, Paraíba, Brazil.

3 Laboratory of Applied Ecology, Department of Botany, Center for Biosciences, Federal University of Pernambuco, Pernambuco, Brazil.

4 Program in Ecology and Evolution, State University of Feira de Santana, Feira de Santana, Bahia, Brazil.

5 Research Group on Multidisciplinary and Socioecological Studies, Institute of Biosciences, Federal University of Mato Grosso do Sul, Campo Grande, Brazil.

6 Brazilian Knowledge Center on Biodiversity, Belo Horizonte, MG, Brazil.

\* Corresponding author ✉. E-mail address: SSS ([suellen\\_santos@ufms.br](mailto:suellen_santos@ufms.br)), HFM ([henrique.fmagalhaes@ufpe.br](mailto:henrique.fmagalhaes@ufpe.br)), EMCN ([eraldomcosta@gmail.com](mailto:eraldomcosta@gmail.com)), FOR ([reinaldo.lucena@ufms.br](mailto:reinaldo.lucena@ufms.br))

## SIGNIFICANCE STATEMENT

To evaluate the current state of ethnozoological research in Brazil, we conducted a systematic review encompassing all studies available in the Web of Science and Scopus databases up to December 2022. A total of 85 articles, written in various languages, were selected, each containing primary data on taxonomic descriptions of diverse animal species and their respective traditional uses. From these selections, we created a distribution map of ethnozoological studies categorized by type of use. Additionally, we applied Spearman and Kruskal-Wallis correlation tests to identify patterns within the data regarding usage, number of studies, and zoological classes. Our findings reveal that: 1) there is a positive relationship between the total number of uses and the number of studies conducted; 2) variations exist in the average number of studies and total uses across different zoological classes; 3) there is a steady increase in the number of ethnozoological studies in Brazil, particularly in the North-east region and the Caatinga biome; 4) the food and medicinal categories were the most frequently considered in the review, with significant emphasis on the utilitarian versatility of herpetofauna and mastozoofauna.

## INTRODUCTION

Biodiversity has undergone scrutiny from various perspectives. The scientific emphasis on its utilitarian aspects, along with considerations of local ecological knowledge (CEL) and local zoological knowledge (LZK), has facilitated an understanding of the connections forged between human populations and faunal resources. Studies have documented and disclosed substantial data on management practices, bioprospecting, the discovery of new species, insights into the impact of biological and ecological phenomena, the development of hypotheses, and the application of methodological tests (Alves and Souto 2015; Braga-Pereira *et al.* 2022; Costa-Neto *et al.* 2013; Gama 2014; Morado *et al.* 2021; Rodríguez *et al.* 2021; Santos *et al.* 2019c; Serrão *et al.* 2019; Pozzi and Ladio 2023).

Ethnozoological studies have contributed to the understanding of the identification and classification of animals by humans, taking into account cultural, environmental, and historical contexts (Alves and Souto 2015). These studies reveal that numerous species within the Brazilian fauna serve the diverse needs of populations. They are utilized as source of protein, zotherapy resources, in the manufacturing of artisanal products, in mystical and religious rituals, and pets (Alencar *et al.* 2012; Santos *et al.* 2019a, b; Santos *et al.* 2020; Santos *et al.* 2021; Soares *et al.* 2018; Tito and Giraldin 2021). Certain species are also considered significant for people due to their role as predictors of climatic events, conveying either good or bad news, which, in turn, leads to some of them being slaughtered over human aversion (Almeida *et al.* 2023; Bonfim *et al.* 2021; Los *et al.* 2013; Margalida *et al.* 2014; Santos *et al.* 2023; Silva *et al.* 2017).

Despite the growing number of studies and recognition of the area's importance for conservation, knowledge remains fragmented and scattered throughout the literature. There is a need to summarize information regarding the current state of fauna species of utilitarian importance for human, particularly in cultural and biodiversity rich countries, such as Brazil. Although

there are reviews regarding ethnozoological knowledge, there is a need to compile data that indicates information such as zoological taxa in greater evidence, areas that require expanding research, and understanding established collaboration networks in studies conducted in Brazil by national and international educational institutions.

When analyzing specific data, systematic reviews can offer syntheses of the knowledge state in a scientific field and future research priorities can be identified, addressing issues that may remain unanswered through individual studies alone (Page *et al.* 2021). Considering Brazil's pivotal role as a trailblazer in ethnozoological research, with a rich legacy of academic contributions (Oliveira and Lopes 2021), we carried out a scientometric survey to systematically explore the taxa that hold representative significance within human communities residing in Brazilian territory. In addition to assessing the correlation between the total use of animals and developed studies, and the average number of studies per zoological class, as well as identifying national and international collaboration networks that have contributed to this science.

By undertaking this comprehensive examination, we aim to not only contribute to the existing body of knowledge but also to guide and inspire future inquiries, ensuring a nuanced understanding of the intricate relationships between humans and the diverse fauna that share their habitat. We focused on ethnozoological studies and addressed several key questions: 1) What is the current landscape of ethnozoological research in Brazilian territory?; 2) Which zoological groups are most frequently studied?; 3) What utilitarian purposes are predominantly explored?; 4) What are the primary thematic approaches evident in the analyzed papers, and what potential gaps exist that warrant further investigation?; and 5) How have institutional scientific collaboration networks been acting in the development of these studies?

## MATERIAL AND METHODS

### Search strategies

The research entails a systematic review conducted in accordance with the PRISMA guidelines (2020) (2020) (Page *et al.* 2021), utilizing the Web of Science database to identify relevant articles (<https://www-webofscience.ez15.periodicos.capes.gov.br/wos/woscc/basic-search>) and Scopus (<https://www-scopus.ez15.periodicos.capes.gov.br/search/form.uri?display=basic#basic>).

Our search strategy involved employing the following keywords: Ethnozoology, Ethnomastozoology, Ethno-ornithology, Ethnoherpetology, Ethnoentomology, Ethnoichthyology, Ethnocarcinology, Ethnomalacology, Zootherapy, and Hunting.

### Selection of studies

In the initial stage of document identification, we conducted searches using a combination of the aforementioned keywords, pairing them with the term “Brazil”. In the case of the term “Hunting”, we added the keyword “Ethnozoology” to this combination. This addition was made to account for articles found outside the scope of the research topic, such as “Hunting AND Brazil AND Ethnozoology”.

During the tracking stage, we performed a preliminary assessment of located articles by reviewing titles, abstracts, and keywords. Articles that did not reference the keywords used in the search within these fields were disregarded. This exclusion criterion applied particularly to articles structured around secondary data, scientific notes, and duplicates.

In the eligibility stage, we thoroughly examined the full articles to identify those containing the necessary information for the analyses included in this review. Consequently, we incorporated articles obtained from the search platforms until December 2022. These articles, written in any language, consisted of primary data and involved the taxonomic identification of Brazilian fauna species along with analyses of their utilization.

We extracted the following data from the studies: the publication year of the article, the biome, state, and municipality where each research was conducted, the described species and zoological group, and the utilitarian purpose associated with each animal.

### Data Analysis

We generated the distribution map of ethnozoological studies using the geographic coordinates provided in the analyzed articles through the QGIS program. In cases where this geographic data was absent, we

conducted a search using Google Maps, utilizing information related to the municipality(ies) specified in each search (Figure 3).

Species nomenclature was cross-verified based on the information presented in the articles, and classifications were revised in accordance with the Brazilian Biodiversity Information System. This was particularly crucial in instances where the records diverged due to changes in species nomenclature (SIBBr 2020).

The recorded uses in the analyzed studies were categorized into the following groups: food, medicinal, artisanal (encompassing decorative items and household utensils), pets, magical-religious (pertaining to species used in mystical practices), veterinary (involving the use of animals or their by-products for medicinal treatment of other wild or domestic species), conflict (involving animals considered pests causing damage to plantations, domestic species, or inducing aversion), and bioindication (pertaining to species indicative of meteorological variations and omens, possibly associated with positive news). (Barbosa and Aguiar *et al.* 2015; Abrantes *et al.* 2011; Montenegro *et al.* 2014).

To assess the correlation between the sum of uses and the number of studies conducted, the Spearman’s correlation test was employed. In addition, the Kruskal-Wallis test was used to determine whether there was a difference between the average number of studies and the total use (sum of all uses by species) by taxonomic zoological class. These analyses were conducted using R Software (R Core Team 2021).

Lastly, to comprehend the main thematic approaches within the articles, we followed the classification proposed by Ritter *et al.* (2015), adapting it to ethnozoological studies. This classification includes: General Ethnozoology (encompassing studies addressing the general use of animal resources), Specialized Ethnozoology (covering works focused on the use of fauna for specific purposes such as food or crafts, excluding medicinal purposes, or aimed at a particular zoological group or species), General Medicinal Use (encompassing studies on the general use of medicinal animals), Specific Medicinal Use (including specific studies on the use of medicinal animals within a zoological group or species), and Hunting (encompassing studies on the use of fauna identified using this keyword, not included in other searches). It is worth noting that medicinal categorization is treated separately due to the existence of numerous articles specifically dedicated to this topic.

### Bibliometric Analysis

To perform the bibliometric analysis, we used only the papers available on the Web of Science Core Collection and then we used the VOSviewer 1.6.18 software. An analysis of terms frequency and co-occurrence al-

lowed mapping relationships between all the papers and identify main addressed topics. This analysis was conducted in the titles, keywords and abstracts. Analysis of authorship affiliations permitted to map the most active countries/territories in terms of publications. The authorship affiliations analysis used the list of affiliations of the authors and co-authors (Van Eck and Waltman 2010).

## RESULTS

### Screening of Articles and Profile of Ethnozoological Studies

A total of 513 articles were initially identified across the aforementioned search platforms. However, adhering to the steps outlined by PRISMA, our study narrowed down to the analysis of 85 articles (Figure 1).

The documents retained in the review encompass at least one taxon identified at the genus level, along with its corresponding utilitarian purpose. For instance, consider the study by Barbosa *et al.* (2022): the authors report on 156 species distributed among mammal fauna (31), avifauna (93), and herpetofauna groups, which include reptiles (27) and amphibians (05), with associated usage citations. However, the document selectively highlights certain scientific identifications, and these animals are included in our database.

Articles that were excluded from the study did not meet the inclusion criteria for various reasons: 1) originating from another country; 2) involving secondary data; 3) lacking mention of animals from the Brazilian fauna; 4) lacking the scientific identification of the species; 5) failing to specify the utilitarian purpose associated with the species; 6) dealing with studies focused on experimental investigations, phytochemical evaluations, and specimen capture techniques.

The publications spanned the years 1999 to 2022, with significant representation observed between 2011 and 2020 (see Figure 2).

Articles are predominantly developed in the Northeast region (76 papers), with a limited number focused on the Central-West and South regions, each comprising only one study (see Figure 3).

Among the Brazilian biomes, the Caatinga (prevalent in the Northeast region) attracts the most significant research investment with 52 papers, followed by the Atlantic Forest (21 papers), the Amazon Forest (8 papers), and the Cerrado (2 papers) (see Figure 3). Some articles were developed in ecotone areas, and others did not specify the biome in which they were conducted. It is important to highlight that, in the case of the Central-West region, the reported research

was conducted in collaboration with academics residing in 14 municipalities within the Middle Araguaia region. The extension of study participant allocations resulted in an increased heat spot displayed on the map for the region, which does not accurately reflect the actual number of studies conducted in the area.

Upon evaluating the attributes assigned to species, we observed a greater representation of vertebrates. A total of 829 species were identified, comprising 270 birds, 229 fish, 102 reptiles, 86 mammals, 58 insects, 27 mollusks, 25 crustaceans, 13 amphibians, 9 echinoderms, 7 arachnids, 1 annelid, 1 cnidarian, and 1 kilopod. Although avifauna and ichthyofauna present the highest number of described species, in terms of utilitarian versatility, herpetofauna and mammal fauna stand out. Species such as *Salvator merianae*, *Boa constrictor*, *Crotalus durissus*, *Cayman latirostris*, *Euphractus sexcinctus*, *Cerdocyon thous*, and *Tamandua tetradactyla* are associated with seven utility categories (excluding bioindication). The highest number of animals is associated with food use, followed of categories reported in Table 1.

Concerning the applied statistics, Spearman's correlation revealed a positive result between the sum of uses and the number of studies, yielding  $S = 4598709$ ,  $p\text{-value} < 2.2e-16$ , and  $\rho = 0.9515689$  (Figure 4).


The Kruskal-Wallis test for the average number of studies per zoological class resulted in a  $p\text{-value}$  of  $8.691e-14$  (Figure 5). This outcome demonstrates that the classes Asterozoa (Echinodermata) (4), Bivalvia (Mollusca) (6), Chilopoda (Arthropoda) (9), Echinozoa (Echinodermata) (12), and Hydrozoa (Cnidaria) (14) (Figure 6) are below average.

For the Total uses by zoological class, a  $p\text{-value}$  of  $4.785e-15$  was obtained. This result indicates that the classes Asterozoa (4), Chilopoda (9), Echinozoa (12), and Hydrozoa (14) are below average (Figure 6).


The predominant thematic approach in the analyzed articles was Specific Ethnozoology (52 papers), followed by General Medicinal Use (17 papers), Hunting (11 papers), Specific Medicinal Use (4 papers), and General Ethnozoology (1 paper) (Figure 1).

### Bibliometric Analysis

Out of the 76 articles predominantly developed in the Northeast region 60 were available in the Core Collection of the Web of Science. The analysis of terms showed the most recurrent terms and the interconnection strength between them over time (Figure 7) and by cluster (Figure 8). The results present the 91 most recurrent terms grouped among 5 clusters, of which the most recurrent terms were "bird", "pet" and "medicine" (51, 45 and 34 times respectively). The clusters were grouped in topics related to health benefits (red cluster), Ecology and Conservation (green cluster), Lo-



**Figure 1.** Number of studies selected for the systematic review, following the guidelines proposed by the PRISMA protocol (Page et al. 2021).




**Figure 2.** Number of ethnozoological studies available in the Web of Science and Scopus databases per publication year from 1999 to 2022.

cal Knowledge and Techniques (purple cluster), Human Consumption (yellow cluster), and Human Animal Conflicts (blue cluster).

The results of the affiliation analysis show the co-authorship by countries of the published papers (Figure 9). The results show the time in terms of publications and their relationship in publications. All the papers were affiliated to Brazil which is the center of the publications and only a few other countries show

up in the research of which the more developed countries present older partnership with Brazil while developing countries present newer partnership (later than 2020).

As for the results of the affiliation analysis of co-authorship by institutions, the institutions from Northeast Brazil were most prominent and have been publishing the longest in this topic (Figure 10). These institutions with more papers are in general older in-



**Figure 3.** Geographic distribution of ethnozoological research conducted in Brazil available in the Web of Science and Scopus databases from 1999 to 2022.

stitutions while institutions that have only recently started publishing newer institutions. The pattern of co-authorship also shows that the institutions that tend to publish together more and cluster together do so by regions, for instance institutions from the Southeast Brazil are mostly in the same cluster, the ones from the Northeast in another, and also the ones outside Brazil in a separate cluster (Figure 11).

## DISCUSSION

### Geographic bias

Our review indicates a trend towards increased Brazilian publications in the field over the last decades especially in the Caatinga and Atlantic Forest biomes.

Our data align with the findings of the review conducted by Alves and Souto (2011), who, in their analysis of scientific materials on ethnozoology in Brazil, identified exponential growth in research from the early 21st century, particularly between 2001 and 2011, and with the Corrêa *et al.* (2022), who, when analyzing publications between 1967 and 2017, observed peaks in publications in 2006, 2012, and 2017, but noted a decline in 2017. The authors found that 38% of the studies were conducted in the Caatinga ( $n = 61$ ), 23% in the Atlantic Forest ( $n = 38$ ), 19% in the Cerrado ( $n = 31$ ), 17% in the Amazon Rainforest ( $n = 27$ ), 2% in the Pampa ( $n = 3$ ), and 1% in the Pantanal ( $n = 2$ ). Despite the overall growth in ethnozoological studies in Brazil, this scenario highlights that there are still regions that demand greater investment.

**Table 1.** Taxonomic classes and their uses categories.


Taxonomic classes	Categories							
	Food	Medicinal	Breeding	Magical-religious	Conflict	Craft purposes	Veterinary	Bioindication
Actinopteri	142	45	-	18	4	18	3	-
Asteroidea	-	6	-	4	-	1	-	-
Arachnida	-	4	-	-	2	-	2	-
Aves	161	77	109	49	32	-	5	15
Bivalvia	-	3	-	3	-	-	-	-
Camarondonta	-	1	-	-	-	-	-	-
Cephalopoda	-	1	-	-	-	-	-	-
Chilopoda	-	1	-	-	-	-	-	-
Chondrichthyes	18	17	-	1	6	-	-	-
Clitellata	-	2	-	-	-	-	-	-
Echinoidea	-	2	-	1	-	-	-	-
Gastropoda	1	4	-	17	-	-	-	-
Hydrozoa	-	1	-	1	-	-	-	-
Insecta	6	41	1	2	12	2	7	1
Lissamphibia	3	11	-	3	-	-	1	-
Malacostraca	5	15	-	5	-	10	-	-
Mammalia	50	64	13	29	34	14	12	-
Reptilia	29	58	55	25	41	21	11	-
Total species	415	353	178	158	131	66	41	16
Total (%)	50	42.5	21.4	19	15.7	7.9	4.9	1.9

The prominence of the Northeast region and institutions from the region can be attributed to consolidated research groups in ethnozoology affiliated with academic institutions in this area (Oliveira and Lopes 2021) which facilitates the expansion of research development as newer institutions start to take part during the last few years and hubs of development, even though it is mostly centralized around older institutions. Another contributing factor to the research expansion in the region is the persistent socioeconomic and environmental challenges, despite ongoing development initiatives and government programs providing tax and economic benefits. The northeastern states rank lower in the national Human Development Index (HDI), have the lowest life expectancy in Brazil, lack environmental sanitation in approximately 55% of residences, and face challenges such as irregular rainfall impacting livestock and agricultural practices (ArboControl 2023).


These factors can influence the strengthening of

connections between social groups in this territorial extension and the local fauna, thereby directing the focus towards conducting ethnozoological research. Furthermore, certain studies span multiple locations, exemplified by Barboza *et al.* (2016), who investigated mammal hunting with rural and urban communities in the Caatinga biome across 10 municipalities in four states. Similarly, Barbosa *et al.* (2022) explored hunting activity in four protected areas of the Atlantic Forest situated in four municipalities. The Southeast region accounted for three articles, while the North region for nine. However, it is noteworthy that several studies were conducted in more than one region.

There necessity of expanding ethnozoological research in biomes with a low concentration of studies, particularly in regions like the Cerrado, which had the fewest studies, and Pantanal, which lacked articles. The Brazilian Cerrado, classified as a global biodiversity hotspot by the Ministry of the Environment (MMA 2023a), emerges as a priority area for stud-



**Figure 4.** Scatterplot of the correlation between the general sum and the number of ethnozoological studies carried out in Brazil available in the Web of Science and Scopus databases from 1999 to 2022.




**Figure 5.** Boxplot of the sum of zoological classes and the number of ethnozoological studies carried out in Brazil available in the Web of Science and Scopus databases from 1999 to 2022. Caption: 1. Actinopteri; 2. Lissamphibia; 3. Arachnida; 4. Asteroidea; 5. Aves; 6. Bivalvia; 7. Camarondonta; 8. Cephalopoda 9. Chilopoda; 10. Chondrichthyes; 11. Clitellata; 12. Echinoidea; 13. Gastropoda; 14. Hydrozoa; 15. Insecta; 16. Malacostraca; 17. Mammalia; 18. Reptilia.

ies aiming to comprehend the socio-environmental and socioeconomic dynamics at play. The Pantanal, recognized as the world's largest continuous floodplain with a rich presence of traditional and native communities, has been impacted by human activities, especially agricultural practices (MMA 2023b), necessitating increased investment in ethnozoological research.


Another ecoregion in the Central-West of Brazil requiring expanded study is the Chaco. Affected by soybean production, the area has experienced health and social conflicts among communities residing near

plantations (Yousefi *et al.* 2018). Lacking official government recognition as a biome, the Chaco is often overlooked in public policies (Campos 2023). The consolidation of data in these locations may enhance their visibility by identifying weaknesses and potentialities related to social, environmental, and economic dimensions, which, in turn, can provide directions for necessary conservation and restoration strategies.





**Figure 6.** Boxplot of the sum of zoological classes and the number of ethnozoological studies carried out in Brazil available in the Web of Science and Scopus databases from 1999 to 2022. Caption: 1. Actinopteri; 2. Lissamphibia; 3. Arachnida; 4. Asteroidea; 5. Aves; 6. Bivalvia; 7. Camarondonta; 8. Cephalopoda 9. Chilopoda; 10. Chondrichthyes; 11. Clitellata; 12. Echinoidea; 13. Gastropoda; 14. Hydrozoa; 15. Insecta; 16. Malacostraca; 17. Mammalia; 18. Reptilia.



**Figure 7.** The image shows the 91 (60% most relevant terms that occurred at least 5 times) out of the 1896 recurrent terms across 60 papers acquired from Web of Science. The most occurrent terms were “bird”, “pet” and “medicine” (51, 45 and 34 times respectively). Colors represent the time period the term was most recurrent according to label and circle sizes the relative occurrences. Lines represent cooccurrences between terms and lines’ thicknesses the relative times terms cooccurred.


## Animal use bias

Research indicates that factors influencing people’s habits include sociocultural and economic considerations, taboos, utilitarian preferences, and/or ecological and biological aspects of the species. This results in the individual or multiple use of animals, contributing valuable data to their conservation efforts (Alves *et al.* 2021; Barbosa *et al.* 2022; Barboza *et al.* 2016; Landim *et al.* 2023; Marques *et al.* 2022; Santos *et al.*


2021; Silva *et al.* 2023; Soares *et al.* 2018; Soares *et al.* 2020).

In the analyzed articles, various purposes are attributed to the identified species, including food, medicinal use, pet keeping, magical-religious significance, conflict resolution, artisanal, veterinary applications, and bioindication, with the first two being the most recurrent.

For food purposes individuals are prepared and consumed based on individual preferences for prepa-



**Figure 8.** The image shows the 91 (60% most relevant terms that occurred at least 5 times) out of the 1896 recurrent terms across 60 papers acquired from Web of Science. The most occurrent terms were “bird”, “pet” and “medicine” (51, 45 and 34 times respectively). Colors represent clusters of terms that most occurred together. Lines represent cocurrences between terms and lines’ thicknesses the relative times terms cocurred.




**Figure 9.** The image shows the six countries across the 60 papers acquired from Web of Science. The country affiliated to the most papers was Brazil (with 60 papers) followed by Argentina, England, Portugal, and USA (with two papers each) and Indonesia and Mexico (with one paper each). Colors represent the time period the country published the most according to label and circle sizes the relative number of papers affiliated to it. Lines represent co-authorship between countries and lines’ thicknesses the relative times countries co-authored papers.


ration of people (e.g. Alves *et al.* 2012a; Barros and Azevedo 2014; Bonifácio *et al.* 2016; Magalhães *et al.* 2006; Pinto *et al.* 2013; Rocha-Mendes *et al.* 2005; Santos *et al.* 2022; Santos and Alves 2016). Certain social groups, such as fishermen, despite having access to protein sources like beef, often turn to fish as their main protein source, contributing to their food security (e.g. Pinto *et al.* 2017). Future assessments should list species genuinely useful for subsistence, in conjunction with other analyses such as studies on the biology and ecology of species, to propose strategies for the sustainable use of these resources.

Several species are used in medicinal practices,

with 584 species described for Latin America (Alves and Alves 2011) and for Brazil alone 354 animal species (Alves *et al.* 2013). Many of these medicinal practices involve the use of live animals, parts of their bodies (e.g., skin and feathers), by-products resulting from their metabolic actions (body secretions such as fat), and materials constructed by them (nests or cocoons). These uses are linked to the treatment of various pathologies, for instance respiratory (e.g. Alves *et al.* 2012a; Alves and Pereira Filho 2007; Bezerra *et al.* 2017; Costa-Neto *et al.* 2002; Modro *et al.* 2009; Seger *et al.* 2019). It is important to specify the type of use and condition to be treated during



**Figure 10.** The image shows the 43 interconnected institutions out of the 57 across the 60 papers acquired from Web of Science, the other 14 institution did not have any connections with other institutions. The institutions affiliated to the most papers were Universidade Estadual Paraíba, Universidade Federal Paraíba and Universidade Federal rural de Pernambuco (with 39, 29 and 9 respectively). Colors represent the time period the institution published the most according to label and circle sizes the relative number of papers affiliated to it. Lines represent co-authorship between institutions and lines' thicknesses the relative times institutions co-authored papers. The 14 institutions that do not appear in the map do not have connections with any other institution.



**Figure 11.** The image shows the 43 interconnected institutions out of the 57 across the 60 papers acquired from Web of Science, the other 14 institution did not have any connections with other institutions. The institutions affiliated to the most papers were Universidade Estadual Paraíba, Universidade Federal Paraíba and Universidade Federal rural de Pernambuco (with 39, 29 and 9 respectively). Colors represent clusters of institutions that most co-authored papers together and circle sizes the relative number of papers affiliated to the institution. Lines represent co-authorship between institutions and lines' thicknesses the relative times institutions co-authored papers. The 14 institutions that do not appear in the map do not have connections with any other institution.

data collection, as it aids in the development of conservation actions (Alves and Alves 2011; Moura and Marques 2008).

Regarding the animals as pets, these studies contain crucial information, including data on commercial networks and species management for commercial purposes (e.g. Alves et al. 2012a; Alves et al.

2016a; Alves et al. 2019; Bezerra et al. 2017, 2020; Fernandes-Ferreira et al. 2012; Licarião et al. 2013; Oliveira et al. 2018; Oliveira et al. 2020; Soares et al. 2020; Sodr e-Andre et al. 2022; Souto et al. 2017; Teixeira et al. 2014). “Pet” is one of the most mentioned term in the term map, right along with birds and in the same category as “trades”, which is also a topic

that is being used more often in the last decade. This shows the importance of surveillance of animal trafficking since it has negative impacts on the environment (Sollund 2013; Duarte et al. 2021). Conscientization of the impacts of having wild animals as pets could have impacts in the demands for such animals, which in turn would lead to less poaching and trafficking, decreasing the offers (Sollund 2013).

To fulfill the purpose magical-religious, live animals or products of animal origin are used and/or sold (e.g. Alves et al. 2012a; Alves and Pereira Filho 2007; Bezerra et al. 2013; Fernandes-Ferreira et al. 2012; Léo Neto et al. 2012). Alves et al. (2012b), when investigating animals used for spiritual and religious purposes by followers of the Afro-Brazilian religion Candomblé, identified the most cited animals in the terreiros, including *Monetaria caputserpentis*, *Monetaria moneta*, *Monetaria annulus*, *Gallus gallus*, *Columba livia*, *Chelonoidis denticulata*, *Numida meleagris*, and *Capra hircus*. Some of these species are domesticated and even brought from other continents and are also raised for human consumption. However, some of the species used for religious purposes are only found in the wild. The authors also found an extensive commercial network involving collectors, middlemen/distributors, store owners, and consumers focused on the commercialization of animal products. Traders cited *Hippocampus reidi*, *Boa constrictor*, *Sotalia fluviatilis*, and *Crotalus durissus* as the most traded species. Product use must be identified by spiritual guides, with information provided by spiritual entities. Regarding the use of live animals, these are sacrificed and offered to the orixás in the terreiros.

In the mystical use of animal resources, some species are employed in crafting handmade objects. For instance, Bonifácio et al. (2016) describe the use of the horn and paws of *Subulo gouazoubira* in the form of keychains and paws kept in homes, respectively, serving as a protective amulet or to expel bad moods. These peculiarities are fundamental characteristics to be analyzed in ethnozoological studies, as conservation and management strategies must integrate the cultural and spiritual values associated with the species.

Conflict situations with animals are also reported by some studies (e.g. Carvalho et al. 2019; Costa et al. 2021; Diniz 2019; Silva et al. 2020a; Fernandes-Ferreira et al. 2012; Lima et al. 2016; Lima et al. 2020; Saiki et al. 2009). Lima et al. (2020) found that motivations for people to kill species considered conflicting, such as *Boa constrictor*, *Panthera onca*, and *Rupornis magnirostris*, include a natural aversion to wildlife, reduction and prevention attacks on livestock, and upon others domestic animals, respectively. Other significant reasons include viewing some animals as disease carriers, such as *Cerdocyon thous*

(Silva et al. 2020b), and causing economic losses to people. Carvalho et al. (2019) reported economic losses triggered by wild animals, mainly by *Psittacara leucophthalmus*, which attacked corn and fruit crops. Lima et al. (2016) recorded that the species *Planococcus* sp., *Aphis gossypii*, *Bemisia tabaci*, and *Diabrotica speciosa* are insects considered pests of okra. It is necessary to outline control strategies for the management of pest species.

The use craft was reported in some research (e.g., Alves et al. 2012a; Blásquez et al. 2009; Mariz et al. 2020; Mendonça et al. 2014; Santos et al. 2019b; Silva et al. 2020a). Santos et al. (2019b) identified that species from the carnivore group (*Cerdocyon thous*, *Puma yagouaroundi*, *Leopardus pardalis*, *Puma concolor*, *Galitictis cuja*, *Procyon cancrivorous*) obtained a significant number of citations for craft purposes, with leather commonly cited as a material used in making musical pieces, such as tambourines and drums, motorcycle seats, chair covers, car seats, shoes and boot soles, bags, hats, keychains, belts, among others. The authors also discuss the use of *Kerodon rupestrus* stomach in cheese production, something also observed by Alves et al. (2012a). Mariz et al. (2020) highlight the capture of fish and crustaceans (for example, *Panulirus* spp., *Carpilius corallinus* e *Ucides cordatus*) intended to produce handicrafts. The species with the highest rates of capture were *Balistes vetula* and *Megalops atlanticus*, with the most cited materials being the whole body or taxidermized body parts and skin, used to make jewelry and souvenirs.

For veterinary use, medicines are crafted using various products and by-products, including bile, bone, fat, feathers, gizzards, heads, homemade butter, honey, horns, leather, meat, milk, rattles (from rattlesnakes), saburá (fermented pollen), skulls, stingers, tails, urine, and viscera, or the live animal. These are intended for the treatment of inflammatory and dermatological diseases or conditions (e.g., respiratory problems, rheumatism, throat problems, furunculosis, dermal nodules, wounds, and swelling), as well as obstetric disorders (prolapse of the uterus) (e.g., Barboza et al. 2007; Confessor et al. 2009; Souto et al. 2011a; Souto et al. 2011b; Souto et al. 2012a; Souto et al. 2012b). Souto et al. (2011b) identified that these practices have a strong relationship with traditional human medicine, presenting converging information on species used, prevalence of use, modes of administration, and techniques applied.

The bioindication category, the species are described as foreshadowing events, such as good or bad news, indicative of bad luck or death, and climate predictions (e.g., Costa-Neto 2006; Galvagne-Loss and Costa-Neto 2017; Melo et al. 2014). In the case of birds, their vocalization and body parts, such as feathers, respectively, aid in classifying the predicted event

and protecting against unpleasant events (e.g., Bezerra *et al.* 2013). Furthermore, the authors recorded that, in some cases, people believe that keeping certain birds as pets helps to avoid unpleasant events, as they can predict such occurrences, as is the case with *Cyanocorax cyanopogon*.

These articles contain valuable information about people's relationships with wild animals. While these practices are considered important for populations, they can result in irreparable damage to fauna. Therefore, they must be scrutinized to identify species requiring attention regarding the intensification of studies, aiming to provide data focused on the sustainability of these resources. It is also crucial for these studies to engage with other scientific areas, such as political ecology, to develop plausible alternatives for addressing emerging socio-environmental problems, promoting actions aimed at building a sustainable future. In this context, the creation of public policies can be crucial tools in preventing environmental damage, considering the importance of fauna for people.

## Taxonomic Bias

Most studies focus on the vertebrate group, which includes animals of large or medium size such as mammals, birds, and fish. In aquatic systems, fish are the most addressed. This pattern can be explained mainly by people's dietary dependence on the resource, but it can also be related to researchers' interest in these groups.

Classes like Asterozoa, Echinozoa, Hydrozoa, and Chilopoda fall below the average when correlated with the mean of developed studies and uses. This is due to the limited number of studies conducted with animals from these taxa. Consequently, further evaluations are imperative to gain a more nuanced understanding of this indicator, employing a specific approach tailored to each zoological theme.

The results of the Kruskal-Wallis test for the average number of studies per zoological class can be attributed to the greater concentration of studies involving rural farming communities, a social group with no direct contact with certain organisms. However, even other populations involved mention these species with low frequency.

The classes with the largest number of included species were Actinopteri, Chondrichthyes, and the Aves group. In addition to ichthyofauna and avifauna, the mammalian group and reptiles are described as animals with great utilitarian importance. The use of these animals may be associated with their geographic distribution and population density, as noted by Santos *et al.* (2019c) in their investigation, where the authors found that the perceived abundance of mammal fauna influenced the capture of the collected animals

by the studied community.

Mendonça *et al.* (2016) also observed this dynamic, emphasizing that in areas where larger animals are absent, people intensify the collection of species with a greater perceived spatial distribution. The flavor of the species is another factor influencing people's choices, as well as low income and education, contributing to animal slaughter and consumption. On the other hand, seasonality did not predict consumption intensity.

The results for the Total uses by zoological class reflects the low diversity of use described for the species present in the taxa (asteroidea, chilopoda, echinoidea, and Hydrozoa), a favorable situation for these animals since the versatility of use, together with their biological conditions and environmental factors, can favor their population decline.

## Main thematic approaches

The high representation of articles focusing on specialized Ethnozoology (61%) can be attributed to several factors. Firstly, there is an evaluative focus in research, exemplified by studies such as Sodr  Neto *et al.* (2022), which assesses local customs and practices specifically related to birdlife. Second, a focus on a particular zoological taxon, as seen in Silva *et al.* (2020b), who directs their investigation to the ecology, interactions, and human perceptions of *Cerdocyon thous* in rural landscapes. Lastly, studies addressing a specific category of use, such as the research by Santos *et al.* (2020), which identifies conflicts between humans and wildlife.

Specific approaches investigated revealed a lower number of studies that delved into the peculiarities of management practices. For instance, Costa-Neto *et al.* (2002) explored *in situ* management of fish and crustaceans for capturing specimens, while Soares *et al.* (2020) described *ex situ* maintenance practices involved in breeding wild birds kept in captivity for commercialization, fights, or appreciation of bird songs. Oliveira *et al.* (2018) highlighted care practices for maintaining birds, including providing food, medicines, and, in some cases, training to improve singing or learn songs from other species. Santos *et al.* (2021) emphasized the increasing importance of this approach for other zoological groups, potentially leading to the incipient process of animal domestication.

The Medicinal approach also demonstrated significant relevance, encompassing research on lists of all species known and used by populations (20%) and specific zoological groups (5%), totaling 21 articles (25%). Example include studies by Alves *et al.* (2006; 2007; 2007a; 2007b; 2008; 2009; 2010; 2016b), Alves and Santana (2008); Andrade and Costa-Neto (2005), Bar-

ros *et al.* (2012), Brito *et al.* (2019), Costa-Neto (1999; 2011), Costa-Neto and Oliveira (2000), Costa-Neto and Resende (2004), Ferreira *et al.* (2009), Nascimento *et al.* (2016) Oliveira *et al.* (2010), among others. These findings underscore the importance of zotherapy in various sociocultural environments.

Despite the wealth of information on medicinal use, Alves *et al.* (2021) recently evaluated zotherapy information from a new hypothetical and methodological perspective. This research sought to understand the influence of ecological and evolutionary factors on the medicinal use of wild mammals, revealing that phylogenetically close species are used to treat similar pathologies. This emphasizes the need for new perspectives to comprehend the relationship between people and animals, considering cultural dynamism and the specific biological and ecological characteristics of each species.

Hunting, as an approach, represented 13% of the analyzed manuscripts (e.g., Alves *et al.* 2012c; Barbosa *et al.* 2018; Barbosa *et al.* 2022; Barboza *et al.* 2016; Fernandes-Ferreira *et al.* 2013; Marques *et al.* 2022; Mendonça *et al.* 2016; Oliveira *et al.* 2017; Souto *et al.* 2019; Souza and Alves 2014; Souza *et al.* 2022). Despite its illegality in Brazil, the studies indicate its prevalence in various municipalities, with records of commercialization of products and by-products derived from hunted species. The activity is often associated with purposes such as medicinal use, crafting, and breeding as pets. Fernandes-Ferreira and Alves (2017), in their analyses on the topic, emphasize the necessity to expand studies concerning the complexity of hunting, including factors influencing the consumption of animals and the sustainability of the fauna involved in the practice.

On the other hand, the General Ethnozoology approach had the lowest representation (1%), featuring only one article by Torres *et al.* (2009). Their study aimed to analyze aspects of the interaction between human populations and biodiversity in the Genipabu Environmental Protection Area (APA-Genipabu) on the coast of the state of Rio Grande do Norte. This scarcity of research highlights the limited number of studies with a broad focus on understanding all zoological groups considered important by people in a particular location. Expanding studies to all groups requires greater involvement among regional and global institutions, enabling the strengthening of collaborative research networks.

## CONCLUSION

The prevalence of ethnozoological research in Brazil has been notable, with a significant focus on data compilation in the Northeast region, particularly within the Caatinga biome. Regions like the Central-

West and South demand increased investment in researches to enhance comprehension of the dynamics between people and the fauna in these areas.

Despite the escalating number of articles produced in Brazilian territory, certain zoological classes receive limited attention, necessitating expansion. Vertebrates stand out as the zoological groups with the highest number of species spotlighted for their importance food and medicinal, with a particular emphasis on avifauna and ichthyofauna. However, herpetofauna and mammal fauna exhibit greater versatility of use, warranting more in-depth assessments capable of gauging the impact stemming from these relationships. In general terms, these studies address popular knowledge about species, without extending to investigations that involve, for example, hypothesis testing. Achieving this requires alignment between ethnozoological studies and other scientific analyses, such as ecological assessments focusing on the biology and ecology of species, as well as political ecology.

Bibliometric analyses demonstrate the need to expand collaboration among national and international research networks, providing a pathway to address the scarcity of studies in biomes such as the Cerrado and Pantanal, and consequently enhancing the understanding of species requiring conservation actions and the development of political strategies for populations that regularly interact with fauna, relying on these resources for their subsistence. Established research groups in northeastern Brazil may play a relevant role in nucleating new research nodes in other regions.

## ACKNOWLEDGMENTS

We express our gratitude to the Coordination for the Improvement of Higher Education Personnel (CAPES) for providing financial support for this research through a scholarship awarded to the first author of this manuscript. To the Postgraduate Program in Development and Environment at the Federal University of Paraíba in João Pessoa, Brazil, and the Research Group on Multidisciplinary and Socioecological Studies at the Federal University of Mato Grosso do Sul in Campo Grande, Brazil, for their invaluable intellectual and logistical support in conducting this research. Additionally, our sincere thanks extend to the Pantanal Network PPBio/CNPq.

## DATA AVAILABILITY

For access to the specific data supporting the findings of this study, interested parties can contact the corresponding author upon reasonable request.

## CONFLICT OF INTEREST

The authors affirm that no conflicts of interest exist in relation to this study.

## CONTRIBUTION STATEMENT

Conceptualization: SSS, RFPL.

Data curation: SSS.

Formal analysis: SSS, HKLS, DGN, FOR.

Investigation: SSS, HKLS, RFPL.

Methodology: SSS, HKLS, RFPL, DGN, FOR.

Visualization: SSS, HKLS, HFM, EMCN, RFPL, DGN, FOR.

Writing – original draft: SSS.

Writing – review and editing: SSS, HKLS, HFM, EMCN, RFPL, DGN, FOR.

## REFERENCES

- Abrantes PM, Souza RF, Lucena CM, Lucena RFP, Pereira DD (2011) **Avisos de chuva e de seca na memória do povo: o caso do Cariri Paraibano.** *Revista de Biologia e Farmácia - BioFar* 5: 255-264.
- Almeida MCS, Ferreira FS, Beltrão-Mendes R (2023) **Game mammals and their uses by local hunters in an atlantic forest region of Northeast Brazil.** *Revista Etnobiologia* 21(1): 31-47.
- Alencar JBR, Silva EF, Santos VM, Soares HKL, Lucena RFP, Brito CH (2012) **Percepção e uso de “insetos” em duas comunidades rurais no semiárido do estado da Paraíba.** *Revista de Biologia e Farmácia - BioFar* Volume Especial.
- Alves RRN, Alves HN (2011) **The faunal drug-store: animal-based remedies used in traditional medicines in Latin America.** *Journal of Ethnobiology and Ethnomedicine* 7(1): 43. doi: doi: 10.1186/1746-4269-7-9.
- Alves MM, Lopes SF, Alves RRN (2016a) **Wild vertebrates kept as pets in the semiarid region of Brazil.** *Tropical Conservation Science* 9(1): 354-368. doi: doi: 10.1177/19400829160090011.
- Alves RRN, Araújo BMC, Policarpo IS, Pereira HM, Borges AKM, Vieira WLS, Vasconcellos A (2019) **Keeping reptiles as pets in Brazil: Ethnobiological and conservation aspects.** *Journal for Nature Conservation* 49: 9-21. doi: doi: 10.1016/j.jnc.2019.02.002.
- Alves RRN, Borges AKM, Barboza RRD, Souto WMS, Gonçalves-Souza T, Provete DB, Albuquerque UP (2021) **A global analysis of ecological and evolutionary drivers of the use of wild mammals in traditional medicine.** *Mammal Review* 51(2): 293–306. doi: doi: 10.1111/mam.12233.
- Alves RRN, Gonçalves MBR, Vieira WLS (2012c) **Caça, uso e conservação de vertebrados no semiárido Brasileiro.** *Tropical Conservation Science* 5(3): 394-416. doi: doi: 10.1177/194008291200500312.
- Alves RRN, Leite RCL, Souto WMS, Bezerra DMM, Loures-Ribeiro (2013) **A Ethno-ornithology and conservation of wild birds in the semi-arid Caatinga of northeastern Brazil.** *Journal of Ethnobiology and Ethnomedicine* 9(14):1-12. doi: doi: 10.1186/1746-4269-9-14.
- Alves RRN, Lima HN, Tavares MC, Souto WMS, Barboza RRD, Vasconcellos A (2008) **Animal-based remedies as complementary medicines in Santa Cruz do Capibaribe, Brazil.** *BMC Complementary and Alternative Medicine* 8(44): 1-9. doi: doi: 10.1186/1472-6882-8-44.
- Alves RRN, Léo Neto NA, Brooks SE, Albuquerque UP (2009) **Commercialization of animal-derived remedies as complementary medicine in the semi-arid region of Northeastern Brazil.** *Journal of Ethnopharmacology* 124: 600–608. doi: doi: 10.1016/j.jep.2009.04.049.
- Alves RRN, Melo MF, Ferreira FS, Trovão DMBM, Dias TLP, Oliveira JV, Lucena RFP, Barboza RRD (2016b) **Healing with animals in a semiarid northeastern area of Brazil.** *Environment, Development and Sustainability* 18: 1733–1747. doi: 10.1007/s10668-015-9715-1.
- Alves RRN, Oliveira MGG, Barboza RRD, Lopez LCS, Oliveira MGG (2010) **An ethnozoological survey of medicinal animals commercialized in the markets of Campina Grande, NE Brazil.** *Human Ecology Review* 17(1): 11-17. .
- Alves RRN, Pereira Filho GA, Lima YCC (2007) **Snakes used in ethnomedicine in Northeast Brazil.** *Environment, Development and Sustainability* 9: 455–464. doi: 10.1007/s10668-006-9031-x.
- Alves RRN; Pereira Filho GA (2007) **Commercialization and use of snakes in North and Northeastern Brazil: implications for conservation and management.** *Biodiversity and Conservation* 16: 969–985. doi: 10.1007/s10531-006-9036-7.
- Alves RRN, Pereira Filho GA, Vieira KS, Souto WMS, Mendonça LET, Montenegro FGP, Almeida WO, Vieira WLS (2012a) **A zoological catalogue of hunted reptiles in the semiarid region of Brazil.** *Journal of Ethnobiology and Ethnomedicine* 8(27): 1-29. doi: 10.1186/1746-4269-8-27.

- Alves RRN, Rosa IL (2006) **From cnidarians to mammals: The use of animals as remedies in fishing communities in NE Brazil.** *Journal of Ethnopharmacology* 107:259–276. doi: [10.1016/j.jep.2006.03.007](https://doi.org/10.1016/j.jep.2006.03.007).
- Alves RRN, Rosa IL (2007a) **Zootherapeutic practices among fishing communities in North and Northeast Brazil: A comparison.** *Journal of Ethnopharmacology* 111: 82–103. doi: [10.1016/j.jep.2006.10.033](https://doi.org/10.1016/j.jep.2006.10.033).
- Alves RRN, Rosa IL (2007b) **Zootherapy goes to town: The use of animal-based remedies in urban areas of NE and N Brazil.** *Journal of Ethnopharmacology* 113: 541–555. doi: [10.1016/j.jep.2007.07.015](https://doi.org/10.1016/j.jep.2007.07.015).
- Alves RRN, Rosa IL, Léo Neto NA, Voeks R (2012b) **Animals for the gods: magical and religious faunal use and trade in Brazil.** *Human Ecology* 40: 751–80. doi: [10.1007/s10745-012-9516-1](https://doi.org/10.1007/s10745-012-9516-1).
- Alves RRN, Santana GG (2008) **Use and commercialization of *Podocnemis expansa* (Schweiger 1812) (Testudines: Podocnemididae) for medicinal purposes in two communities in North of Brazil.** *Journal of Ethnobiology and Ethnomedicine* 4(3): 1–6. doi: [10.1186/1746-4269-4-3](https://doi.org/10.1186/1746-4269-4-3).
- Alves RRN, Souto WMS (2011) **Ethnozoology in Brazil: Current status and perspectives.** *Journal of Ethnobiology and Ethnomedicine* 7(22): 1–18. doi: [10.1186/1746-4269-7-22](https://doi.org/10.1186/1746-4269-7-22).
- Alves RRN, Souto WMS (2015) **Ethnozoology: A Brief Introduction.** *Ethnobiology and Conservation* 4: 1–13. doi: [10.15451/ec2015-1-4.1-1-13](https://doi.org/10.15451/ec2015-1-4.1-1-13).
- Andrade JN, Costa-Neto EM (2005) **Primeiro registro da utilização medicinal de recursos pesqueiros na cidade de São Félix, Estado da Bahia, Brasil.** *Acta Scientiarum Biological Sciences* 27(2): 177–183. doi: [10.4025/actasci-biolsci.v27i2.1328](https://doi.org/10.4025/actasci-biolsci.v27i2.1328).
- Barbosa JAA, Aguiar JO (2015) **Conhecimentos e usos da fauna por caçadores no semiárido brasileiro: um estudo de caso no estado da Paraíba, Nordeste do Brasil.** *Biotemas* 28(2): 137–48. doi: [10.5007/2175-7925.2015v28n2p137](https://doi.org/10.5007/2175-7925.2015v28n2p137).
- Barbosa JAA, Aguiar JO, Alves RRN (2018) **Medicinal use of animals by hunters in North eastern Brazil.** *Indian Journal of Traditional Knowledge* 17(3): 485–493. .
- Barbosa JAA, Aguiar JO, Alves RRN (2022) **Hunting and wildlife use in protected areas of the Atlantic rainforest, northeastern Brazil.** *Desenvolvimento e Meio Ambiente* 60: 249–270. doi: [10.5380/dma.v60i0.74388](https://doi.org/10.5380/dma.v60i0.74388) e-ISSN 2176-9109.
- Barboza RD, Lopes SF, Souto WMS, Fernandes-Ferreira H, Alves RRN (2016) **The role of game mammals as bushmeat In the Caatinga, north-east Brazil.** *Ecology and Society* 21(2): 2. doi: [10.5751/ES-08358-210202](https://doi.org/10.5751/ES-08358-210202).
- Barboza RRD, Souto WMS, Mourão JS (2007) **The use of zootherapeutics in folk veterinary medicine in the district of Cubati, Paraíba State, Brazil.** *Journal of Ethnobiology and Ethnomedicine* 3(32):1–12. doi: [10.1186/1746-4269-3-32](https://doi.org/10.1186/1746-4269-3-32).
- Barros FB, Azevedo PA (2014) **Common opossum (*Didelphis marsupialis* Linnaeus, 1758): food and medicine for people in the Amazon.** *Journal of Ethnobiology and Ethnomedicine* 10(65): 1–14. doi: [10.1186/1746-4269-10-65](https://doi.org/10.1186/1746-4269-10-65).
- Barros FB, Varela SAM, Pereira HM, Vicente L (2012) **Medicinal use of fauna by a traditional community in the Brazilian Amazonia.** *Journal of Ethnobiology and Ethnomedicine* 8(37): 1–20. doi: [10.1186/1746-4269-8-37](https://doi.org/10.1186/1746-4269-8-37).
- Bezerra DMM, Araújo HFP, Alves AGC, Alves RRN (2013) **Birds and people in semiarid northeastern Brazil: symbolic and medicinal relationships.** *Journal of Ethnobiology and Ethnomedicine* 9(3): 1–11. doi: [10.1186/1746-4269-9-3](https://doi.org/10.1186/1746-4269-9-3).
- Bezerra DMM, Araújo HFP, Alves RRN (2017) **Keeping wild birds as pets in a semiarid region of Rio Grande do Norte State, Northeastern Brazil.** *Hornero* 32(1): 85–93. doi: [10.56178/eh.v32i1.543](https://doi.org/10.56178/eh.v32i1.543).
- Bezerra DMM, Araújo HFP, Alves RRN (2020) **Understanding the use of wild birds in a priority conservation area of Caatinga, a Brazilian tropical dry forest.** *Environment, Development and Sustainability* 22: 5297–5316. doi: [10.1007/s10668-019-00425-1](https://doi.org/10.1007/s10668-019-00425-1).
- Blásquez JR, Costa-Neto EM, Landero-Torres I (2009) **Comparison of edible bee species from Sieras of Jibóia (Bahia, Brazil) and Zongolica, (Veracruz, Mexico).** *Revista Colombiana de Entomología* 35(2): 217–223. doi: [10.25100/so-colen.v35i2.9222](https://doi.org/10.25100/so-colen.v35i2.9222).
- Bonfim AF, Andrea MV, Almeida R, Oliveira KN, Lemos MS, Aniceto ES (2021) **Saber sobre pássaros: Um entendimento etnoornitológico dos moradores do povoado de Catunida estrada, município de Jaguarari, no Sertão Baiano.** *Revista Ouricuri* 11(1): 029–050. doi: [10.29327/ouricuri.11.1-4](https://doi.org/10.29327/ouricuri.11.1-4).
- Bonifácio KM, Freire EMX, Schiavetti A (2016) **Cul-**



tural keystone species of faunaasa method for assessing conservation priorities in a Protected Area of the Brazilian semiarid. *Biota Neotropica* 16(2): e20140106. doi: [10.1590/1676-0611-BN-2014-0106](https://doi.org/10.1590/1676-0611-BN-2014-0106).

Braga-Pereira F, Morcatty TQ, El Bizri HR, Tavares AS, Mere-Roncal C, González-Crespo C, Bertsch C, Rodriguez CR, Bardales-Alvites C, von Mühlen EM, Bernárdez-Rodríguez GF, Paim FP, Tamayo JS, Valsecchi J, Gonçalves J, Torres-Oyarce L, Lemos LP, Vieira MAR, Bowler M, Gilmore MP, Perez NCA, Alves RR, Peres CA, Pérez-Peña P, Mayor P (2022) **Congruence of local ecological knowledge (LEK)-based methods and line-transect surveys in estimating wildlife abundance in tropical forests.** *Methods in Ecology and Evolution* 13:743-756. doi: [10.1111/2041-210X.13773](https://doi.org/10.1111/2041-210X.13773).

Brito ISPB, Borges AKM, Lopes SF, Dias TLP, Alves RRN (2019) **Environmental influence on the choice of medicinal animals: a case study from northeastern Brazil.** *Journal of Ethnobiology and Ethnomedicine* 15(55): 1-10. doi: [10.1186/s13002-019-0337-9](https://doi.org/10.1186/s13002-019-0337-9).

Campos L (2023) **Gran Chaco: conheça o bioma esquecido.** Disponível em <https://ecoa.org.br/gran-chaco-o-bioma-esquecido/> Acesso em 24-10-23.

Carvalho ALC, Araújo AR, Machado TMM, Ribon R, Lopes LEB (2019) **Wildlife and damage to agriculture: na ethnobiological approach with rural producers in southeastern Brazil.** *Revista Brasileira de Ornitologia* 27(1): 17-26. doi: [10.1007/BF03544442](https://doi.org/10.1007/BF03544442).

Confessor MVA, Mendonça LET, Mourão JS, Alves RRN (2009) **Animals to heal animals: ethnoveterinary practices in semiarid region, Northeastern Brazil.** *Journal of Ethnobiology and Ethnomedicine* 5(37): 1-9. doi: [10.1186/1746-4269-5-37](https://doi.org/10.1186/1746-4269-5-37).

Costa MKB, Alves RRN, Navoni JA, Freire EMX (2021) **Ethnozoology of snakebite victims in a risk area in Northeast Brazil.** *Toxicon* 201: 155-163. doi: [10.1016/j.toxicon.2021.08.021](https://doi.org/10.1016/j.toxicon.2021.08.021).

Costa-Neto EM (1999) **Healing with animals in Feira de Santana City, Bahia, Brazil.** *Journal of Ethnopharmacology* 65: 225-230. doi: [10.1016/S0378-8741\(98\)00158-5](https://doi.org/10.1016/S0378-8741(98)00158-5).

Costa-Neto EM (2006) **“Cricket singing means rain”: semiotic meaning of insects in the district of Pedra Branca, Bahia State, Northeastern Brazil.** *Anais da Academia Brasileira de Ciências* 78(1): 59-68. doi: [10.1590/S0001-37652006000100007](https://doi.org/10.1590/S0001-37652006000100007).

Costa-Neto EM (2011) **A zooterapia popular no Estado da Bahia: registro de novas espécies animais utilizadas como recursos medicinais.** *Ciências & Saúde Coletiva* 16(Supl. 1): 1639-1650. doi: [10.1590/S1413-81232011000700100](https://doi.org/10.1590/S1413-81232011000700100).

Costa-Neto EM, Dias CV, Melo MN (2002) **O conhecimento ictiológico tradicional dos pescadores da Cidade de Barra, Região do Médio São Francisco, Estado da Bahia, Brasil.** *Acta Scientiarum* 24 (2): 561-572. doi: [10.4025/actasciobiolsci.v24i0.2360](https://doi.org/10.4025/actasciobiolsci.v24i0.2360).

Costa-Neto EM, Oliveira MVM (2000) **Cockroach is Good for Asthma: Zootherapeutic Practices in Northeastern Brazil.** *Research in Human Ecology* 7(2): 41-51. .

Costa-Neto EM, Resende JJ (2004) **A percepção de animais como “insetos” e sua utilização como recursos medicinais na cidade de Feira de Santana, Estado da Bahia, Brasil.** *Acta Scientiarum. Biological Sciences* 26(2): 143-149. doi: [10.4025/actasciobiolsci.v26i2.1612](https://doi.org/10.4025/actasciobiolsci.v26i2.1612).

Costa-Neto EM (2013) **Análise Etnossemântica de nomes comuns de abelhas e vespas (insecta, hymenoptera) na terra indígena pankararé, bahia, brasil.** *Cadernos de Linguagem e Sociodidática* 14 (1): 237-251. doi: [10.26512/les.v14i1.22248](https://doi.org/10.26512/les.v14i1.22248).

Corrêa CSL, Tavares RL, Calvão LB, Oliveira Junior JMB (2022). **Estudos da etnozoológia no Brasil: uma análise cienciométrica de 1967 a 2017.** In: Oliveira-Junior, JMB (Orgs.) **Zoologia: panorama atual e desafios futuros.** Ponta Grossa - PR: Atena, 2022. doi: [10.22533/at.ed.4972227051](https://doi.org/10.22533/at.ed.4972227051) .

Diniz RF (2019) **Etnosaberes e culturas tradicionais afrobrasileiras: farmacopeia, magia e reprodução material e simbólica de comunidades quilombolas do Vale do Jequitinhonha-MG.** *Geographia* 21(47). doi: [10.22409/GEOgraphia2019.v21i47.a28178](https://doi.org/10.22409/GEOgraphia2019.v21i47.a28178).

Duarte DF, Fernandes TAS, Waldige AA, Silva TS, Santos BRGD, Ferreira TF, Alves JN, Cruz da Silva AY, Scherer A (2021). **Wild animals trafficking and its impacts on the environment.** .

Ferreira FS, Brito SV, Ribeiro SC, Almeida WO, Alves RRN (2009) **Zootherapeutics utilized by residents of the community Poço Dantas, Crato-CE, Brazil.** *Journal of Ethnobiology and Ethnomedicine* 5(21): 1-10. doi: [10.1186/1746-4269-5-21](https://doi.org/10.1186/1746-4269-5-21).

Fernandes-Ferreira H, Alves RRN (2017) **The researches on the hunting in Brazil: a brief overview.** *Ethnobiology and Conservation* 6: 1-6. doi: [10.15451/ec2017-07-6.6-1-6](https://doi.org/10.15451/ec2017-07-6.6-1-6).

- Fernandes-Ferreira H, Mendonça SV, Albano C, Ferreira FS, Alves RRN (2012) **Hunting, use and conservation of birds in Northeast Brazil.** *Biodiversity and Conservation* 21: 221–244. doi: [10.1007/s10531-011-0179-9](https://doi.org/10.1007/s10531-011-0179-9).
- Fernandes-Ferreira H, Mendonça SV, Cruz RL, Borges-Nojasa DM, Alves RRN (2013) **Hunting of Herpetofauna in Montane, Coastal, and dry-land areas of north Eastern Brazil.** *Herpetological Conservation and Biology* 8(3): 652–666.
- Galvagne-Loss AT, Costa Neto EM (2017) **Ecología de las aves de acuerdo a los habitantes del poblado de Pedra Branca, Santa Teresinha (Bahía. Noreste de Brasil).** *Hornero* 32(1): 73–84.
- Gama CS (2014) **Confirmation of sexual dimorphism in *Hoplias aimara* (Valenciennes, 1847) (Erythrinidae: Characiformes) proposed by local expertise in Amapá, Brazil.** *Brazilian Journal of Biology* 74 (3): 687–690. doi: [10.1590/bjb.2014.0076](https://doi.org/10.1590/bjb.2014.0076).
- Landim AS, Souza JM, Santos LB, Lins-Neto EMF, Silva DT, Ferreira FS (2023) **Food taboos and animal conservation: a systematic review on how cultural expressions influence interaction with wildlife species.** *Journal of Ethnobiology and Ethnomedicine* 19(31): 1–14. doi: [10.1186/s13002-023-00600-9](https://doi.org/10.1186/s13002-023-00600-9).
- Léo Neto NA, Voeks RA, Dias TLP, Alves RRN (2012) **Mollusks of Candomblé: symbolic and ritualistic importance.** *Journal of Ethnobiology and Ethnomedicine* 8(10): 1–10. doi: [10.1186/1746-4269-8-10](https://doi.org/10.1186/1746-4269-8-10).
- Licarião MR, Bezerra DMM, Alves RRN (2013) **Wild birds as pets in Campina Grande, Paraíba State, Brazil: An Ethnozoological Approach.** *Anais da Academia Brasileira de Ciências* 85(1): 201–213. doi: [10.1590/S0001-37652013000100011](https://doi.org/10.1590/S0001-37652013000100011).
- Lima DCO, Ramos MA, Silva HCH, Alves AGC (2016) **Rapid assessment of insect fauna based on local knowledge: comparing ecological and ethnozoological methods.** *Journal of Ethnobiology and Ethnomedicine* 12(15):1–8. doi: [10.1186/s13002-016-0085-z](https://doi.org/10.1186/s13002-016-0085-z).
- Lima NS, Napiwoski SJ, Oliveira MA (2020) **Human-wildlife conflict in the Southwestern Amazon: poaching and its motivations.** *Nature Conservation Research* 5(1): 109–114. doi: [10.24189/ncr.2020.006](https://doi.org/10.24189/ncr.2020.006).
- Loss ATG, Costa-Neto EM, Flores FM (2013) **Ornitoáugures no Povoado de Pedra Branca, Santa Teresinha, Estado da Bahia, Nordeste do Brasil.** *Revista Etnobiología* 11( 3): 201–213.
- Magalhães C, Barbosa UC, Py-Daniel V (2006) **Decapod crustaceans used as food by the Yanomami Indians of the Balawa-ú village, State of Amazonas, Brazil.** *Acta Amazonica* 36(3): 369 – 374. doi: [10.1590/S0044-59672006000300013](https://doi.org/10.1590/S0044-59672006000300013).
- Mariz D, Souza ACFF, Teixeira SF, Campos SS, Lucena RFP, Alves RRN (2020) **Knowledge on the use of catch material for craftwork/handicrafts by an urban fishing community.** *Indian Journal of Traditional Knowledge* 19(4): 902–909. doi: [10.56042/ijtk.v19i4.25551](https://doi.org/10.56042/ijtk.v19i4.25551).
- Margalida A, Campión D, Donázar J A (2014) **Vultures vs livestock: Conservation relationships in an emerging conflict between humans and wildlife.** *Oryx* 48(2): 172–176. doi: [10.1017/S0030605312000889](https://doi.org/10.1017/S0030605312000889).
- Marques AK, Novato TS, Albuquerque UP, Soldati GT (2022) **Can Socioeconomic Variables Influence Bird Hunting Activity in the Brazil's Semi- Arid Region?.** *Human Ecology* 50: 515–530. doi: [10.1007/s10745-022-00330-8](https://doi.org/10.1007/s10745-022-00330-8).
- Melo RS, Silva OC, Souto A, Alves RRN, Schie N (2014) **The role of mammals in local communities living in conservation areas in the Northeast of Brazil: an ethnozoological approach.** *Tropical Conservation Science* 7(3): 423–439. doi: [10.1177/194008291400700305](https://doi.org/10.1177/194008291400700305).
- Mendonça LET, Vasconcellos A, Souto CM, Oliveira TPR, Alves RRN (2016) **Bushmeat consumption and its implications for wildlife conservation in the semi-arid region of Brazil.** *Regional Environmental Change* 16: 1–9. doi: [10.1007/s10113-015-0901-3](https://doi.org/10.1007/s10113-015-0901-3).
- Mendonça LET, Vieira WLS, ALVES RRN (2014) **Caatinga Ethnoherpetology: Relationships between herpetofauna and people in a semiarid region of northeastern Brazil.** *Amphibian & Reptile Conservation* 8(1): 24–32.
- Ministério do Meio Ambiente - MMA (2023a) Disponível em <https://antigo.mma.gov.br/biomas/cerrado.html> Acesso em 25-07-23.
- Ministério do Meio Ambiente - MMA (2023b) Disponível em <https://www.gov.br/mma/pt-br/as-suntos/ecossistemas-1/biomas/pantanal> Acesso em 05-10-23.
- Modro AFH, Souza S, Aburaya FH, Maia E (2009) **Conhecimento dos moradores do médio Araguaia, Estado do Mato Grosso, sobre a utilização de produtos de abelhas (Hymenoptera, Apidae).** *Acta Scientiarum. Biological Sciences* 31(4):

421-424. doi: [10.4025/actasciobiolsci.v31i4.4518](https://doi.org/10.4025/actasciobiolsci.v31i4.4518).

Montenegro IF, Alencar JBR, Silva EF, Lucena RFP, Brito CH (2014) **Conhecimento, percepção e uso de animais categorizados como “insetos” em uma comunidade rural no semiárido do estado da Paraíba, Nordeste do Brasil.** *Gaia Scientia* Volume Especial - Populações tradicionais.

Morado CN, Andrade-Tubino MF, Araújo FG (2021) **Local ecological knowledge indicates: There is another breeding period in the summer for the mullet *Mugil lizaina* Brazilian tropical bay.** *Ocean and Coastal Management* 105569. doi: [10.1016/j.ocecoaman.2021.105569](https://doi.org/10.1016/j.ocecoaman.2021.105569).

Moura FPB, Marques JGW (2008) **Folk medicine using animals in the Chapada Diamantina: incidental medicine?** *Ciência Saúde Coletiva* 13(2): 2179-2188. doi: [10.1590/S1413-81232008000900023](https://doi.org/10.1590/S1413-81232008000900023).

Nascimento ALB, Lozano A, Melo JG, Alves RRN, Albuquerque UP (2016) **Functional aspects of the use of plants and animals in local medical systems and their implications for resilience.** *Journal of Ethnopharmacology* 194: 348-357. doi: [10.1016/j.jep.2016.08.017](https://doi.org/10.1016/j.jep.2016.08.017).

Oliveira WSL, Borges AKM, Lopes SF, Vasconcellos A, Alves RRN (2020) **Illegal trade of songbirds: an analysis of the activity in an area of Northeast Brazil.** *Journal of Ethnobiology and Ethnomedicine* 16: 16. doi: [10.1186/s13002-020-00365-5](https://doi.org/10.1186/s13002-020-00365-5).

Oliveira AMV, Lopes WH (2021) **Trends in scientific publication on ethnozoology: Brazil's highlight in international science.** *Revista Ibero Americana de Ciências Ambientais* 12(1): 691-698. doi: [10.6008/CBPC2179-6858.2021.001.0055](https://doi.org/10.6008/CBPC2179-6858.2021.001.0055).

Oliveira WSL, Lopes SF, Alves RRN (2018) **Understanding the motivations for keeping wild birds in the semi-arid region of Brazil.** *Journal of Ethnobiology and Ethnomedicine* 14(41): 1-14. doi: [10.1186/s13002-018-0243-6](https://doi.org/10.1186/s13002-018-0243-6).

Oliveira WSL, Luna MSO, Souto WMS, Alves RRN (2017) **Interactions between people and game mammals in a Brazilian semi-arid area.** *Indian Journal of Traditional Knowledge* 16(2): 221-228. .

Oliveira ES, Torres DF, Brooks SE, Alves RRN (2010) **The medicinal animal markets in the metropolitan region of Natal City, northeastern Brazil.** *Journal of Ethnopharmacology* 130: 54-60. doi: [10.1016/j.jep.2010.04.010](https://doi.org/10.1016/j.jep.2010.04.010).

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-

Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D (2021) **The PRISMA 2020 statement: An updated guideline for reporting systematic reviews.** *International Journal of Surgery* 29(71): 372. doi: [10.1136/bmj.n71](https://doi.org/10.1136/bmj.n71).

Pinto MF, Mourão JS, Alves RRN (2013) **Ethnotaxonomical considerations and usage of ichthyofauna in a fishing community in Ceará State, Northeast Brazil.** *Journal of Ethnobiology and Ethnomedicine* 9(17): 1-11. doi: [10.1186/1746-4269-9-17](https://doi.org/10.1186/1746-4269-9-17).

Pinto MF, Mourão JS, Alves RRN (2017) **Animal source foods consumed in two fishing communities on the northeast coast of Brazil.** *Environment, Development and Sustainability* 19: 679-692. doi: [10.1007/s10668-016-9758-y](https://doi.org/10.1007/s10668-016-9758-y).

Pozzi CM, Ladio AH (2023) **Variation of local zoological knowledge about Southern river otter and other semi-aquatic mammals in Nahuel Huapi National Park (Argentina).** *Journal of Ethnobiology and Ethnomedicine* 19(15): 1-12. doi: [10.1186/s13002-023-00590-8](https://doi.org/10.1186/s13002-023-00590-8) .

*R Core Team R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.* 2021. <https://www.Rproject.org>.

Ritter MR, Silva TC, Araújo EL, Albuquerque UP (2015) **Bibliometric analysis of ethnobotanical research in Brazil (1988–2013).** *Acta Botanica Brasilica* 29(1): 113-119. doi: [10.1590/0102-33062014abb3524](https://doi.org/10.1590/0102-33062014abb3524).

Rocha-Mendes F, Mikich SB, Bianconi GV, Pedro WA (2005) **Mamíferos do município de Fênix, Paraná, Brasil: etnozologia e conservação.** *Revista Brasileira de Zoologia* 22(4): 991-1002. doi: [10.1590/S0101-81752005000400027](https://doi.org/10.1590/S0101-81752005000400027).

Rodríguez GF, Paim FP, Tamayo JS, Valsecchi J, Gonçalves J, Torres-Oyarce L, Lemos LP, Vieira MAR, Bowler M, Gilmore MP, Perez NCA, Alves RR, Peres CA, Pérez-Peña P, Mayor P (2021) **Congruence of local ecological knowledge (LEK)-based methods and line-transect surveys in estimating wildlife abundance in tropical forests.** *Methods in Ecology and Evolution* 13: 743-756. doi: [10.1111/2041-210X.13773](https://doi.org/10.1111/2041-210X.13773).

Saiki PTO, Guido LFE, Cunha AMO (2009) **Etnoecologia, etnotaxonomia e valoração cultural de Psittacidae em distritos rurais do Triângulo Mineiro, Brasil.** *Revista Brasileira de Ornitologia* 17(1): 41-52.

Santos SS, Bonifácio KM, Lucena RFP (2021)

- Use and traditional management of *Galea spixii* (Wagler 1831) and *Kerodon rupestris* (Wied-Neuwied 1820) in the municipalities of Pau dos Ferros, Encanto, and Francisco Dantas (Northeastern Brazil). *Environment, Development and Sustainability*, 23: 1-17.
- Santos CAB, Alves RRN (2016) **Ethnoichthyology of the indigenous Truká people, Northeast Brazil.** *Journal of Ethnobiology and Ethnomedicine* 12(1): 1-10. doi: [10.1186/s13002-015-0076-5](https://doi.org/10.1186/s13002-015-0076-5).
- Santos JS, Teixeira JVS, Guanaes DHA, Rocha WD, Schiavetti Alexandre (2020) **Conflicts between humans and wild animals in and surrounding protected area (Bahia, Brazil): an ethnozoological approach.** *Ethnobiology and Conservation* 9(5): 1-22. doi: [10.15451/ec2020-02-9.05-1-22](https://doi.org/10.15451/ec2020-02-9.05-1-22).
- Santos CP, Braga-Pereira F, Borges AKM, Van Vliet N, Alves RRN (2022) **Consumption and Preferences for Wild and Domestic Meat in Indigenous Communities in the Brazilian Atlantic Forest.** *Ecology and Evolution* 10: 1-13. doi: [10.3389/fevo.2022.900398](https://doi.org/10.3389/fevo.2022.900398).
- Santos SS, Lucena RFP, Soares HKL, Soares VMS, Sales NS, Mendonça LET (2019c) **Use of mammals in a Semi-Arid Region of Brazil: An approach to the use value and data analysis for conservation.** *Journal of Ethnobiology and Ethnomedicine* 15(33): 1- 14. doi: [10.1186/s13002-019-0313-4](https://doi.org/10.1186/s13002-019-0313-4).
- Santos SS, Soares HKL, Soares VMS, Lucena RFP (2019a) **Conhecimento tradicional e utilização da fauna silvestre em São José da Lagoa Tapada, Paraíba, Brasil.** *Revista Etnobiología* 17 (1): 31–48.
- Santos SS, Soares HKL, Soares VMS, Lucena CM, Lucena RFP (2023) **Conhecimento de agricultores sobre aves em uma comunidade rural no Semiárido da Paraíba: uma abordagem da etnoornitologia.** *Revista Brasileira de Gestão Ambiental e Sustentabilidade* 10: 33-56. doi: [10.21438/rbgas\(2023\)102403](https://doi.org/10.21438/rbgas(2023)102403).
- Santos SS, Soares HKL, Soares VMS, Lucena RFP (2019b) **Traditional knowledge and use of mammals in a rural community in the Sertaneja depression (Paraíba State, Northeast Brazil).** *Indian Journal of Traditional Knowledge* 18 (1): 94-103.
- Santos SS, Lucena RFP, Soares HKL, Soares VMS, Sales NS, Mendonça LET (2019c) **Use of mammals in a Semi-Arid Region of Brazil: An approach to the use value and data analysis for conservation.** *Journal of Ethnobiology and Ethnomedicine* 15(33): 1- 14. doi: [10.1186/s13002-019-0313-4](https://doi.org/10.1186/s13002-019-0313-4).
- Seeger KR, Solé M, Martínez RA, Schiavetti Alexandre (2019) **Of people and toads: Local knowledge about amphibians around a protected area in the Brazilian Atlantic Forest.** *Neotropical Biology and Conservation* 14(4): 439–457. doi: [10.3897/neotropical.14.e47020](https://doi.org/10.3897/neotropical.14.e47020).
- Serrão EM, Braga TMP, Coêlho YKS, Campos DPF, Santos AA, Imbiriba LC, Zacardi DM (2019) **Traditional knowledge of fishermen of the reproductive behavior of fish in a flood lake in western Pará, Brazil.** *Sociedade e Natureza* 31: e45133. doi: [10.14393/SN-v31-2019-45133](https://doi.org/10.14393/SN-v31-2019-45133).
- Silva C, Silva LT, White BLA (2017) **Aversão a espécies de aves por moradores da zona urbana e rural do Município de Itabaiana, Sergipe, Brasil.** *Revista Etnobiología* 15(2): 5-15.
- Silva AR, Braga-Pereira F, Borges AKM, Oliveira JV, Silva MXG, Alves RRN (2023) **Bioecological representations and social characteristics of students influence their attitudes toward wild vertebrates.** *Journal of Ethnobiology and Ethnomedicine* 19(25): 1-14. doi: [10.1186/s13002-023-00593-5](https://doi.org/10.1186/s13002-023-00593-5).
- Silva JS, Nascimento ALB, Alves RRN, Albuquerque UP (2020a) **Use of game fauna by Fulni-ô people in Northeastern Brazil: implications for conservation.** *Journal of Ethnobiology and Ethnomedicine* 16: 18. doi: [10.1186/s13002-020-00367-3](https://doi.org/10.1186/s13002-020-00367-3).
- Silva LT, Souza ACFF, Silva LAM (2020b) **Ecology, interactions and human perceptions of *Cerdoyon thous* in rural landscapes in the state of Pernambuco, Brazil.** *Anais da Academia Brasileira de Ciências* 92(3): e20180890. doi: [10.1590/0001-3765202020180890](https://doi.org/10.1590/0001-3765202020180890).
- Sistema de Informações de Informações sobre a Biodiversidade Brasileira (SIBBr), Catálogo Taxonômico da Fauna do Brasil e Lista da Flora do Brasil 2020.* Acesso em: <https://ala-bie.sibbr.gov.br/ala-bie/species/183551>.
- Soares HKL, Soares VMS, Lopes SF, Lucena RFP, Barboza RRD (2020) **Rearing and trade of wild birds in a semiarid region of Brazil.** *Environment, Development and Sustainability* 22: 4323–4339. doi: [10.1007/s10668-019-00386-5](https://doi.org/10.1007/s10668-019-00386-5).
- Soares VMS, Soares HKL, Santos SS, Lucena RFP (2018) **Local knowledge, use, and conservation of wild birds in the semi-arid region of Paraíba state, northeastern Brazil.** *Journal of Ethnobiology and Ethnomedicine* 14–77. doi: [10.1186/s13002-018-0276-x](https://doi.org/10.1186/s13002-018-0276-x).
- Sodré Neto AI, Fraga RE, Schiavetti A (2022) **Tradition and trade: culture and exploitation of avian fauna by a rural community surrounding protected areas in**

thesouthofBahia's State, Northeastern Brazil. *Journal of Ethnobiology and Ethnomedicine* 18(12): 1-14. doi: [10.1186/s13002-022-00515-x](https://doi.org/10.1186/s13002-022-00515-x).

Sollund R (2013). Animal trafficking and trade: Abuse and species injustice. In *Emerging issues in green criminology: exploring power, justice and harm* 72-92. London: Palgrave Macmillan UK.

Souto WMS, Barboza RRD, Rocha MSP, Alves RRN, Mourão JS (2012a) **Animal-based medicines used in ethnoveterinary practices in the semi-arid region of Northeastern Brazil.** *Anais da Academia Brasileira de Ciências* 84(3): 669-678. doi: [10.1590/S0001-37652012005000038](https://doi.org/10.1590/S0001-37652012005000038).

Souto WMS, Barboza RRD, Mourão JS, Alves RRN (2012b) **Traditional knowledge of sertanejos about Zootherapeutic practices used in ethnoveterinary medicine of NE Brazil.** *Indian Journal of Traditional Knowledge* 11(2): 259-265. .

Souto WMS, Lima RN, Sousa BFCF (2019) **Illegal bushmeat hunting and trade dynamics in a major road-hub region of the Brazilian Mid North.** *Indian Journal of Traditional Knowledge* 18(2): 402-411. .

Souto WMS, Mourão JS, Barboza RRD, Mendonça LET, Lucena RFP, Confessor MVA, Vieira WLS, Montenegro PFGP, Lopez LCS, Alves RRN (2011a) **Medicinal animals used in ethnoveterinary practices of the 'Cariri Paraibano', NE Brazil.** *Journal of Ethnobiology and Ethnomedicine* 7(30): 1-20. doi: [10.1186/1746-4269-7-30](https://doi.org/10.1186/1746-4269-7-30).

Souto WMS, Mourão JS, Barboza RRD, Alves RRN (2011b) **Parallels between zootherapeutic practices in ethnoveterinary and human complementary medicine in northeastern Brazil.** *Journal of Ethnopharmacology* 134: 753-767. doi: [10.1016/j.jep.2011.01.041](https://doi.org/10.1016/j.jep.2011.01.041).

Souto WMS, Torres MAR, Sousa BFCF, Lima KGGC, Vieira LTS, Pereira GA, Guzzi A, Silva MV, Pralon BGN (2017) **Singing for Cages: The Use and Trade of Passeriformes as Wild Pets in an Economic Center of the Amazon—NE Brazil Route.** *Tropical Conservation Science* 10: 1-19. doi: [10.1177/1940082917689898](https://doi.org/10.1177/1940082917689898).

Souza JB, Alves RRN (2014) **Hunting and wildlife use in an Atlantic Forest remnant of northeast-**

**ern Brazil.** *Tropical Conservation Science* 7(1): 145-160. doi: [10.1177/194008291400700105](https://doi.org/10.1177/194008291400700105).

Souza JM, Lins Neto EMF, Ferreira FS (2022) **Influence of the sociodemographic profile of hunters on the knowledge and use of faunistic resources.** *Journal of Ethnobiology and Ethnomedicine* 18(38): 1-13. doi: [10.1186/s13002-022-00538-4](https://doi.org/10.1186/s13002-022-00538-4).

Teixeira PHR, Thel TN, Ferreira JMR, Azevedo Jr SM, Telino Junior WR, Lyra-Neves RM (2014) **Local knowledge and exploitation of the avian fauna by a rural community in the semi-arid zone of northeastern Brazil.** *Journal of Ethnobiology and Ethnomedicine* 10(81): 1-10. doi: [10.1186/1746-4269-10-81](https://doi.org/10.1186/1746-4269-10-81).

Tito MCPS, Giraldin O (2021) **Etnozoologia no Brasil central: os animais na cultura do povo In-/Javaé.** *Revista Ibero-Americana de Ciências Ambientais* 12(10): 527-544. doi: [10.6008/CBPC2179-6858.2021.010.0042](https://doi.org/10.6008/CBPC2179-6858.2021.010.0042).

Torres DF, Oliveira ES, Alves RRN, Vasconcellos A (2009) **Ethnobotany and ethnozoology in conservation areas: Use of biodiversity in the apagenipabu, Rio Grande do Norte, Brazil.** *Inter-ciência* 34(9): 624-629.

Van Eck NJ, Waltman L (2010) **Software survey: VOSviewer, a computer program for bibliometric mapping.** *Scientometrics* 84(2): 523-538. doi: [10.1007/s11192-009-0146-3](https://doi.org/10.1007/s11192-009-0146-3).

Yousefi A, Bellantonio M, Higonnet E, Hurowitz G (2018) **The Avoidable Crisis: The European Meat Industry's Environmental Catastrophe –** *Mighty Earth* 1-16.

Received: 01 June 2023

Accepted: 29 July 2023

Published: 14 August 2024

Editor: Rômulo Alves

